



AUSTRALIAN DRUG TRENDS 2026

Key Findings from the National Illicit Drug
Reporting System (IDRS) Interviews



AUSTRALIAN DRUG TRENDS 2026: KEY FINDINGS FROM THE NATIONAL ILLICIT DRUG REPORTING SYSTEM (IDRS) INTERVIEWS

Rachel Sutherland¹, Julia Uporova¹, Lily Palmer¹, Antonia Karlsson¹, Agata Chrzanowska¹, Udesha Chandrasena¹, Raimondo Bruno^{1,2}, Paul Dietze^{1,3,4}, Simon Lenton^{1,4}, Caroline Salom¹, Sophie Radke², Jack Curran³, Joanna Wilson³, Dorian Raffaele³, Matthew Cowman³, Louisa Degenhardt¹, Michael Farrell¹ & Amy Peacock^{1,2}

¹ National Drug and Alcohol Research Centre, University of New South Wales Sydney

² School of Psychology, University of Tasmania

³ Disease Elimination Program, Burnet

⁴ National Drug Research Institute and enAble Institute, Curtin University



ISSN 2981-9326 ©NDARC 2026

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. All other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to the National Drug and Alcohol Research Centre, University of New South Wales, Sydney, NSW 2052, Australia.

Suggested citation: Sutherland R, Uporova J, Palmer L, Karlsson A, Chrzanowska A, Chandrasena U, Bruno R, Dietze P, Lenton S, Salom C, Radke S, Curran J, Wilson J, Raffaele D, Cowman M, Degenhardt L, Farrell M, & Peacock A. Australian Drug Trends 2026: Key Findings from the National Illicit Drug Reporting System (IDRS) Interviews. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney; 2026. Available from: <https://doi.org/10.26190/unsworks/32706>

Please note that, as with all statistical reports, the data in this report may be subject to minor revisions over time. For the most up-to-date information, please refer to the online version at [Drug Trends](#).

This report was prepared by the National Drug and Alcohol Research Centre, UNSW Sydney. Please contact the Drug Trends team with any queries regarding this publication: drugtrends@unsw.edu.au

Table of Contents

BACKGROUND AND METHODS	15
SAMPLE CHARACTERISTICS	19
HEROIN	26
METHAMPHETAMINE	33
COCAINE	49
CANNABIS AND/OR CANNABINOID-RELATED	56
PHARMACEUTICAL OPIOIDS	68
OTHER DRUGS	83
DRUG-RELATED HARMS AND OTHER BEHAVIOURS	93

List of Tables

TABLE 1: GUIDE TO TABLE/FIGURE NOTES.....	18
TABLE 2: DEMOGRAPHIC CHARACTERISTICS OF THE SAMPLE, NATIONALLY, 2025-2026, AND BY CAPITAL CITY, 2026.....	21
TABLE 3: PAST SIX MONTH USE OF HEROIN, BY CAPITAL CITY, 2007–2026	29
TABLE 4: PAST SIX MONTH USE OF ANY METHAMPHETAMINE, BY CAPITAL CITY, 2007–2026	37
TABLE 5: PAST SIX MONTH USE OF METHAMPHETAMINE POWDER, BY CAPITAL CITY, 2007-2026	40
TABLE 6: PAST SIX MONTH USE OF METHAMPHETAMINE CRYSTAL, BY CAPITAL CITY, 2007-2026.....	41
TABLE 7: PAST SIX MONTH USE OF COCAINE, BY CAPITAL CITY, 2007-2026	52
TABLE 8: PAST SIX MONTH NON-PRESCRIBED USE OF CANNABIS AND CANNABINOID-RELATED PRODUCTS, BY CAPITAL CITY, 2007-2026	60
TABLE 9: PAST SIX MONTH NON-PRESCRIBED USE OF METHADONE, BY CAPITAL CITY, 2007-2026.....	71
TABLE 10: PAST SIX MONTH NON-PRESCRIBED USE OF BUPRENORPHINE TABLET, BY CAPITAL CITY, 2007-2026	73
TABLE 11: PAST SIX MONTH NON-PRESCRIBED USE OF BUPRENORPHINE-NALOXONE (ANY FORM), BY CAPITAL CITY, 2007-2026.....	75
TABLE 12: PAST SIX MONTH NON-PRESCRIBED USE OF MORPHINE, BY CAPITAL CITY, 2007-2026.....	77
TABLE 13: PAST SIX MONTH NON-PRESCRIBED USE OF OXYCODONE, BY CAPITAL CITY, 2007-2026	79
TABLE 14: PAST SIX MONTH NON-PRESCRIBED USE OF FENTANYL, BY CAPITAL CITY, 2018-2026.....	81
TABLE 15: PAST SIX MONTH USE OF OTHER OPIOIDS, NATIONALLY, 2019-2026	82
TABLE 16: PAST SIX MONTH USE OF NEW PSYCHOACTIVE SUBSTANCES, NATIONALLY, 2013-2026	85
TABLE 17: PAST 12-MONTH NON-FATAL OVERDOSE BY DRUG TYPE, NATIONALLY, 2025-2026, AND BY CAPITAL CITY, 2026.....	99
TABLE 18: AUDIT-C TOTAL SCORES AND PER CENT OF PARTICIPANTS SCORING ABOVE RECOMMENDED LEVELS, NATIONALLY, 2010-2026	101
TABLE 19: LIFETIME AWARENESS OF NALOXONE, AND PAST YEAR ACCESS, BY CAPITAL CITY, 2026	104
TABLE 20: INJECTING EQUIPMENT ACCESS IN PAST MONTH, NATIONALLY, 2025-2026, AND BY CAPITAL CITY, 2026.....	106
TABLE 21: INJECTING BEHAVIOURS IN THE PAST MONTH, AND LOCATION AND SITE OF LAST INJECTION, NATIONALLY, 2025-2026, AND BY CAPITAL CITY, 2026.....	108
TABLE 22: INJECTION-RELATED ISSUES IN THE PAST MONTH, NATIONALLY, 2025-2026, AND BY CAPITAL CITY, 2026.....	109
TABLE 23: ANY CURRENT DRUG TREATMENT, NATIONALLY, 2025-2026, AND BY CAPITAL CITY, 2026.....	110
TABLE 24: TOTAL OPIOID AND METHAMPHETAMINE SDS SCORES AND PER CENT OF PARTICIPANTS SCORING ABOVE CUT-OFF SCORES INDICATIVE OF DEPENDENCE, AMONG THOSE WHO REPORTED PAST SIX MONTH USE, NATIONALLY, 2015-2026.....	111
TABLE 25: HCV AND HIV TESTING AND TREATMENT, NATIONALLY, 2018-2026	113
TABLE 26: SEXUAL HEALTH BEHAVIOURS, NATIONALLY, 2022-2026.....	114
TABLE 27: NATURE OF DISABILITY, AMONG THOSE WHO SELF-IDENTIFIED AS A PERSON LIVING WITH A DISABILITY, NATIONALLY, 2026, AND BY CAPITAL CITY, 2026.....	119
TABLE 28: TYPES OF HEALTH SERVICES ACCESSED FOR ALCOHOL AND OTHER DRUG REASONS AND FOR ANY REASON IN THE PAST SIX MONTHS, NATIONALLY, 2022-2026	121
TABLE 29: PURCHASING APPROACHES IN THE PAST 12 MONTHS, NATIONALLY, 2022-2026	130

List of Figures

FIGURE 1: DRUG OF CHOICE, NATIONALLY, 2007-2026	23
FIGURE 2: DRUG INJECTED MOST OFTEN IN THE PAST MONTH, NATIONALLY, 2007-2026	24
FIGURE 3: WEEKLY OR MORE FREQUENT SUBSTANCE USE IN THE PAST SIX MONTHS, NATIONALLY, 2007-2026	25
FIGURE 4: PAST SIX MONTH USE AND FREQUENCY OF USE OF HEROIN, NATIONALLY, 2007-2026.....	28
FIGURE 5: MEDIAN PRICE OF HEROIN PER CAP, POINT AND GRAM, NATIONALLY, 2007-2026	30
FIGURE 6: CURRENT PERCEIVED PURITY OF HEROIN, NATIONALLY, 2007-2026	31
FIGURE 7: CURRENT PERCEIVED AVAILABILITY OF HEROIN, NATIONALLY, 2007-2026	32
FIGURE 8: PAST SIX MONTH USE OF ANY METHAMPHETAMINE AND OF METHAMPHETAMINE POWDER, BASE, AND CRYSTAL, NATIONALLY, 2007-2026	35
FIGURE 9: FREQUENCY OF USE OF ANY METHAMPHETAMINE AND OF METHAMPHETAMINE POWDER, BASE, AND CRYSTAL, NATIONALLY, 2007-2026	36
FIGURE 10: MEDIAN PRICE OF METHAMPHETAMINE POWDER PER POINT AND GRAM, NATIONALLY, 2007-2026.....	43
FIGURE 11: MEDIAN PRICE OF METHAMPHETAMINE CRYSTAL PER POINT AND GRAM, NATIONALLY, 2007-2026	44
FIGURE 12: CURRENT PERCEIVED PURITY OF METHAMPHETAMINE POWDER, NATIONALLY, 2007-2026	45
FIGURE 13: CURRENT PERCEIVED PURITY OF METHAMPHETAMINE CRYSTAL, NATIONALLY, 2007-2026	46
FIGURE 14: CURRENT PERCEIVED AVAILABILITY OF METHAMPHETAMINE POWDER, NATIONALLY, 2007-2026	47
FIGURE 15: CURRENT PERCEIVED AVAILABILITY OF METHAMPHETAMINE CRYSTAL, NATIONALLY, 2007-2026	48
FIGURE 16: PAST SIX MONTH USE AND FREQUENCY OF USE OF COCAINE, NATIONALLY, 2007-2026.....	51
FIGURE 17: MEDIAN PRICE OF COCAINE PER POINT AND GRAM, NATIONALLY, 2007-2026.....	53
FIGURE 18: CURRENT PERCEIVED PURITY OF COCAINE, NATIONALLY, 2007-2026	54
FIGURE 19: CURRENT PERCEIVED AVAILABILITY OF COCAINE, NATIONALLY, 2007-2026	55
FIGURE 20: PAST SIX MONTH USE AND FREQUENCY OF USE OF NON-PRESCRIBED CANNABIS AND CANNABINOID-RELATED PRODUCTS, NATIONALLY, 2007-2026	59
FIGURE 21: PAST SIX MONTH USE OF DIFFERENT FORMS OF NON-PRESCRIBED CANNABIS AND/OR CANNABINOID-RELATED PRODUCTS, AMONG THOSE WHO REPORTED RECENT NON-PRESCRIBED USE, NATIONALLY, 2018-2026	61
FIGURE 22: MEDIAN PRICE OF NON-PRESCRIBED HYDROPONIC (A) AND BUSH (B) CANNABIS PER OUNCE AND GRAM, NATIONALLY, 2007-2026.....	63
FIGURE 23: CURRENT PERCEIVED POTENCY OF NON-PRESCRIBED HYDROPONIC (A) AND BUSH (B) CANNABIS, NATIONALLY, 2007-2026	64
FIGURE 24: CURRENT PERCEIVED AVAILABILITY OF NON-PRESCRIBED HYDROPONIC (A) AND BUSH (B) CANNABIS, NATIONALLY, 2007-2026	66
FIGURE 25: PAST SIX MONTH USE (PRESCRIBED AND NON-PRESCRIBED) AND FREQUENCY OF USE OF NON-PRESCRIBED METHADONE, NATIONALLY, 2007-2026.....	70
FIGURE 26: PAST SIX MONTH USE (PRESCRIBED AND NON-PRESCRIBED) AND FREQUENCY OF USE OF NON-PRESCRIBED BUPRENORPHINE TABLET, NATIONALLY, 2007-2026	72
FIGURE 27: PAST SIX MONTH USE (PRESCRIBED AND NON-PRESCRIBED) AND FREQUENCY OF USE OF NON-PRESCRIBED BUPRENORPHINE-NALOXONE, NATIONALLY, 2007-2026	74

FIGURE 28: PAST SIX MONTH USE (PRESCRIBED AND NON-PRESCRIBED) AND FREQUENCY OF USE OF NON-PRESCRIBED MORPHINE, NATIONALLY, 2007-2026	76
FIGURE 29: PAST SIX MONTH USE (PRESCRIBED AND NON-PRESCRIBED) AND FREQUENCY OF USE OF NON-PRESCRIBED OXYCODONE, NATIONALLY, 2007-2026	78
FIGURE 30: PAST SIX MONTH USE (PRESCRIBED AND NON-PRESCRIBED) AND FREQUENCY OF USE OF NON-PRESCRIBED FENTANYL, NATIONALLY, 2013-2026.....	80
FIGURE 31: PAST SIX MONTH USE OF NON-PRESCRIBED PHARMACEUTICAL DRUGS, NATIONALLY, 2007-2026	88
FIGURE 32: PAST SIX MONTH USE OF LICIT AND OTHER DRUGS, NATIONALLY, 2007-2026	92
FIGURE 33: USE OF OPIOIDS, STIMULANTS, BENZODIAZEPINES AND CANNABIS ON THE DAY PRECEDING INTERVIEW AND MOST COMMON DRUG PATTERN PROFILES, NATIONALLY, 2026	94
FIGURE 34: PAST SIX MONTH USE OF DRUGS FOR 48 HOURS OR MORE CONTINUOUSLY WITHOUT SLEEP ('BINGE'), NATIONALLY, 2018-2026.....	95
FIGURE 35: PAST 12-MONTH ANY NON-FATAL OVERDOSE, NATIONALLY, 2007-2026	98
FIGURE 36: LIFETIME AWARENESS OF NALOXONE AND NALOXONE TAKE-HOME PROGRAMS, NATIONALLY, 2013-2026.....	104
FIGURE 37: PAST 12 MONTH OBTAINMENT OF NALOXONE, NATIONALLY, 2020-2026	104
FIGURE 38: BORROWING AND LENDING OF NEEDLES AND SHARING OF INJECTING EQUIPMENT IN THE PAST MONTH, NATIONALLY, 2007-2026	107
FIGURE 39: SELF-REPORTED MENTAL HEALTH PROBLEMS AND TREATMENT SEEKING IN THE PAST SIX MONTHS, NATIONALLY, 2007-2026	116
FIGURE 40: K10 PSYCHOLOGICAL DISTRESS SCORES, NATIONALLY, 2007-2026, AND AMONG THE GENERAL POPULATION, 2022-2023	118
FIGURE 41: HEALTH SERVICE ACCESS FOR ALCOHOL AND OTHER DRUG REASONS, AND FOR ANY REASON IN THE PAST SIX MONTHS, NATIONALLY, 2020-2026	120
FIGURE 42: SELF-REPORTED TESTING, AND DRIVING OVER THE (PERCEIVED) LEGAL LIMIT FOR ALCOHOL OR WITHIN THREE HOURS FOLLOWING ILLICIT DRUG USE, AMONG THOSE WHO HAD DRIVEN IN THE LAST SIX MONTHS, NATIONALLY, 2007-2026	123
FIGURE 43: PAST YEAR ENGAGEMENT IN DRUG CHECKING, NATIONALLY, 2019-2026	125
FIGURE 44: SELF-REPORTED CRIMINAL ACTIVITY IN THE PAST MONTH, NATIONALLY, 2007-2026	127
FIGURE 45: VICTIM OF CRIME INVOLVING VIOLENCE IN THE PAST MONTH, NATIONALLY, 2019-2026	128
FIGURE 46: LIFETIME INCARCERATION, AND PAST 12 MONTH ARREST AND DRUG-RELATED ENCOUNTERS WITH POLICE THAT DID NOT RESULT IN ARREST, NATIONALLY, 2007-2026.....	129

Acknowledgements

Funding

In 2026, the Illicit Drug Reporting System (IDRS), falling within the Drug Trends program of work, was supported by funding from the Australian Government Department of Health, Disability and Ageing under the Drug and Alcohol Program.

Research Team

The National Drug and Alcohol Research Centre (NDARC), UNSW Sydney, coordinated the IDRS. The following researchers and research institutions contributed to the IDRS in 2026:

- Dr Rachel Sutherland, Antonia Karlsson, Julia Uporova, Udesha Chandrasena, Lily Palmer, Agata Chrzanowska, Professor Louisa Degenhardt, Professor Michael Farrell and Associate Professor Amy Peacock, National Drug and Alcohol Research Centre, University of New South Wales Sydney, New South Wales;
- Joanna Wilson, Dorian Raffaele, Matthew Cowman and Professor Paul Dietze, Disease Elimination Program, Burnet, Victoria;
- Sophie Radke and Professor Raimondo Bruno, School of Psychology, University of Tasmania, Tasmania; and
- Jack Curran and Professor Simon Lenton, National Drug Research Institute and EnAble Institute, Curtin University, Western Australia.

We would like to thank past and present members of the research team.

Participants

We would like to thank all the participants who were interviewed for the IDRS in the present and in previous years.

Contributors

We thank all the individuals who contributed to questionnaire development and assisted with the collection and input of data at a jurisdictional and national level. We would like to thank the members of the Drug Trends Advisory Committee, as well as the Australian Injecting & Illicit Drug Users League (AIVL), for their contribution to the IDRS.

We acknowledge the Traditional Custodians of the lands on which this work was undertaken. We pay our respects to Elders past and present, and extend that same respect to all Aboriginal and Torres Strait Islander peoples. We recognise their continuing connection to Country, culture and community.

Abbreviations

ACT	Australian Capital Territory
Ade	Adelaide
AIVL	Australian Injecting & Illicit Drug Users League
ADHD	Attention-deficit hyperactivity disorder
Alpha PVP	α -Pyrrolidinopentiophenone
AOD	Alcohol and Other Drugs
AUDIT-C	Alcohol Use Disorders Identification Test-Concise
Bri/GC	Brisbane/Gold Coast
Can	Canberra
CBD	Cannabidiol
COVID-19	Coronavirus Disease 2019
Dar	Darwin
DSM	Diagnostic and Statistical Manual of Mental Disorders
EDRS	Ecstasy and Related Drugs Reporting System
GHB/GBL/1,4-BD	Gamma-hydroxybutyrate/Gamma-butyrolactone/1,4-Butanediol
GP	General Practitioner
HCV	Hepatitis C Virus
HIV	Human immunodeficiency virus
Hob	Hobart
IDRS	Illicit Drug Reporting System
IQR	Interquartile Range
K10	Kessler Psychological Distress Scale
LSD	<i>d</i> -lysergic acid
MDA	3,4-methylenedioxyamphetamine
MDPV	Methylenedioxypropylone
Mel	Melbourne
N (or n)	Number of Participants
NDARC	National Drug and Alcohol Research Centre
NPS	New Psychoactive Substances
NSP	Needle and Syringe Program
NSW	New South Wales
NT	Northern Territory
OTC	Over-the-Counter
PBS	Pharmaceutical Benefits Scheme
PCR	Polymerase Chain Reaction
Per	Perth
PTSD	Post-traumatic stress disorder

QLD	Queensland
RNA	Ribonucleic Acid
SA	South Australia
SD	Standard Deviation
SDS	Severity of Dependence Scale
STI	Sexually Transmitted Infection
Syd	Sydney
TAS	Tasmania
TGA	Therapeutic Goods Administration
THC	Tetrahydrocannabinol
UNSW	University of New South Wales
VIC	Victoria
WA	Western Australia
WHO	World Health Organization

Executive Summary

The IDRS comprises a sentinel sample of people who regularly inject illicit drugs, recruited via advertisements in needle syringe programs and other harm reduction services, as well as via peer referral, across each capital city of Australia. The results are not representative of all people who inject drugs, nor of use in the general population. **Data were collected in 2026 from May-July. Interviews from 2020 onwards were delivered face-to-face as well as via telephone; all interviews prior to 2020 were conducted face-to-face. This methodological change should be factored into all comparisons of data from the 2020-2026 samples relative to previous years.**

Sample Characteristics

The 2026 IDRS sample (N=902) was mostly similar to the sample in 2025 and in previous years. Participants predominantly identified as male (63%) with a median age of 47 years, both stable from 2025. In 2026, most participants reported being unemployed at the time of the interview (89%; 88% in 2025). However, the median weekly income significantly increased from \$465 in 2025 to \$500 in 2026 ($p<0.001$). Drug of choice remained stable, with 44% nominating methamphetamine (44% in 2025) and 41% nominating heroin (42% in 2025) as their drug of choice. Most participants continued to nominate methamphetamine as the drug injected most often in the past month (57%; 57% in 2025).

Heroin

In 2026, half (51%) of the sample reported recent heroin use, stable relative to 2025 (54%). However, frequency of use significantly increased, from a median of 120 days in the

past six months in 2025 to a median of 180 days in 2026 ($p=0.011$). The median price for one point of heroin in 2026 was \$50 (IQR=50-80; \$50 in 2025; IQR=50-90; $p=0.041$). Perceived purity and perceived availability remained stable, with 33% reporting 'medium' purity (33% in 2025) and 95% reporting 'easy' or 'very easy' obtainment (93% in 2025).

Methamphetamine

Recent use of any methamphetamine gradually increased between 2010 and 2021, however has remained relatively stable since. In 2026, 81% of the sample reported recent use on a median of 90 days, stable from 2025 (82%; median 80 days). Crystal continued to be the most commonly used form of methamphetamine (79%; 79% in 2025), with use of methamphetamine powder (6%; 8% in 2025) and methamphetamine base (1%; 3% in 2025) remaining low and stable. The reported median price for one point of crystal was \$50, stable from 2025. Perceived purity and perceived availability of methamphetamine crystal remained stable, with 31% reporting 'medium' purity (34% in 2025) and 96% reporting 'easy' or 'very easy' obtainment (96% in 2025).

Cocaine

In 2026, 18% of participants reported recent cocaine use on a median of three days, stable relative to 2025 (19%; median 2 days). The median price for one gram of cocaine remained stable at \$350 (\$300 in 2025). Perceived purity of cocaine also remained stable, with 49% of participants reporting 'high' purity (36% in 2025). However, perceived availability significantly changed in 2026, relative to 2025 ($p=0.014$), with more participants perceiving

availability to be 'easy' or 'very easy' (79%; 62% in 2025).

Cannabis and/or Cannabinoid-Related Products

Recent use of non-prescribed cannabis and/or cannabinoid-related products has gradually declined over time. In 2026, 61% of participants reported recent non-prescribed use, the lowest per cent observed over the past 20 years, although stable relative to 2025 (65% in 2025). Forty-five per cent of those who had recently used non-prescribed cannabis and/or cannabinoid-related products reported daily use (50% in 2025). Perceived purity and perceived availability of bush and hydroponic cannabis remained stable in 2026, relative to 2025. The median price per gram of hydro and bush cannabis also remained stable at \$20, respectively.

Pharmaceutical Opioids

Overall, non-prescribed use of most forms of pharmaceutical opioids has either declined or remained stable over the past two decades. In 2026, oxycodone, methadone and morphine were the most commonly used non-prescribed pharmaceutical opioids (8%, 7% and 7%, respectively), with significant decreases observed in recent use of non-prescribed buprenorphine-naloxone (4%; 6% in 2025; $p=0.034$), non-prescribed buprenorphine tablet (2%; 4% in 2025; $p=0.003$), and non-prescribed morphine (7%; 9% in 2025; $p=0.047$).

Other Drugs

Use of any NPS has gradually declined since monitoring commenced in 2013, with 4% reporting recent use in 2026 (6% in 2025). Benzodiazepines remained the most common non-prescribed pharmaceutical drug reported

by participants in 2026 (21%; 23% in 2025), followed by pregabalin (14%; 16% in 2025). Recent use of non-prescribed pharmaceutical stimulants significantly decreased in 2026 (5%; 8% in 2025; $p=0.021$). Recent use of tobacco remained high yet stable (86%; 87% in 2025), however use of illicit tobacco increased significantly (70%; 63% in 2025; $p=0.009$). Illicit e-cigarettes were used by one quarter (28%) of the sample in 2026, stable relative to 2025 (25%), with participants reporting use on a median of 90 days in the six months preceding interview. Use of all other monitored drugs remained stable.

Drug-Related Harms and Other Behaviours

Polysubstance use and bingeing

In 2026, 63% of the sample reported using two or more drugs (excluding tobacco and e-cigarettes) on the day preceding interview.

Three fifths (60%) of the sample had binged on one or more drugs for 48 hours or more in the six months preceding interview.

Injecting behaviours and equipment access

In 2026, 5% of participants reported receptive sharing of a needle or syringe (5% in 2025) and 8% reported distributive sharing (8% in 2025) in the past month. One-third (33%) of participants reported re-using their own needles in the past month, stable relative to 2025 (31%), and one fifth (20%) reported sharing other equipment (19% in 2025).

Twenty-eight per cent of the sample reported having an injection-related problem in the past month (30% in 2025), most commonly any infection/abscess (12%; 14% in 2025) and any nerve damage (9%; 12% in 2025).

Overdose, naloxone and drug checking

In 2026, one fifth (19%) of the sample reported any past 12-month non-fatal overdose, stable relative to 2025 (19%), most commonly from heroin (10%; 10% in 2025).

Awareness of take-home naloxone remained stable in 2026 (79%) relative to 2025 (80%), as did past year access (56%; 55% in 2025).

In 2026, 12% of participants reported that they or someone else had tested the contents and/or purity of their illicit drugs in Australia in the past year (15% in 2025).

Dependence, drug treatment and HCV

In 2026, 34% of male participants obtained an AUDIT-C score of ≥ 4 , and 33% of female participants obtained a score of ≥ 3 , indicative of hazardous alcohol use.

Forty-seven per cent of those who reported recent methamphetamine use obtained an SDS score of ≥ 4 , a significant decrease from 54% in 2025 ($p=0.014$), while 66% of participants reporting recent opioid use obtained a score of ≥ 5 , a significant increase from 59% in 2025 ($p=0.030$), indicating possible dependence on these substances.

Thirty-seven per cent of the sample reported that they were in any drug treatment for their substance use at the time of interview (38% in 2025), with the most common treatment being methadone (21%).

In 2026, 55% of participants reported that they had received a hepatitis C virus (HCV) antibody test in the past year (51% in 2025), and 43% had received an PCR or RNA test (39% in 2025). Three per cent reported having a current HCV infection (4% in 2025) and 6% reported that they had received HCV treatment in the past year (5% in 2025). Almost one third (31%) of the

sample reported having had a test for human immunodeficiency virus (HIV) in the past six months, stable relative to 2025 (30%), while the proportion reporting lifetime testing increased significantly (84%; 79%; $p=0.030$).

Sexual activity, mental health, disability and health service access

In 2026, 45% of the sample reported engaging in sexual activity in the past four weeks and 24% reported having a sexual health check-up in the past six months.

Self-reported mental health problems remained stable in 2026 (53%) relative to 2025 (55%). Three fifths (63%) reported high/very high psychological distress (62% in 2025). Half (53%) self-identified as a person living with a disability (52% in 2025).

Most participants (87%) reported accessing any health service for alcohol and/or drug support in the six months preceding interview in 2026 (90% in 2025).

Driving, contact with police and modes of purchasing drugs

Seventy-seven per cent of those who had recently driven reported driving within three hours of consuming an illicit or non-prescribed drug in the past six months (77% in 2025) and 8% reported driving while over the perceived legal limit of alcohol (9% in 2025).

One third (33%) of participants reported a drug-related encounter with police which did not result in charge or arrest, stable relative to 2025 (29%).

In 2026, the most popular means of arranging the purchase of illicit or non-prescribed drugs in the 12 months preceding interview was face-to-face (84%), a significant increase relative to 2025 (79%; $p=0.019$).

2026 SAMPLE CHARACTERISTICS




Between May-July, 902 participants, recruited from all capital cities across Australia, were interviewed.



47 years **Male**

Median age and per cent who identified as male.



89% **19%**

Unemployed No fixed address

Per cent who were unemployed or had no fixed address at the time of interview.




Injected heroin

Injected methamphetamine

Injected other illicit or non-prescribed drugs


Participants were recruited on the basis that they had injected drugs on at least 6 days in the previous 6 months.

INJECTING-RELATED RISKS AND HARMS



5% **8%**


5% of participants reported receptive sharing in the past month, and 8% reported distributive sharing.



31% **33%**


2025 2026

Percentage who reported re-using their own needles in the past month.




36%

Percentage who reported injecting someone else in the past month.



28% of participants reported having an injection-related health issue in the past month, stable from 30% in 2025.

OTHER HARMS



Two or more drugs 63%

Stimulants and cannabis 16%

Opioids and cannabis 10%

The per cent who reported using ≥2 drugs on the day preceding interview, and the 2 most common polysubstance use profiles.




19% **19%**

2025 2026

Past year non-fatal overdose remained stable in 2026 relative to 2025.



53% of participants self-reported a mental health problem in the 6 months preceding interview.




Depression 61%

Anxiety 55%

PTSD 34%

Among those who reported a mental health problem, the 3 most common mental health issues were depression, anxiety and PTSD.

NALOXONE AND OTHER HARM REDUCTION STRATEGIES



79% **56%**


Heard of take-home naloxone Naloxone obtained

Knowledge of take-home naloxone and past year naloxone obtained remained stable in 2026 relative to 2025.




36%

36% reported ever using naloxone to resuscitate someone who had overdosed, with 21% having done so in the past year.



12%

Percentage who reported that they or someone else had tested the content and/or purity of their illicit drugs in Australia in the past year.



37% **87%**


Drug treatment Health service access

Percentage who reported current drug treatment and health service access for AOD support in the past 6 months.

HEROIN



Past 6 month use of heroin remained stable in 2026 (51%) relative to 2025 (54%).




Of those who had recently consumed heroin, 51% reported daily use, a significant increase from 42% in 2025.



\$50 2025 **\$50** 2026


The median reported price for 1 point of heroin.




Percentage who perceived heroin as being 'easy' or 'very easy' to obtain.

METHAMPHETAMINE

FORM of methamphetamine



Past 6 month use of any methamphetamine, crystal, powder and base remained stable in 2026, relative to 2025.




Of those who had recently used any form of methamphetamine, 79% reported weekly or more frequent use, stable from 2025 (79%).




\$50 2025 **\$50** 2026

The median reported price for 1 point of methamphetamine crystal.




Percentage who perceived methamphetamine crystal as being 'easy' or 'very easy' to obtain.

CANNABIS AND/OR CANNABINOID-RELATED PRODUCTS



Past non-prescribed 6 month use remained stable in 2026 (61%) relative to 2025 (65%).




Of those who had recently used non-prescribed cannabis/cannabinoid-related products, 45% reported daily use, stable from 2025 (50%).



\$20 Hydroponic **\$20** Bush

The median reported price for 1 gram of hydroponic and bush cannabis.




Percentage who perceived cannabis/cannabinoid-related products as being 'easy' or 'very easy' to obtain.


PAST 6 MONTH USE OF OTHER DRUGS

* $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$.


Non-prescribed morphine




Non-prescribed fentanyl



Non-prescribed oxycodone



GHB/GBL/1,4-BD



Background and Methods

The Illicit Drug Reporting System (IDRS) interviews are conducted annually with a sentinel cross-sectional group of people who regularly inject drugs, recruited from all capital cities of Australia (N=902 in 2026). The results from the IDRS interviews are not representative of all people who consume drugs, nor of illicit drug use in the general population, but this is not the aim of these data. Rather, these data are intended to provide evidence indicative of trends that warrant further monitoring. These findings should be interpreted alongside analyses of other data sources for a more complete profile of trends in illicit drug use, market features, and harms in Australia.



1

Background

The [Illicit Drug Reporting System \(IDRS\)](#) is an ongoing illicit drug monitoring system which has been conducted in all states and territories of Australia since 2000, and forms part of [Drug Trends](#). The purpose of the IDRS is to provide a coordinated approach to monitoring the use, market features, and harms of illicit drugs.

The IDRS is designed to be sensitive to emerging trends, providing data in a timely manner, rather than describing issues in extensive detail. It does this by studying a range of data sources, including data from annual interviews with people who regularly inject illicit and/or non-prescribed drugs and from secondary analyses of routinely-collected indicator data. This report focuses on the key results from the annual interview component of the IDRS.

Methods

IDRS 2000-2019

Full details of the [methods for the annual interviews](#) are available for download. To briefly summarise, participants were recruited using multiple methods (e.g., via needle and syringe programs (NSP) and peer referral) and needed to: i) be at least 17 years of age (due to ethical requirements); ii) have injected psychoactive illicit or non-prescribed drugs at least six days during the six months preceding interview; and iii) have been a resident of the capital city in which the interview took place for ten of the past 12 months. Interviews took place in varied locations negotiated with participants (e.g., treatment services, coffee shops or parks), and in later years were conducted using REDCap (Research Electronic Data Capture), a software program to collect data on laptops or tablets. Following provision of written informed consent and completion of a structured interview, participants were reimbursed \$40 cash for their time and expenses incurred.

IDRS 2020-2026

Given the emergence of COVID-19 and the resulting restrictions on travel and people's movement in Australia (which first came into effect in March 2020), face-to-face interviews were not always possible due to the risk of infection transmission for both interviewers and participants. For this reason, all methods in 2020 were similar to previous years as detailed above, with the exception of:

1. Means of data collection: Interviews were conducted via telephone across all capital cities in 2020, with some capital cities (Darwin, Northern Territory (NT) and Hobart, Tasmania (TAS)) also offering face-to-face interviews;
2. Means of consenting participants: Participants' consent to participate was collected verbally prior to beginning the interview;
3. Means of reimbursement: Participants were given the option of receiving \$40 reimbursement via one of three methods, comprising bank transfer, PayID or gift voucher, where completing the interview via telephone; and
4. Age eligibility criterion: Changed from 17 years old (16 years old in Perth, Western Australia (WA)) to 18 years old.

From 2021 onwards, a hybrid approach was used whereby interviews were conducted either face-to-face (with participants reimbursed with cash) or via telephone/videoconference (with participants reimbursed via bank transfer or other electronic means). Face-to-face interviews were the preferred methodology, however telephone interviews were conducted when required (i.e., in accordance with government directives) or when requested by services. Consent was collected verbally for all participants.

In 2026, participant reimbursement increased from \$40 (2000-2025) to \$50 to provide more appropriate remuneration for participation and to better align with community expectations and contemporary research reimbursement practices.

2026 IDRS Sample

Between 12 May-10 July 2026, a total of 902 participants were recruited across capital cities nationally. The number of participants recruited from the capital city in each jurisdiction were: Sydney, NSW, n=151; Melbourne, VIC, n=149; Adelaide, SA, n=100; Canberra, ACT, n=100; Hobart, TAS, n=102; Brisbane and Gold Coast, QLD, n=100; Darwin, NT, n=100; and Perth, WA, n=100. One tenth (10%; n=88) of all 2026 interviews were conducted via telephone/videoconference: Sydney, NSW n=10, Adelaide, SA n=2; Hobart, TAS n=17; Brisbane/Gold Coast, QLD, n=15; and Perth, WA n=45.

In 2026, most participants continued to be recruited via needle and syringe programs (NSP) (57%; 55% in 2025) and word-of-mouth (36%; 37% in 2025). Few participants (3%) were recruited via a treatment provider (2% in 2025) or via social media (1%; 2% in 2025). One fifth (19%) of the 2026 sample had taken part in the 2025 interview (24% of the 2025 sample had taken part in the 2024 interview; $p=0.040$).

Data Analysis

For normally distributed continuous variables, means and standard deviations (SD) are reported; for skewed data (i.e., skewness $> \pm 1$ or kurtosis $> \pm 3$), medians and interquartile ranges (IQR) are reported. Tests of statistical significance have been conducted between estimates for 2025 and 2026. Note that no corrections for multiple comparisons have been made and thus comparisons should be treated with caution. Values where cell sizes are ≤ 5 have been suppressed with corresponding notation (zero values are reported). References to 'recent' use and behaviours refers to the six months preceding interview. The response options 'Don't know' and 'Skip question', which were available to select throughout the interview, were excluded from analysis.

Although national IDRS data collection commenced in 2000, this report presents data from the last 20 years only (2007-2026). Data from earlier years can be found in historical IDRS reports, available via the [Drug Trends webpage](#).

Guide to Table/Figure Notes

Table 1: Guide to Table/Figure Notes

Legend	
	Empty cell(s) indicate question not asked in respective year (for figures)
/	Question not asked in respective year (for tables)
-	Per cent suppressed due to small cell size ($n \leq 5$ but not 0) (for figures and tables)
*$p < 0.050$; **$p < 0.010$; ***$p < 0.001$	Statistical significance between 2025 and 2026
Syd	Sydney
Can	Canberra
Mel	Melbourne
Hob	Hobart
Ade	Adelaide
Per	Perth
Dar	Darwin
Bri/GC	Brisbane and the Gold Coast (and the Sunshine Coast 2014-16)

Interpretation of Findings

Caveats to interpretation of findings are discussed more completely in the [methods for the annual interviews](#) but it should be noted that these data are from participants recruited in capital cities, and thus do not reflect trends in regional and remote areas. Further, the results are not representative of all people who consume illicit drugs, nor of illicit drug use in the general population, but rather are intended to provide evidence indicative of emerging issues that warrant further monitoring.

This report covers a subset of items asked of participants and does not include jurisdictional-level results beyond estimates of recent use of various substances (comprehensive jurisdictional findings are provided separately; see below), nor does it include implications of findings. These findings should be interpreted alongside analyses of other data sources for a more complete profile of trends in illicit drug use, market features, and harms in Australia (see section on 'Additional Outputs' below for details of other outputs providing such profiles).

Additional Outputs

[Infographics](#) and the [executive summary](#) from this report are available for download. There are a range of outputs from the IDRS which triangulate key findings from the annual interviews and other data sources, including national reports, jurisdictional reports, bulletins, and other resources available via the [Drug Trends webpage](#). This includes results from the [Ecstasy and Related Drugs Reporting System \(EDRS\)](#), which focuses on the use of ecstasy and other illicit stimulants.

Please contact the research team at drugtrends@unsw.edu.au with any queries, to request additional analyses using these data, or to discuss the possibility of including items in future interviews.

2 Sample Characteristics


Participants were asked questions about select sociodemographic characteristics, as well as key drug use characteristics of interest.

N=902

Median age



Gender



Born in Australia



Sexual identity



Mean years of school education



Post-school qualification(s)




Employment status




Median weekly income




Accommodation




Drug of choice



Drug injected most often in the past month



Weekly or more frequent use in the past 6 months



Sample Characteristics

In 2026, the median age of the sample was 47 years (IQR=41-54; 47 years in 2025; IQR=41-54; $p=0.693$). Gender remained stable compared to 2025 ($p=0.466$), with almost two thirds (63%) of the sample identifying as male (66% in 2025) (Table 2). In 2026, current employment status remained stable relative to 2025 ($p=0.552$), with most participants reporting that they were unemployed at the time of the interview (89%; 88% in 2025). However, the median weekly income significantly increased from \$465 (IQR=375-598) in 2025 to \$500 (IQR=400-600) in 2026 ($p<0.001$). Current accommodation remained stable between 2025 and 2026 ($p=0.994$), with two thirds of participants reporting living in their own house/flat (67%; 66% in 2025). One fifth (19%) reported having 'no fixed address' in 2026 (19% in 2025), followed by 5% residing in a parent's/family home (5% in 2025). In 2026, the per cent of participants that reported English as their primary language spoken at home remained stable compared to the 2025 sample (97%; 98% in 2025; $p=0.215$), as did the percentage who reported being born in Australia (90%; 91% in 2025; $p=0.569$). In 2026, significantly more participants reported having a post-school qualification/s (62%) relative to 2025 (57%; $p=0.035$) (Table 2).

In 2026, drug of choice remained stable compared to 2025 ($p=0.164$), with 44% of the sample nominating methamphetamine as their drug of choice (44% in 2025), followed by two fifths (41%) nominating heroin (42% in 2025) as their drug of choice (Figure 1). The drug injected most often in the past month also remained stable in 2026 relative to 2025 ($p=0.233$), with methamphetamine reported as the drug injected most often by 57% of the sample (57% in 2025) (Figure 2), followed by heroin (37%; 37% in 2025).

Weekly or more frequent consumption of 'any' methamphetamine (63% versus 65%; $p=0.657$) and heroin (42% in both years) remained stable in 2026 relative to 2025. Weekly or more frequent use of non-prescribed cannabis also remained stable in 2026 (49%) relative to 2025 (52%; $p=0.159$) (Figure 3).


Table 2: Demographic characteristics of the sample, nationally, 2025-2026, and by capital city, 2026

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=884	N=902	N=151	N=100	N=149	N=102	N=100	N=100	N=100	N=100
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
Median age (years; IQR)	47 (41-54)	47 (41-54)	47 (41-54)	47 (42-54)	48 (43-53)	45 (40-52)	48 (41-56)	48 (40-56)	44 (37-51)	49 (43-53)
% Gender										
Female	34	36	32	28	37	36	41	38	41	39
Male	66	63	68	70	62	61	59	62	59	59
Non-binary	1	1	-	-	-	-	0	0	0	-
% Aboriginal and/or Torres Strait Islander	29	27	35	20	29	21	26	15	48	19
% Born in Australia	91	90	86	93	89	97	95	86	92	86
% English primary language spoken at home	98	97	95	98	93	100	99	99	96	100
% Sexual identity										
Heterosexual	82	83	72	85	84	86	90	91	88	74
Homosexual	5	4	10	-	6	-	-	-	-	-
Bisexual	9	10	11	9	8	11	8	8	7	17
Queer	2	-	-	-	-	0	0	0	-	0
Other identity	1	2	5	-	-	-	-	0	-	-
Mean years of school education (range)	10 (1-12)	10 (1-12)	10 (5-12)	10 (4-12)	10 (4-12)	10 (3-12)	10 (1-12)	10 (3-12)	10 (5-12)	11 (5-12)
% Post-school qualification(s) ^	57	62*	55	64	57	71	70	39	63	81
% Current employment status										
Unemployed	88	89	91	92	95	92	84	91	82	82
Employed full time	3	2	-	0	0	-	-	-	-	7
Part time/casual	6	6	4	-	-	-	8	6	10	10
Self-employed	2	3	-	-	-	-	7	-	-	-
Other	0	0	0	0	0	0	0	0	0	0
% Past month gov't pension, allowance or benefit	94	94	94	97	95	98	92	94	89	90
Current median weekly income (\$; IQR)	\$465 (375-598)	\$500*** (400-600)	473 (390-550)	500 (400-600)	516 (400-600)	500 (405-600)	450 (400-600)	450 (400-500)	500 (450-650)	500 (433-600)
% Current accommodation										
Own home (incl. renting)	66	67	72	75	58	63	65	64	63	76
Parents'/family home	5	5	-	-	6	6	10	6	6	-

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=884	N=902	N=151	N=100	N=149	N=102	N=100	N=100	N=100	N=100
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
Boarding house/hostel	4	4	-	-	4	6	-	-	7	-
Shelter/refuge	4	4	-	0	5	6	0	15	-	-
No fixed address+	19	19	21	18	23	21	20	12	20	14
Other	1	2	0	-	4	-	-	-	-	-


Note. ^Includes trade/technical and university qualifications. +No fixed address includes couch surfing and rough sleeping or squatting. Statistical significance for 2025 versus 2026 is presented in table for national estimates; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. For sample characteristics over the whole duration of the project, see [methods for the annual interviews](#). Please refer to Table 1 for a guide to table/figure notes.

Figure 1: Drug of choice, nationally, 2007-2026




Note. Participants could only endorse one substance. Substances listed in this figure are the primary endorsed; a nominal per cent endorsed other substances. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 2: Drug injected most often in the past month, nationally, 2007-2026



Note. Participants could only endorse one substance. Substances listed in this figure are the primary endorsed; a nominal per cent endorsed other substances. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 3: Weekly or more frequent substance use in the past six months, nationally, 2007-2026



Note. Computed of the entire sample regardless of whether they had used the substance in the past six months. Prior to 2021, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2017-2020 figures include some participants who were using prescribed cannabis only (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low. Further, from 2022, we captured use of 'cannabis and/or cannabinoid-related products', while in previous years questions referred only to 'cannabis'. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Past 6 months use

51%

3 Heroin

Participants were asked about their recent (past six month) use of heroin and homebake heroin.

Participants typically describe heroin as white/off-white rock, brown/beige rock or white/off-white powder. Homebake is a form of heroin illicitly produced from pharmaceutical opioid products.

Median days*

180 days

*Among those whom reported past 6 month use in 2026:

Routes of administration

Injecting
98%

Smoking

Snorting

Swallowing

Median Amount Used

Typical day:

0.20 grams

Maximum:

0.40 grams

F
O
R
M
S

72% white/off-white rock

54% white/off-white powder

25% brown/beige rock

19% brown/beige powder

2% homebake

Median Price

\$ 50
Per point

\$ 300
Per gram

Perceived Purity

13% fluctuates

31%
high

23%
low

33%
medium

Perceived Availability

4% difficult

30%
easy

65%
very
easy

n ≤ 5 very difficult

Patterns of Consumption

Recent Use (past 6 months)

Over the past 20 years, past six-month heroin use has ranged from a low of 50% to a high of 64%. In 2026, half (51%) of the sample reported recent use of heroin (54% in 2025; $p=0.343$) (Figure 4). Recent use of heroin remained stable in all capital city samples (Table 3).

Frequency of Use

Among those who reported recent heroin use and commented in 2026 ($n=463$), participants reported use on a median of 180 days in the past six months (i.e., daily use; $IQR=48-180$). This was a significant increase relative to 2025 (120 days; $IQR=24-180$; $n=465$; $p=0.011$) and is the highest frequency of use observed in the past 20 years (Figure 4). Weekly or more frequent use among those who reported recent use was stable in 2026 (82%) compared to 2025 (78%; $p=0.106$), however, daily use significantly increased (51%; 42% in 2025; $p=0.009$).

Routes of Administration

Injecting remained the most common route of administration among participants who consumed heroin in 2026 (98%; 99% in 2025; $p=0.236$). Participants who reported injecting did so on a median of 180 days ($IQR=48-180$), a significant increase relative to 2025 (120 days; $IQR=24-180$; $p=0.003$). Fewer participants reported smoking (8%; 9% in 2025; $p=0.807$) or snorting (1%; 1% in 2025) as routes of administration.


Quantity

Of those who reported recent use and responded ($n=439$), the median amount of heroin used on a 'typical' day in the last six months was 0.20 grams ($IQR=0.10-0.40$; 0.20 grams in 2025; $IQR=0.10-0.40$; $n=443$; $p=0.026$). The median maximum amount of heroin used per day in the last six months was 0.40 grams ($IQR=0.20-0.80$; $n=437$; 0.40 grams in 2025; $IQR=0.15-1.00$; $n=437$; $p=0.921$).

Forms Used

Among participants who reported recent use of heroin and commented ($n=458$), 72% reported using white/off-white rock (72% in 2025; $p=0.374$), and 54% reported using white/off-white powder (47% in 2025; $p=0.284$). One quarter (25%) reported using brown/beige rock (28% in 2025; $p=0.271$), and one fifth (19%) reported using brown/beige powder (16% in 2025; $p=0.565$). Two per cent reported using homebake in 2026 (2% in 2025; $p=0.808$), and few participants ($n\leq 5$) reported using purple rock or purple powder ($n\leq 5$ in 2025).

Figure 4: Past six month use and frequency of use of heroin, nationally, 2007-2026



Note. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 3: Past six month use of heroin, by capital city, 2007–2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	88	72	85	-	67	57	7	65
2008	83	86	85	-	51	59	14	74
2009	94	78	79	12	72	71	13	75
2010	92	78	85	8	64	69	5	81
2011	87	79	81	19	57	79	9	65
2012	89	74	84	9	52	80	11	65
2013	83	75	83	10	41	75	17	72
2014	85	75	83	13	43	79	7	66
2015	91	79	74	-	49	75	14	50
2016	86	70	77	7	37	78	7	58
2017	80	74	80	15	52	66	13	55
2018	83	75	83	8	35	67	9	45
2019	82	77	85	15	28	62	-	63
2020	78	85	86	24	47	69	-	64
2021	75	78	76	11	23	61	-	43
2022	72	66	78	22	35	60	-	51
2023	67	81	87	11	22	63	~	46
2024	73	74	83	12	21	70	-	37
2025	69	81	81	16	28	65	-	39
2026	77	69	79	14	27	61	10	48

Note. ~Due to the particularly small sample recruited in Darwin in 2023 ($n < 50$), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Price, Perceived Purity and Perceived Availability

Price

In 2026, the reported median price for one point of heroin was \$50 (IQR=50-80; $n=204$; \$50 in 2025; IQR=50-90; $n=215$; $p=0.041$). The median price per cap remained stable (\$50; IQR=50-50; $n=19$; \$50 in 2025; IQR=50-50; $n=21$; $p=0.967$), as did the median price of one gram (\$300 in 2026; IQR=250-400; $n=53$; \$400 in 2025; IQR=273-500; $n=32$; $p=0.243$) (Figure 5).


Perceived Purity

Among those who were able to comment in 2026 ($n=413$), the perceived purity of heroin remained stable relative to 2025 ($p=0.981$). One third (33%) perceived the purity of heroin to be 'medium' in 2026 (33% in 2025), 31% reported 'high' purity (30% in 2025), and 23% reported 'low' purity (23% in 2025) (Figure 6).

Perceived Availability


Among those who were able to comment in 2026 ($n=428$), the perceived availability of heroin remained stable relative to 2025 ($p=0.356$). Two thirds (65%) of participants perceived heroin to be 'very easy' to obtain (60% in 2025), and 30% reported that it was 'easy' to obtain (33% in 2025). Four per cent perceived it to be 'difficult' to obtain (6% in 2025) (Figure 7).

Figure 5: Median price of heroin per cap, point and gram, nationally, 2007-2026




Note. Among those who commented. Between 2009-2017 a cap was referred to as cap/point (thus, the same values appear for cap and point during this period); in 2018 these measures were separated out into their own response options. The error bars represent the IQR. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 6: Current perceived purity of heroin, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 7: Current perceived availability of heroin, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Past 6 months use

81%

4

Methamphetamine

Participants were asked about their recent (past six month) use of various forms of methamphetamine, including powder (white particles, described as speed), base (wet, oily powder) and crystal (clear, ice-like crystals).

Median days*

90 days

*Among those whom reported past 6 month use in 2026:

FORMS

79% crystal

6% powder

1% base

Median Amount Used (Crystal)

Typical day:

0.20 grams

Maximum:

0.40 grams

Routes of administration (Crystal)

Injecting
96%

Smoking
46%

Snorting

Swallowing

Median Price (Crystal)

\$ 50
Per point

\$ 300
Per gram

(Crystal)

Perceived Purity

17% fluctuates

26% high

31% medium

26% low

4% difficult

26% easy

70% very easy

(Crystal)

Perceived Availability

Patterns of Consumption (Any Methamphetamine)

Recent Use (past 6 months)

The per cent of participants reporting recent use of any methamphetamine gradually increased between 2010 and 2021, before subsequently stabilising. In 2026, 81% reported recent use, stable from 2025 (82%; $p=0.671$) (Figure 8). Any recent methamphetamine use remained high and stable across all capital city samples (Table 4).

Frequency of Use

In 2026, participants reported using methamphetamine on a median of 90 days (i.e. every other day; IQR=24-180; $n=727$), the highest frequency of use observed in the past 20 years, although stable relative to 2025 (80 days; IQR=24-180; $n=703$; $p=0.445$) (Figure 9). Weekly or more frequent use, among those who reported recent use and commented ($n=727$), also remained stable in 2026 (79%; 79% in 2025; $p=0.843$), as did the per cent reporting daily use (28%; 26% in 2025; $p=0.437$).


Forms Used

The forms of methamphetamine used by participants have shifted over time, with use of methamphetamine powder and base decreasing and use of methamphetamine crystal increasing (Figure 8). In 2026, among those who reported recent use of any methamphetamine ($n=729$), nearly all (98%; 97% in 2025; $p=0.492$) reported using methamphetamine crystal, followed by 7% reporting methamphetamine powder (10% in 2025; $p=0.073$) and 2% reporting methamphetamine base (3% in 2025; $p=0.096$).

Number of Forms Used


Among participants who had recently consumed any methamphetamine and commented in 2026 ($n=729$), the median number of forms of methamphetamine used was one (IQR=1-1), stable relative to 2025 (1 form; IQR=1-1; $n=706$; $p=0.052$). This was consistent across jurisdictions.

Figure 8: Past six month use of any methamphetamine and of methamphetamine powder, base, and crystal, nationally, 2007-2026



Note. 'Any methamphetamine' includes crystal, powder, base and liquid methamphetamine combined from 2007-2018, and crystal, powder and base methamphetamine combined from 2019 onwards. Questions regarding liquid methamphetamine not asked from 2019. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 9: Frequency of use of any methamphetamine and of methamphetamine powder, base, and crystal, nationally, 2007-2026



Note. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Y axis reduced to 90 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 4: Past six month use of any methamphetamine, by capital city, 2007–2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	62	83	74	88	74	70	68	78
2008	74	74	68	74	69	74	57	59
2009	57	75	70	80	61	63	55	70
2010	57	59	60	70	74	64	36	59
2011	60	73	65	77	66	64	55	71
2012	72	77	67	77	79	72	48	53
2013	75	66	61	74	75	72	43	58
2014	75	76	77	70	75	66	37	72
2015	66	81	74	72	76	71	67	67
2016	77	83	73	75	77	65	71	70
2017	69	80	66	69	76	70	66	74
2018	76	85	78	79	83	67	75	72
2019	76	79	70	81	90	79	90	68
2020	77	65	66	77	81	73	83	63
2021	74	75	79	89	88	82	76	79
2022	87	81	75	84	90	78	80	70
2023	86	75	77	88	91	79	~	73
2024	83	85	70	91	96	65	87	85
2025	78	73	73	95	93	71	96	88
2026	76	83	69	95	92	70	91	78

Note. ~Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Patterns of Consumption (by form)

Methamphetamine Powder

Recent Use (past 6 months): Use of methamphetamine powder has decreased substantially over time. In 2026, 6% reported recent use, the lowest per cent recorded in the past two decades, though stable relative to 2025 (8%; $p=0.062$) (Figure 8). The per cent reporting recent use remained low and stable across all capital city samples (Table 5).

Frequency of Use: Participants reported using methamphetamine powder on a median of 30 days in the preceding six months (IQR=6-95; $n=51$), stable relative to 2025 (40 days; IQR=10-120; $n=68$; $p=0.549$) (Figure 9). In 2026, three fifths (59%) of those who had recently used methamphetamine powder reported weekly or more frequent use, stable from 2025 (62%; $p=0.847$), and 16% reported daily use, also stable from 2025 (24%; $p=0.361$).

Routes of Administration: Among participants who had recently consumed methamphetamine powder and commented ($n=51$), most (94%) participants reported injecting powder in the past six months, stable relative to 2025 (97%; $p=0.649$). Participants who reported injecting methamphetamine powder did so on a median of 33 days (IQR=6-90), stable relative to 2025 (36 days; IQR=10-120; $p=0.512$). One third (33%) reported smoking methamphetamine powder (30% in 2025; $p=0.839$).

Quantity: Of those who reported recent use and responded ($n=48$), the median amount of powder used on a 'typical' day in the past six months was 0.20 grams (IQR=0.10-0.30; 0.20

grams in 2025; IQR=0.10-0.30; $n=67$; $p=0.543$). The median maximum amount of powder used per day in the last six months was 0.50 grams (IQR=0.20-0.60; $n=47$; 0.30 grams in 2025; IQR=0.20-0.50; $n=66$; $p=0.068$).

Methamphetamine Base

Recent Use (past 6 months): Base has typically been the least commonly used form of methamphetamine and has gradually declined over time. In 2026, the lowest per cent reported recent use in the past 20 years (1%; 3% in 2025; $p=0.096$) (Figure 8). The per cent reporting recent use remained stable across all capital city samples.

Frequency of Use: Participants reported using methamphetamine base on a median of 12 days in the preceding six months (IQR=3-30; $n=13$; 6 days in 2025; IQR=2-14; $n=22$; $p=0.296$) (Figure 9).

Routes of Administration: Among participants who had recently consumed methamphetamine base and commented ($n=13$), injecting remained the most common route of administration (92%; 78% in 2025). Few participants ($n\leq 5$) reported smoking, swallowing or snorting. Due to small numbers reporting recent use, significance testing for routes of administration were not undertaken.

Quantity: Of those who reported recent use and responded ($n=13$), the median amount of base used on a 'typical' day in the past six months was 0.20 grams (IQR=0.20-0.50; 0.30 grams in 2025; IQR=0.10-0.50; $n=17$; $p=0.899$). The median maximum amount of base used per day in the last six months was 0.30 grams (IQR=0.20-0.50; $n=13$; 0.30 grams in 2025; IQR=0.10-1.00; $n=17$; $p=0.849$).

Methamphetamine Crystal

Recent Use (past 6 months): Reports of recent use of methamphetamine crystal more than doubled between 2009 (37%) and 2021 (78%), surpassing methamphetamine powder from 2011 onwards, but remaining relatively stable since 2021 (Figure 8). Indeed, in 2026, 79% of the sample reported recent use of methamphetamine crystal, stable from 2025 (79%; $p=0.903$). Recent use remained high and stable in all capital city samples (Table 6).

Frequency of Use: In 2026, median days of use remained stable at 82 days (IQR=24-180; $n=713$; 72 days in 2025; IQR=24-180; $n=685$; $p=0.412$), although this represents the highest frequency of use observed in the past 20 years (Figure 9). Almost four fifths (78%) of those who had recently used methamphetamine crystal reported weekly or more frequent use (78% in 2025; $p=0.845$), including 28% who reported daily use (26% in 2025; $p=0.431$).

Routes of Administration: Among participants who had recently consumed methamphetamine crystal and commented ($n=714$), the most common route of administration was injecting (96%; 95% in 2025; $p=0.442$), consistent with previous years. Participants who reported injecting did so on a median of 72 days in the preceding six months (IQR=20-175), stable relative to 2025 (72 days; IQR=24-170; $p=0.794$). Forty-six per cent reported smoking methamphetamine crystal in 2026, stable relative to 2025 (44%; $p=0.394$).

Quantity: Of those who reported recent use and responded ($n=693$), the median amount of crystal used on a 'typical' day of consumption in the past six months was 0.20 grams (IQR=0.10-0.30; 0.20 grams in 2025; IQR=0.10-0.30; $n=671$; $p=0.012$). The median maximum amount of crystal used per day in the last six months was 0.40 grams (IQR=0.20-0.70; $n=680$; 0.30 grams in 2025; IQR=0.20-0.70; $n=656$; $p=0.064$).

Table 5: Past six month use of methamphetamine powder, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	35	55	65	63	42	61	58	62
2008	38	37	64	61	34	61	50	35
2009	33	46	65	56	33	54	50	46
2010	29	48	53	56	29	51	25	41
2011	30	46	49	67	36	43	43	40
2012	17	42	39	70	34	45	46	30
2013	14	29	23	61	40	48	31	37
2014	17	36	25	50	34	39	16	31
2015	13	15	18	49	32	34	25	27
2016	17	18	9	33	19	18	24	27
2017	10	20	15	30	18	16	19	34
2018	11	23	16	22	31	12	17	34
2019	13	27	11	35	44	26	15	20
2020	11	13	10	43	35	36	-	19
2021	-	13	7	16	34	9	-	19
2022	11	-	5	18	31	13	-	11
2023	7	10	7	14	23	10	~	7
2024	12	9	-	13	15	-	-	10
2025	5	11	-	12	23	7	0	-
2026	-	12	-	9	15	-	-	6

Note. ~Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.005$. Please refer to Table 1 for a guide to table/figure notes.

Table 6: Past six month use of methamphetamine crystal, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	50	80	43	38	41	56	29	39
2008	69	68	39	32	49	61	28	40
2009	46	57	32	26	30	43	15	46
2010	48	48	36	20	60	40	18	37
2011	53	57	53	26	44	46	28	50
2012	68	66	59	43	56	64	26	44
2013	74	61	55	45	57	59	30	50
2014	74	72	75	54	60	53	26	58
2015	65	79	71	59	70	64	60	62
2016	77	78	73	73	73	75	62	69
2017	69	79	63	65	72	69	60	69
2018	76	85	77	76	79	64	74	70
2019	74	77	68	76	89	75	87	65
2020	75	63	64	77	80	69	83	63
2021	74	74	78	85	83	80	74	78
2022	87	81	75	84	83	77	80	70
2023	86	74	77	85	88	79	~	70
2024	82	83	70	89	92	65	87	85
2025	77	71	72	92	83	70	96	88
2026	75	81	69	90	87	70	89	78

Note. ~Due to the particularly small samples recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Price, Perceived Purity and Perceived Availability

Methamphetamine Powder

Price: The median price for one point (0.10 of a gram) of methamphetamine powder remained stable in 2026 (\$50; IQR=50-69; n=22; \$50 in 2025; IQR=50-50; n=35; $p=0.251$), as did the median price for one gram of methamphetamine powder (\$200; IQR=125-250; n=7; \$300 in 2025; IQR=200-300; n=9; $p=0.350$) (Figure 10).

Perceived Purity: Among those that were able to comment in 2026 (n=53), the perceived purity of methamphetamine powder significantly changed relative to 2025 ($p=0.049$). Two fifths (40%) of participants perceived purity to be 'medium', an increase from 28% in 2025, followed by 38% reporting 'high' purity, an increase from 25% in 2025 (Figure 12).

Perceived Availability: Among those were able to comment in 2026 (n=55), the perceived availability of methamphetamine powder remained stable relative to 2025 ($p=0.122$). In 2026, 62% perceived methamphetamine powder to be 'very easy' to obtain (43% in 2025), 16% perceived it to be 'easy' to obtain (24% in 2025) and 11% perceived it to be 'difficult' or 'very difficult' to obtain, respectively (23% and 10%, respectively in 2025) (Figure 14).

Methamphetamine Base

Questions pertaining to the price, perceived purity and perceived availability of methamphetamine base were not asked of participants from 2020 onwards. For historical information, please refer to the [2019 IDRS National Report](#).

Methamphetamine Crystal


Price: The median price for one point (0.10 of a gram) of methamphetamine crystal in 2026 was \$50 (IQR=50-50; n=295; \$50 in 2025; IQR=50-50; n=328; $p=0.828$). Across the years, the median price for one gram of crystal has fluctuated between \$250 and \$600. In 2026, the median price for one gram of crystal was \$300 (IQR=200-300; n=71), stable relative to 2025 (\$250; IQR=200-300; n=67; $p=0.516$) (Figure 11).

Perceived Purity: Among those that were able to comment in 2026 (n=644), the perceived purity of methamphetamine crystal remained stable relative to 2025 ($p=0.152$). Thirty-one per cent of participants perceived the purity of crystal to be 'medium' (34% in 2025), followed by 26% reporting 'high' purity (28% in 2025) and another 26% reporting 'low' purity (21% in 2025). Seventeen per cent reported 'fluctuating' purity (17% in 2025) (Figure 13).

Perceived Availability: Among those who were able to comment in 2026 (n=666), the perceived availability of methamphetamine crystal remained stable relative to 2025 ($p=0.939$). Seventy per cent of participants perceived that methamphetamine crystal was 'very easy' to obtain (70% in 2025), and a further 26% perceived it to be 'easy' to obtain (26% in 2025). Four per cent perceived methamphetamine crystal as 'difficult' to obtain (4% in 2025) (Please refer to Table 1 for a guide to table/figure notes.


Figure 15).

Figure 10: Median price of methamphetamine powder per point and gram, nationally, 2007-2026




Note. Among those who commented. The error bars represent the IQR. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 11: Median price of methamphetamine crystal per point and gram, nationally, 2007-2026




Note. Among those who commented. The error bars represent the IQR. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 12: Current perceived purity of methamphetamine powder, nationally, 2007-2026




Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 13: Current perceived purity of methamphetamine crystal, nationally, 2007-2026




Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 14: Current perceived availability of methamphetamine powder, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 15: Current perceived availability of methamphetamine crystal, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Past 6 months use

18%

5 Cocaine

Participants were asked about their recent (past six month) use of various forms of cocaine, including powder and crack/rock cocaine. Cocaine hydrochloride, a salt derived from the coca plant, is the most common form of cocaine available in Australia. 'Crack' cocaine is a form of freebase cocaine (hydrochloride removed), which is particularly pure. 'Crack' is most prevalent in North America and historically, has been infrequently encountered in Australia.

Median days*

3 days

*Among those whom reported past 6 month use in 2026:

Routes of administration

Snorting
65%

Injecting
43%

Smoking

Swallowing

Median Amount Used

Typical day:

0.30 grams

F
O
R
M
S

87% powder



21% crack/rock

Median Price

\$350
Per gram

Perceived Purity

n ≤ 5 fluctuates



Perceived Availability

Patterns of Consumption

Recent Use (past 6 months)

Recent use of cocaine declined from 22% in 2007 to 11% in 2016, before plateauing. In more recent years, reported use has increased slightly, with 18% reporting use in 2026 (19% in 2025; $p=0.538$) (Figure 16). The per cent reporting use in each capital city sample remained stable in 2026 (Table 7).

Frequency of Use

Frequency of use has remained low over the past two decades, with a median of three days of use observed in 2026 (IQR=1-7; $n=164$), stable relative to 2025 (2 days; IQR=1-6; $n=167$; $p=0.336$) (Figure 16). Of those who had recently used cocaine in 2026 and commented ($n=164$), 10% reported weekly or more frequent use, stable relative to 2025 (8%; $p=0.564$).

Routes of Administration

Among participants who had recently consumed cocaine and commented ($n=164$), snorting (65%; 64% in 2025; $p=0.905$) and injecting (43%; 43% in 2025; $p=0.908$) were the two most common routes of administration in 2026, consistent with previous years. Participants who reported injecting cocaine in the preceding six months had done so on a median of two days (IQR=1-8), stable relative to 2025 (3 days; IQR=1-6; $p=0.762$). The per cent who reported smoking cocaine remained low and stable at 6% (5% in 2025; $p=0.811$).


Quantity

Of those who reported recent use and responded ($n=134$), the median amount of cocaine used on a 'typical' day of consumption in the six months preceding interview was 0.30 grams (IQR=0.20-0.50; 0.20 grams in 2025; IQR=0.10-1.00; $n=102$; $p=0.598$).

Forms Used

Among participants who had recently consumed cocaine and commented in 2026 ($n=163$), the vast majority reported using powder cocaine (87%; 86% in 2025; $p=0.872$), with approximately one fifth (21%) reporting use of crack/rock cocaine (24% in 2025; $p=0.505$).

Figure 16: Past six month use and frequency of use of cocaine, nationally, 2007-2026



Note. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Secondary Y axis reduced to 10 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 7: Past six month use of cocaine, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	63	18	22	-	7	16	9	15
2008	58	18	24	-	-	15	-	13
2009	61	22	15	-	10	12	12	15
2010	57	6	14	-	12	15	-	13
2011	47	8	17	7	12	10	-	13
2012	44	16	9	11	7	15	-	-
2013	41	16	11	-	9	15	7	11
2014	32	15	10	8	7	7	-	9
2015	34	12	9	-	13	11	-	8
2016	25	8	10	6	6	10	-	9
2017	21	18	12	11	10	10	9	9
2018	26	14	15	11	10	12	6	9
2019	21	15	10	6	16	12	9	10
2020	23	19	17	16	14	18	-	19
2021	15	16	18	16	16	17	-	12
2022	16	17	19	14	10	12	12	17
2023	23	12	17	15	12	21	~	13
2024	29	18	14	18	13	9	14	20
2025	25	23	17	13	16	13	15	29
2026	25	24	18	16	17	11	13	18

Note. ~Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Price, Perceived Purity and Perceived Availability

Price

The median price for one gram of cocaine has fluctuated between \$300 and \$400 over the past two decades. In 2026, the median price of cocaine was \$350 for one gram (IQR=300-400; n=34), stable relative to 2025 (\$300; IQR=288-350; n=36; $p=0.088$). Few participants (n≤5) commented on the price of one point of cocaine in 2026 (\$50 in 2025; IQR=50-100; n=10) (Figure 17).


Perceived Purity

Among those who responded in 2026 (n=72), the perceived purity of cocaine remained stable relative to 2025 ($p=0.357$). Half (49%) perceived cocaine to be of 'high' purity (36% in 2025), while 22% perceived it to be of 'low' purity (32% in 2025) and 22% perceived it to be of 'medium' purity (24% in 2025) (Figure 18).

Perceived Availability


Among those able to comment in 2026 (n=71), the perceived availability of cocaine significantly changed relative to 2025 ($p=0.014$). Specifically, in 2026, more participants reported cocaine to be 'easy' (45%; 36% in 2025) or 'very easy' (34%; 26% in 2025) to obtain, while fewer participants perceived it to be 'difficult' (21%; 29% in 2025) or 'very difficult' (0%; 10% in 2025) to obtain (Figure 19).

Figure 17: Median price of cocaine per point and gram, nationally, 2007-2026




Note. Among those who commented. The error bars represent IQR. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 18: Current perceived purity of cocaine, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 19: Current perceived availability of cocaine, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; $p < 0.050$; $**p < 0.010$; $***p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Past 6 months
non-prescribed use

61%

6 Cannabis and/or cannabinoid-related products

Participants were asked about their recent (past six month) use of various forms of cannabis, including indoor-cultivated cannabis via a hydroponic system ('hydroponic'), outdoor-cultivated cannabis ('bush'), hashish, hash oil, commercially prepared edibles and CBD and THC extract.

Median days*

120 days

*Among those whom reported past 6 month non-prescribed use in 2026:

Routes of administration

Smoked

97%

Swallowing

Inhaled/vaped

Median Amount Used

Typical day:

1 gram

2 cones

1 joint

F
O
R
M
S

88% hydroponic

33% bush

8% hash/hash oil

5% CBD or THC extract

5% edibles

Median Price


Hydroponic:

\$ 20
Per gram


Bush:

\$ 20
Per gram

Hydroponic:
Perceived
Potency



10% fluctuates



Bush:
Perceived
Potency

Patterns of Consumption

From 2022, participants were asked about their use of both prescribed and non-prescribed cannabis and/or cannabinoid-related products. In 2026, 9% (n=79) of participants reported prescribed use in the six months preceding interview (7% in 2025; n=62; $p=0.255$), of which 97% reported using products that contained THC (not asked in 2025).

In the remainder of this chapter, data from 2021-2026, and between 2007-2016, refers to non-prescribed cannabis use only, while data from 2017-2020 refers to 'any' cannabis use (including hydroponic and bush cannabis, hashish and hash oil). While comparison between 2021-2026 and previous years should be treated with caution, the relatively recent legalisation of medicinal cannabis in Australia and the small percentage reporting prescribed use in 2022 (1%) and 2023 (3%) lends confidence that estimates are relatively comparable.

Recent Use (past 6 months)

Over the past two decades, recent use of non-prescribed cannabis and/or cannabinoid-related products has gradually declined. In 2026, 61% of the sample reported recent use, the lowest per cent observed in the past 20 years, although stable relative to 2025 (65%; $p=0.098$) (Figure 20). The per cent reporting non-prescribed use remained high and stable across most capital city samples, though a significant decrease was observed in the Sydney sample (59%; 71% in 2025; $p=0.026$) (Table 8).

Frequency of Use

In 2026, the median frequency of use in the past six months was 120 days (IQR=24-180; n=552), stable relative to 2025 (180 days; IQR=30-180; n=565; $p=0.168$) (Figure 20). Forty-five per cent of those who had recently used non-prescribed cannabis and/or cannabinoid-related products reported daily use (50% in 2025; $p=0.125$).

Routes of Administration

Among participants who had recently consumed non-prescribed cannabis and/or cannabinoid-related products and commented (n=554), smoking remained the most common route of administration (97%; 97% in 2025; $p=0.726$). Smaller percentages reported swallowing (8%; 6% in 2025; $p=0.141$) or inhaling/vaporising (6%; 4% in 2025; $p=0.052$) cannabis and/or cannabinoid-related products.


Quantity

Among those who reported recent use of non-prescribed cannabis and/or cannabinoid-related products and commented (n=526), the median 'typical' amount used on the last occasion of use was one gram (IQR=1.00-2.00; n=242; 1.00 gram in 2025; IQR=1.00-2.00; n=241; $p=0.338$) or two cones (IQR=1-4; n=211; 2 cones in 2025; IQR=2-4; n=216; $p=0.063$) or one joint (IQR=1-2; n=52; 1 joint in 2025; IQR=1-2; n=59; $p=0.376$).

Forms Used

Of those who had used non-prescribed cannabis and/or cannabinoid-related products in the past six months and commented ($n=526$), 88% reported recent use of hydroponic cannabis (86% in 2025; $p=0.367$), and one third (33%) reported recent use of outdoor-grown 'bush' cannabis, a significant decrease relative to 2025 (40%; $p=0.029$). Smaller percentages reported having used 'hashish' (5%; 3% in 2025; $p=0.216$), commercially prepared edibles (5%; 3% in 2025; $p=0.152$), hash oil (3%; 3% in 2025; $p=0.848$), non-prescribed THC extract (3%; 2% in 2025; $p=0.546$) or non-prescribed CBD extract (2%; 2% in 2025; $p=0.630$) (Figure 21).

Figure 20: Past six month use and frequency of use of non-prescribed cannabis and cannabinoid-related products, nationally, 2007-2026




Note. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Prior to 2021, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2017-2020 figures include some participants who were using prescribed cannabis only (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low (in 2022 few ($n \leq 5$) participants reported use of prescribed cannabis only). Further, from 2022, we captured use of 'cannabis and/or cannabinoid-related products', while in previous years questions referred only to 'cannabis'. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 8: Past six month non-prescribed use of cannabis and cannabinoid-related products, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	79	83	83	87	81	69	83	84
2008	80	80	74	86	75	64	78	82
2009	79	81	79	89	61	72	79	69
2010	72	81	81	79	66	70	72	77
2011	81	87	85	78	69	71	71	79
2012	72	81	85	81	61	79	71	70
2013	80	75	80	71	61	61	67	67
2014	77	74	75	82	75	69	62	70
2015	79	81	76	73	74	60	72	60
2016	76	69	77	74	73	70	72	64
2017	79	76	71	73	73	73	59	70
2018	76	79	70	81	70	77	60	67
2019	73	79	76	76	79	72	72	65
2020	64	77	69	72	67	66	60	64
2021	65	75	66	67	67	69	59	68
2022	72	77	82	70	72	60	70	64
2023	65	75	79	73	71	66	~	59
2024	68	74	69	74	76	60	69	58
2025	71	66	64	68	63	66	60	61
2026	59*	59	63	73	67	56	59	56

Note. ~Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Prior to 2021, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2017-2020 figures include some participants who were using prescribed cannabis only (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low. Further, in 2022, we captured use of 'cannabis and/or cannabinoid-related products', while in previous years questions referred only to 'cannabis'. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 21: Past six month use of different forms of non-prescribed cannabis and/or cannabinoid-related products, among those who reported recent non-prescribed use, nationally, 2018-2026



Note. Prior to 2021, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2018-2020 figures include some participants who were using prescribed forms of cannabis (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Price, Perceived Potency and Perceived Availability

Hydroponic Cannabis

Price: Consistent with previous years, the median price per gram of hydroponic cannabis was \$20 (IQR=20-25; n=123; \$20 in 2025; IQR=20-20; n=104; $p=0.186$). The price per ounce of hydroponic cannabis was \$270 (IQR=220-300; n=45), stable from 2025 (\$250; IQR=200-300; n=37; $p=0.584$) (Figure 22A).

Perceived Potency: Among those that were able to comment in 2026 (n=382), the perceived potency of non-prescribed cannabis remained stable relative to 2025 ($p=0.627$). Three fifths (62%) perceived hydroponic cannabis to be of 'high' potency (64% in 2025) followed by 28% perceiving it to be of 'medium' potency (26% in 2025). Five per cent perceived it to be of 'low' potency (4% in 2025) (Figure 23A).

Perceived Availability: Among those that were able to comment in 2026 (n=388), the perceived availability of non-prescribed cannabis remained stable relative to 2025 ($p=0.685$). Most participants perceived the availability of hydroponic to be 'very easy' (64%; 61% in 2025), followed by 'easy' (29%; 31% in 2025) (Figure 24A).

Bush Cannabis


Price: Consistent with previous years, the median price per gram of bush cannabis was \$20 (IQR=20-20; n=27; \$20 in 2025; IQR=20-20; n=39; $p=0.592$). The price per ounce of bush cannabis significantly decreased, from \$240 in 2025 (IQR=200-300; n=13) to \$178 in 2026 (IQR=150-200; n=14; $p=0.014$), the lowest price observed in the past 20 years (Figure 22B).

Perceived Potency: Among those who were able to comment in 2026 (n=98), the perceived potency of non-prescribed cannabis remained stable relative to 2025 ($p=0.295$). In 2026, 46% perceived bush to be of 'high' potency (40% in 2025) and 38% perceived it to be of 'medium' potency (39% in 2025) (Figure 23B).


Perceived Availability: Among those that were able to comment in 2026 (n=101), the perceived availability of non-prescribed cannabis remained stable relative to 2025 ($p=0.475$). Most participants perceived the availability of bush cannabis to be 'very easy' (52%; 58% in 2025), followed by 'easy' (36%; 27% in 2025) (Figure 24B).

Figure 22: Median price of non-prescribed hydroponic (A) and bush (B) cannabis per ounce and gram, nationally, 2007-2026

(A) Hydroponic cannabis




(B) Bush cannabis




Note. Among those who commented. Data from 2022 onwards refers to non-prescribed cannabis only; prior to 2022, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2017-2021 figures include some participants who reported on the price of prescribed cannabis (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low. The error bars represent the IQR. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 23: Current perceived potency of non-prescribed hydroponic (A) and bush (B) cannabis, nationally, 2007-2026

(A) Hydroponic cannabis




(B) Bush cannabis




Note. Among those who commented. Data from 2022 onwards refers to non-prescribed cannabis only: prior to 2022, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2017-2021 figures include some participants who are reporting on the potency of prescribed cannabis (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 24: Current perceived availability of non-prescribed hydroponic (A) and bush (B) cannabis, nationally, 2007-2026

(A) Hydroponic cannabis



(B) Bush cannabis



Note. Among those who commented. Data from 2022 onwards refers to non-prescribed cannabis only: prior to 2022, we did not distinguish between prescribed and non-prescribed cannabis, and as such it is possible that 2017-2021 figures include some participants who are reporting on the availability of prescribed cannabis (with medicinal cannabis first legalised in Australia in November 2016), although we anticipate these numbers would be very low. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.



Pharmaceutical Opioids

The following section describes recent (past six month) use of pharmaceutical opioids amongst the sample. Terminology throughout this chapter refers to prescribed use: use of pharmaceutical opioids obtained by a prescription in the person's name; non-prescribed use: use of pharmaceutical opioids obtained from a prescription in someone else's name; and any use: use of pharmaceutical opioids obtained through either of the above means. Contact the Drug Trends team (drugtrends@unsw.edu.au) for information on price and perceived availability of non-prescribed pharmaceutical opioids.

7

Past 6 month non-prescribed use in 2026:



Methadone




Buprenorphine Tablet: (Subutex)



Buprenorphine-Naloxone: (Suboxone)





Morphine:



Oxycodone:



Fentanyl:




Methadone

Any Recent Use (past 6 months): Over the past two decades, there has been a gradual decline in any methadone use, which has stabilised in recent years. In 2026, 27% of the sample reported any recent use, stable relative to 2025 (28%; $p=0.787$) (Figure 25). Historically, methadone use has largely consisted of prescribed use (22% in 2026; 21% in 2025; $p=0.956$) with the per cent reporting non-prescribed methadone use remaining low and stable in 2026, relative to 2025 (7%; 8% in 2025; $p=0.782$) (Figure 25). The per cent reporting non-prescribed use remained stable across all capital city samples in 2026, relative to 2025 (Table 9).

Frequency of Use: Frequency of non-prescribed methadone syrup use in the six months preceding interview remained stable (median 3 days in 2026; IQR=2-40; $n=57$; 6 days in 2025 (IQR=2-24; $n=61$) (Figure 25).

Recent Injecting Use: Of those who had recently used methadone syrup or tablets and commented ($n=243$), 28% reported injecting methadone, stable relative to 2025 (24%; $p=0.296$). Participants who reported injecting methadone did so on a median of 12 days (IQR=3-88), stable relative to 2025 (24 days; IQR=7-90; $p=0.149$).

Figure 25: Past six month use (prescribed and non-prescribed) and frequency of use of non-prescribed methadone, nationally, 2007-2026



Note. Includes methadone syrup and tablets except where otherwise specified. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Secondary Y axis reduced to 30 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 9: Past six month non-prescribed use of methadone, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	24	34	21	66	27	31	33	20
2008	27	35	21	70	17	19	45	27
2009	36	26	20	68	10	11	32	11
2010	27	25	19	58	17	13	27	15
2011	25	25	22	53	15	27	30	16
2012	26	27	21	47	14	31	27	12
2013	29	29	12	51	20	24	13	16
2014	29	27	21	51	9	20	16	17
2015	25	16	17	36	11	14	17	14
2016	21	12	13	40	6	13	14	19
2017	19	13	7	39	6	-	18	19
2018	20	13	11	42	-	9	8	18
2019	22	15	7	29	8	-	13	19
2020	17	7	10	26	9	11	-	20
2021	19	14	5	32	7	6	10	13
2022	18	8	7	16	-	9	-	10
2023	12	10	12	11	-	7	~	7
2024	12	6	5	10	-	7	-	8
2025	10	8	9	16	6	-	0	7
2026	13	8	5	8	-	-	-	13

Note. Includes methadone syrup and tablets. Due to the particularly small sample recruited in Darwin in 2023 ($n < 50$), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.


Buprenorphine Tablet

Any Recent Use (past 6 months): The per cent reporting recent use of any buprenorphine in tablet form has gradually declined over time. In 2026, 4% of the sample reported recent use of any buprenorphine tablet, the lowest per cent observed in the past 20 years, however, stable relative to 2025 (6%; $p = 0.053$) (Figure 26). Two per cent reported prescribed use (2% in 2025), and 2% reported non-prescribed use, a significant decrease relative to 2025 (4%; $p = 0.003$) (Figure 26). Recent non-prescribed use remained low across all capital city samples, with a significant decrease observed in the Sydney sample ($n \leq 5$; 6% in 2025; $p = 0.020$) and the Canberra sample (0%; 6% in 2025; $p = 0.014$) (Table 10).

Frequency of Use: Participants reported use of non-prescribed buprenorphine in tablet form on a median of 11 days in the six months preceding interview (IQR=5-54; $n = 16$; 6 days in 2025; IQR=3-69; $n = 35$; $p = 0.507$) (Figure 26).

Recent Injecting Use: Of those who had recently used buprenorphine in tablet form in 2026 and commented ($n = 36$), 56% reported recent injection, stable relative to 2025 (66%; $p = 0.375$). Participants who reported injecting buprenorphine tablet did so on a median of 10 days (IQR=6-72) in the six months preceding interview, also stable from 2025 (15 days; IQR=5-138; $p = 0.469$).

Figure 26: Past six month use (prescribed and non-prescribed) and frequency of use of non-prescribed buprenorphine tablet, nationally, 2007-2026



Note. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Secondary Y axis reduced to 40 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 10: Past six month non-prescribed use of buprenorphine tablet, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	16	28	26	6	11	19	-	31
2008	7	25	19	-	12	18	18	25
2009	18	23	25	12	9	16	-	31
2010	13	27	21	-	9	18	8	27
2011	12	21	18	6	8	11	8	33
2012	13	20	19	6	9	14	10	22
2013	11	16	9	9	7	10	20	16
2014	22	12	12	11	-	19	12	19
2015	9	11	12	13	6	8	10	17
2016	11	8	4	10	-	9	16	26
2017	13	14	6	9	7	10	-	25
2018	-	9	5	11	-	8	-	12
2019	4	-	-	-	0	-	-	15
2020	5	0	0	11	-	9	0	14
2021	5	-	-	11	-	-	-	20
2022	-	-	4	7	-	-	-	15
2023	5	-	-	20	-	-	~	11
2024	-	-	-	8	-	6	-	16
2025	6	6	-	7	-	0	0	9
2026	~*	0*	-	6	-	0	-	-

Note. ~Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.


Buprenorphine-Naloxone

Any Recent Use (past 6 months): After remaining relatively stable from 2009-2015, the per cent reporting any recent buprenorphine-naloxone use has been gradually declining. In 2026, 7% of the sample reported recent use of any buprenorphine-naloxone, a significant decrease relative to 2025 (11%; $p<0.001$) and the lowest percentage observed in the past two decades. Significant decreases were observed in both prescribed use of buprenorphine-naloxone (3%; 5% in 2025; $p=0.029$) and non-prescribed use (4%; 6% in 2025; $p=0.034$) (Figure 27). Recent non-prescribed buprenorphine-naloxone use remained low and stable in all capital city samples in 2026 (Table 11).

Frequency of Use: In 2026, participants reported using non-prescribed buprenorphine-naloxone on a median of four days (IQR=2-9; n=36) in the six months preceding interview, stable from three days in 2025 (IQR=2-21; n=54; $p=0.997$) (Figure 27).

Recent Injecting Use: Of those who had recently used buprenorphine-naloxone and commented in 2026 (n=61), 30% reported injecting as a route of administration, stable from 44% in 2025 ($p=0.098$). Participants reported injecting buprenorphine-naloxone on a median of four days (IQR=1-7) in the six months preceding interview (4 days in 2025; IQR=2-24; $p=0.395$).

Figure 27: Past six month use (prescribed and non-prescribed) and frequency of use of non-prescribed buprenorphine-naloxone, nationally, 2007-2026



Note. From 2007-2011 participants were asked about the use of buprenorphine-naloxone tablet; from 2012-2016 participants were asked about the use of buprenorphine-naloxone tablet and film; from 2017 onwards, participants were asked about the use of buprenorphine-naloxone film only. Median days of non-prescribed use computed among those who reported recent use (maximum 180 days) and is only reported from 2012 onwards to capture film use. Median days rounded to the nearest whole number. Secondary Y axis reduced to 30 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 11: Past six month non-prescribed use of buprenorphine-naloxone (any form), by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	-	6	13	-	-	15	-	24
2008	-	10	18	-	-	12	-	16
2009	6	11	14	-	9	28	8	22
2010	-	12	24	-	8	17	15	21
2011	8	12	29	-	-	14	14	11
2012 [#]	9	9	23	11	18	22	8	15
2013	9	11	17	9	9	22	19	22
2014	15	16	15	11	9	18	20	16
2015	11	12	17	13	15	19	22	27
2016	11	7	14	7	6	-	9	23
2017 [^]	14	13	11	14	14	16	10	24
2018	9	16	12	12	-	7	-	18
2019	11	14	10	7	8	16	10	22
2020	-	-	4	23	11	12	-	15
2021	-	9	5	21	10	13	-	11
2022	5	-	7	13	8	9	-	10
2023	8	-	6	18	8	9	~	15
2024	8	-	5	10	10	14	11	13
2025	5	-	6	11	9	8	-	7
2026	-	-	6	-	-	-	-	-

Note. # Includes 'tablet' and 'film' forms from 2012-2016. ^ Includes only 'film' form from 2017 onwards. ~ Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.


Morphine

Any Recent Use (past 6 months): The per cent reporting recent morphine use has been declining over time (Figure 28). In 2026, 8% of the sample had recently used any morphine (11% in 2025; $p=0.063$), representing the lowest percentage reporting recent use over the past two decades. In 2026, 7% reported recent use of non-prescribed morphine, a significant decrease from 9% in 2025 ($p=0.047$) and 2% reported recent prescribed use (2% in 2025; $p=0.713$). Non-prescribed morphine use remained stable across all capital city samples in 2026 compared to 2025 (Table 12).

Frequency of Use: Since peaking at a median of 26 days of use in 2014, an overall decline in the frequency of non-prescribed morphine use has been observed. In 2026, participants reported using non-prescribed morphine on a median of 12 days of (IQR=3-48; n=60), stable relative to 2025 (12 days; IQR=2-60; n=81; $p=0.908$) (Figure 28).

Recent Injecting Use: Of those who had recently used morphine and commented (n=73), three quarters (77%) reported injecting as a route of administration, stable relative to 2025 (81%; $p=0.569$). Those who reported injecting morphine did so on a median of 24 days (IQR=5-90) in the six months preceding interview, stable relative to 2025 (20 days; IQR=3-84; $p=0.703$).

Figure 28: Past six month use (prescribed and non-prescribed) and frequency of use of non-prescribed morphine, nationally, 2007-2026



Note. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Secondary Y axis reduced to 30 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 12: Past six month non-prescribed use of morphine, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	34	53	37	67	41	45	73	57
2008	31	35	40	81	30	31	85	51
2009	28	38	31	81	22	33	61	38
2010	31	36	30	73	24	28	89	38
2011	21	30	33	73	20	33	72	39
2012	21	30	27	64	23	43	69	34
2013	19	23	20	65	22	37	74	38
2014	25	12	24	71	20	27	80	32
2015	19	20	13	47	20	19	69	29
2016	16	12	10	51	18	16	71	33
2017	16	21	7	42	12	18	60	26
2018	17	10	10	47	7	14	54	29
2019	13	11	9	26	10	15	40	28
2020	7	8	8	38	11	18	32	21
2021	9	9	6	40	8	16	36	18
2022	11	10	7	27	10	7	27	19
2023	8	6	5	20	8	11	~	8
2024	9	8	-	26	9	6	14	11
2025	6	9	6	22	9	-	15	13
2026	-	7	-	13	7	-	18	6

Note. Due to the particularly small sample recruited in Darwin in 2023 ($n < 50$), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.


Oxycodone

Any Recent Use (past 6 months): The percentage of participants reporting recent oxycodone use peaked at 39% in 2012, gradually decreased from 2013 to 2019, and has since remained relatively stable (Figure 29). In 2026, 12% of the sample had recently used any oxycodone, stable relative to 2025 (14%; $p = 0.226$). Four per cent of the sample reported prescribed use (5% in 2025; $p = 0.419$), and 8% reported non-prescribed use (10% in 2025; $p = 0.241$). The per cent reporting recent non-prescribed oxycodone use remained stable in all capital city samples in 2026 compared to 2025 (Table 13).

Frequency of Use: In 2026, participants reported using non-prescribed oxycodone on a median of four days in the six months preceding interview (IQR=2-12; $n = 71$; 5 days in 2025; IQR=2-14; $n = 82$; $p = 0.532$) (Figure 29).

Recent Injecting Use: Of those who had recently used oxycodone and commented ($n = 105$), one third (36%) reported injecting as a route of administration, stable relative to 2025 (38%; $p = 0.779$). Participants reported injecting oxycodone on a median of seven days in the six months preceding interview (IQR=3-14), stable relative to 2025 (6 days; IQR=3-21; $p = 0.773$).

Figure 29: Past six month use (prescribed and non-prescribed) and frequency of use of non-prescribed oxycodone, nationally, 2007-2026



Note. From 2007-2015, participants were asked about recent use and frequency of use for any oxycodone; from 2016-2018, recent use and frequency of use for oxycodone was broken down into three types: tamper resistant ('OP'), non-tamper proof (generic) and 'other oxycodone' (median days non-prescribed use missing from 2016-2018). From 2019, recent use for oxycodone was broken down into four types: tamper resistant ('OP'), non-tamper proof (generic), 'other oxycodone' and oxycodone-naloxone, while frequency of use was asked for any oxycodone. From 2023, participants were asked about recent use and frequency of use for any oxycodone. Median days of non-prescribed use computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Secondary Y axis reduced to 14 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 13: Past six month non-prescribed use of oxycodone, by capital city, 2007-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2007	26	23	28	36	20	44	11	39
2008	27	27	25	53	15	23	28	26
2009	27	27	25	56	9	29	35	34
2010	33	13	28	60	17	20	22	26
2011	34	23	37	45	23	30	26	34
2012	46	34	26	56	26	48	19	29
2013	40	17	23	61	18	33	23	37
2014	40	16	22	47	21	27	22	38
2015	21	15	19	27	25	18	23	24
2016	23	12	10	28	16	17	18	22
2017	27	9	8	29	13	14	14	18
2018	16	10	10	28	-	15	11	18
2019	21	14	5	22	13	11	12	20
2020	9	8	7	24	11	8	9	15
2021	9	-	7	17	9	15	-	10
2022	11	6	10	12	10	11	-	11
2023	12	10	7	15	6	10	~	11
2024	9	9	5	21	8	11	0	16
2025	12	14	4	13	10	7	-	12
2026	11	14	-	15	8	-	-	9

Note. ~Due to the particularly small sample recruited in Darwin in 2023 (n<50), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.


Fentanyl

Any Recent Use (past 6 months): The per cent reporting any recent use of fentanyl has remained low and stable since monitoring commenced in 2013, with 4% reporting any recent use in 2026 (5% in 2025; $p=0.363$) (Figure 30). Three per cent reported non-prescribed use (4% in 2025; $p=0.072$) and 1% reported prescribed use (1% in 2025; $p=0.379$). Non-prescribed fentanyl use remained low and stable in all capital city samples in 2026 (Table 14).

Frequency of Use: In 2026, participants reported non-prescribed fentanyl use on a median of six days (IQR=2-13; n=24) in the six months preceding interview, stable relative to 2025 (3 days; IQR=1-12; n=37; $p=0.349$) (Figure 30).

Recent Injecting Use: Of those who had recently used fentanyl and commented in 2026 (n=36), 56% of participants reported injecting as a route of administration, stable relative to 2025 (75%; $p=0.100$). Participants reported injecting fentanyl on a median of three days (IQR=2-6) in the past six months (3 days in 2025; IQR=1-7; $p=0.870$).

Figure 30: Past six month use (prescribed and non-prescribed) and frequency of use of non-prescribed fentanyl, nationally, 2013-2026



Note. Collection of data on fentanyl use commenced in 2013; from 2013-2017, the IDRS did not distinguish between prescribed and non-prescribed use. Median days computed among those who reported recent use (maximum 180 days). Median days rounded to the nearest whole number. Secondary Y axis reduced to 10 days to improve visibility of trends. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 14: Past six month non-prescribed use of fentanyl, by capital city, 2018-2026

%	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
2018	6	6	8	0	-	8	-	16
2019	11	10	7	-	-	9	13	13
2020	8	9	-	-	10	11	-	-
2021	7	10	-	12	6	6	-	-
2022	4	9	-	10	6	-	0	-
2023	7	7	5	11	-	10	~	-
2024	7	14	-	-	-	-	-	-
2025	6	11	-	-	-	-	0	-
2026	5	-	-	-	-	0	-	-

Note. ~Due to the particularly small sample recruited in Darwin in 2023 ($n < 50$), data are not presented in this table. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Other Opioids

Participants were asked about prescribed and non-prescribed use of other opioids (Table 15). In 2026, 7% of participants reported any recent use of codeine (8% in 2025; $p = 0.652$), with 5% reporting prescribed use (4% in 2025; $p = 0.287$), and 3% reporting non-prescribed use (4% in 2025; $p = 0.052$). Of those who reported recent use ($n = 64$), few participants ($n \leq 5$) reported injecting as a route of administration (9% in 2025; $n = 67$; $p = 0.493$).

In 2026, 3% reported any recent use of tramadol, stable relative to 2025 (3%; $p = 0.890$). One per cent reported recent prescribed use (2% in 2025; $p = 0.573$) and 2% reported recent non-prescribed use, stable relative to 2025 (2%; $p = 0.593$). Of those who reported recent use ($n = 30$), few participants ($n \leq 5$) reported injecting as a route of administration in 2026 ($n \leq 5$ in 2025).

Two per cent of the sample reported recent use of tapentadol in 2026, stable relative to 2025 (2%; $p = 0.618$). One per cent of participants reported prescribed use (1% in 2025; $p = 0.782$) and 1% reported non-prescribed use (2% in 2025; $p = 0.684$). Of those who reported recent use in 2026 ($n = 17$), few participants ($n \leq 5$) reported injecting as a route of administration (30% in 2025; $n = 20$; $p = 0.725$).

Few participants ($n \leq 5$) reported any recent use of other opioids in 2026 ($n \leq 5$ in 2025) (not listed in Table 15).

Table 15: Past six month use of other opioids, nationally, 2019-2026

%	2019 (N=896)	2020 (N=880)	2021 (N=887)	2022 (n=878)	2023 (N=819)	2024 (N=881)	2025 (N=863)	2026 (N=900)
Codeine^								
Any use	9	10	10	9	10	8	8	7
Non-prescribed use	9	4	5	4	4	4	4	3
Any injecting use [#]	5	7	2	-	-	-	9	-
Tramadol								
Any use	16	7	8	10	9	6	3	3
Non-prescribed use	7	4	5	5	4	4	2	2
Any injecting use [#]	9	8	11	12	10	-	-	-
Tapentadol								
Any use	2	1	2	1	1	2	2	2
Non-prescribed use	1	-	-	1	0	1	2	1
Any injecting use [#]	-	-	0	0	0	0	30	-

Note. ^Includes high and low dose. [#]Of those who reported past six month use. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

8 Other Drugs

Participants were asked about their recent (past six month) use of various other drugs, including use of new psychoactive substances, non-prescribed use (i.e., use of a medicine obtained from a prescription in someone else's name) of other pharmaceutical drugs, and use of licit substances (e.g., alcohol, tobacco).

Past 6 month use in 2026 (median days among those that reported use):

New Psychoactive Substances (NPS) 4% 7 days	Non-prescribed benzodiazepines 21% 9 days	Non-prescribed pharmaceutical stimulants 5% 5 days	Non-prescribed antipsychotics 3%
Non-prescribed pregabalin 14% 6 days	Non-prescribed gabapentin 1% 4 days	Alcohol 48% 24 days	Tobacco 86% 180 days
Illicit e-cigarettes/vapes 28% 90 days	Nicotine pouches 4% 7 days	GHB/GBL/1,4-BD 16% 5 days	Unisom 3%

New Psychoactive Substances (NPS)

NPS are often defined as substances which do not fall under international drug control, but which may pose a public health threat. However, there is no universally accepted definition, and in practicality the term has come to include drugs which have previously not been well-established in recreational drug markets.

Recent Use (past 6 months): In 2026, 4% of the sample reported any recent NPS use, stable relative to 2025 (6%; $p=0.104$) (Table 16). 'New' drugs that mimic the effects of cannabis (2%; 2% in 2025; $p=0.851$), 'new' drugs that mimic the effects of opioids (1%; 2% in 2025; $p=0.088$), 'new' drugs that mimic the effects of amphetamines/cocaine (1%; 1% in 2025) and 'new' drugs that mimic the effects of psychedelics (1%; 1% in 2025) were the most commonly used NPS classes. In 2026, no participants reported use of 'new' drugs that mimic the effects of benzodiazepines, a significant decrease relative to 2025 (2%; $p<0.001$) (Table 16).

Table 16: Past six month use of new psychoactive substances, nationally, 2013-2026

%	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026
	N=887	N=898	N=888	N=877	N=888	N=905	N=902	N=884	N=887	N=870	N=818	N=883	N=865	N=902
'New' drugs that mimic the effects of opioids	/	/	/	/	-	-	2	1	1	1	1	1	2	1
'New' drugs that mimic the effects of ecstasy	/	/	/	/	1 [#]	1	2	-	1	1	-	1	1	-
'New' drugs that mimic the effects of amphetamine or cocaine	4	4	3	4	/	2	1	2	1	2	2	1	1	1
'New' drugs that mimic the effects of cannabis	9	8	8	8	5	5	6	5	4	2	3	2	2	2
'New' drugs that mimic the effects of psychedelic drugs	/	/	/	/	1 [#]	2	1	1	0	1	2	2	1	1
'New' drugs that mimic the effects of benzodiazepines	/	/	/	/	/	-	1	-	1	0	-	-	2	0***
'New' drugs that mimic the effects of dissociatives	/	/	/	/	/	/	/	/	/	/	/	/	-	-
Any of the above	12	11	10	11	8	11	11	8	7	6	7	5	6	4

Note. [#]In 2017, participants were asked about use of 'new drugs that mimic the effects of ecstasy or psychedelic drugs', thus the same value appears in both 'new' drugs that mimic the effects of ecstasy and 'new' drugs that mimic the effects of psychedelic drugs; from 2018 onwards, these two NPS classes were separated out. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Non-Prescribed Pharmaceutical Drugs

Benzodiazepines

Recent Use (past 6 months): Recent use of non-prescribed benzodiazepines has fluctuated over time, peaking at 53% in 2011, and declining thereafter. In 2026, one fifth (21%) of the sample reported recent use of non-prescribed benzodiazepines (e.g., Valium, Diazepam, Xanax, Kalma) in the six months preceding interview, stable relative to 2025 (23%; $p=0.188$) (Figure 31).

Frequency of Use: In 2026, participants reported using non-prescribed benzodiazepines on a median of seven days (IQR=3-24; $n=185$) in the six months preceding interview, stable relative to 2025 (10 days; IQR=3-30; $n=199$; $p=0.500$).

Recent Injecting Use: In 2026, 7% of participants who had recently used any non-prescribed benzodiazepines reported injecting as a route of administration (4% in 2025; $p=0.377$), and had done so on a median of two days in the past six months (IQR=1-9; 6 days in 2025; IQR=2-20; $p=0.208$).

Forms used: Among those who reported non-prescribed benzodiazepine use and responded in 2026 ($n=181$), the most commonly used benzodiazepine was Valium (diazepam) (61%), followed by Xanax (alprazolam) (30%), Diazepam (generic) (19%), Clonazepam (generic) (9%) and Serepax (oxazepam) (9%).

Pharmaceutical Stimulants

Recent Use (past 6 months): Non-prescribed use of pharmaceutical stimulants (e.g., Ritalin, dexamphetamine, Modafinil, Concerta, Vyvanse) gradually decreased between 2007 and 2020 before stabilising between 2020 to 2025 (Figure 31). In 2026, 5% reported recent use, the lowest per cent observed in the past 20 years and a significant decrease relative to 2025 (8%; $p=0.021$).

Frequency of Use: In 2026, participants reported using non-prescribed pharmaceutical stimulants on a median of nine days (IQR=2-24; $n=44$) in the six months preceding interview, stable relative to 2025 (5 days; IQR=2-10; $n=66$; $p=0.112$).

Recent Injecting Use: Almost one quarter (23%) of those who had recently used non-prescribed pharmaceutical stimulants reported injecting as a route of administration (18% in 2025; $p=0.615$) and had done so on a median of five days in the past six months (IQR=2-11; 4 days in 2025; IQR=1-41).

Antipsychotics

Recent Use (past 6 months): The per cent of the sample reporting recent use of non-prescribed antipsychotics (asked as 'Seroquel' from 2011-2018) gradually decreased between 2011 and 2020 before stabilising thereafter. In 2026, 3% reported recent use, stable relative to 2025 (3%; $p=0.777$) (Figure 31).

Frequency of Use: Frequency of non-prescribed use remained stable at five days in 2026 (IQR=2-20; $n=25$; 6 days in 2025; IQR=2-24; $n=26$; $p=0.570$).

Recent Injecting Use: No participants reported recently injecting antipsychotics in 2026 (0% in 2025).

Pregabalin

Recent Use (past 6 months): The per cent of the sample reporting recent use of non-prescribed pregabalin has remained relatively stable over the past decade, with 14% reporting recent use in 2026 (16% in 2025; $p=0.151$) (Figure 31).

Frequency of Use: Non-prescribed use was infrequent, with participants reporting use on a median of six days in the preceding six months (IQR=2-24; $n=124$), stable relative to 2025 (median 6 days; IQR=2-24; $n=140$; $p=0.790$).

Recent Injecting Use: Of those who had recently used non-prescribed pregabalin and commented ($n=124$), few participants ($n\leq 5$) reported injecting as a route of administration, stable relative to 2025 (8%; $p=0.119$).


Gabapentin

Recent Use (past 6 months): In 2026, 1% of the sample reported non-prescribed gabapentin use in the six months preceding interview (1% in 2025; $p=0.636$) (Figure 31).

Frequency of Use: Participants reported use on a median of four days in the preceding six months (IQR=2-8; $n=8$), stable relative to 2025 (median 6 days; IQR=2-15; $n=8$; $p=0.783$).

Recent Injecting Use: No participants reported recently injecting non-prescribed gabapentin ($n\leq 5$ in 2025).

Figure 31: Past six month use of non-prescribed pharmaceutical drugs, nationally, 2007-2026



Note. From 2019 to 2023, participants were asked about their use of non-prescribed alprazolam and non-prescribed use of 'other' benzodiazepines (e.g., diazepam), separately. In 2024, these categories were combined, and as such, participants were asked about non-prescribed use of any benzodiazepines. Antipsychotics was asked as 'Seroquel' from 2011-2018. Non-prescribed use of all these substances is reported. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Licit and Other Drugs

Alcohol

Recent Use (past 6 months): In 2026, 48% of the sample reported recent use of alcohol, the lowest per cent observed in the past 20 years, although stable relative to 2025 (53%; $p=0.060$) (Figure 32).

Frequency of Use: Participants who reported recent alcohol use in 2026 reported use on a median of 24 days (IQR=6-90; $n=433$), stable relative to 2025 (24 days; IQR=6-90; $n=456$; $p=0.836$), with 17% reporting daily use (16% in 2025; $p=0.710$).

Tobacco

From 2024, questions about illicit tobacco were included for the first time. Illicit tobacco was defined as products sold illegally without the necessary taxes added to the price.

Recent Use (past 6 months): Tobacco use has remained high over the past 20 years. In 2026, 86% of the sample reported recent use, stable from 2025 (87%; $p=0.622$) (Figure 32). Seventy per cent reported recent use of smoked or non-smoked illicit tobacco, a significant increase from 63% in 2025 ($p=0.009$).

Frequency of Use: Among those reporting recent use in 2026, frequency of use remained high and stable at a median of 180 days (i.e., daily; IQR=180-180; $n=779$; 180 days in 2025; IQR=180-180; $n=752$; $p=0.402$), with 88% reporting daily use (89% in 2025; $p=0.460$).

E-cigarettes/'Vapes'

[Legislation regulating e-cigarettes](#) (also known as vapes) has changed markedly in recent years. From October 2021, Australians were required to have a prescription to legally access nicotine containing e-cigarette products for any purpose, and from 1 July 2024, all e-cigarette products, regardless of whether they contained nicotine, could only legally be sold in a pharmacy. From 1 October 2024, people 18 years and older could buy e-cigarettes from participating pharmacies with a nicotine concentration of 20 mg/mL or less *without a prescription*, where state and territory laws allowed: products with a nicotine concentration of >20 mg/mL still required a prescription.

To capture these changes, in 2022, participants were asked for the first time about their use of both prescribed and non-prescribed e-cigarettes. From 2025, participants were asked about their use of e-cigarettes obtained from pharmacy (with or without a prescription) and 'non-pharmacy' locations.

In 2026, few participants (1%; $n=7$) reported recent use of e-cigarettes that were obtained from a pharmacy (1% in 2025; $n=8$; $p=0.796$). Between 2022 and 2024, few participants reported recent use of prescribed e-cigarettes ($n\leq 5$ in 2022; 1% in 2023 and 2024, respectively). The 2025-2026 data presented below refers only to use of e-cigarettes that were obtained from non-pharmacy locations, 2022-2024 data refers to non-prescribed e-cigarette use, while data from 2021 and earlier captures any e-cigarette use (collectively referred to as 'illicit use' from herein).

Recent Use (past 6 months): In 2026, 28% of the sample reported illicit e-cigarette use, stable relative to 2025 (25%; $p=0.167$) (Figure 32).

Frequency of Use: Among those reporting illicit use in 2026, frequency of use remained stable relative to 2025, with participants reporting a median of 90 days of use in the past six months (i.e., every

second day; IQR=14-180; n=247; 93 days in 2025; IQR=24-180; n=204; $p=0.519$), and 40% reporting daily use (42% in 2025; $p=0.702$).

Contents and Forms Used: Among those who reported recent illicit use and commented (n=249), the majority (94%) reported using disposable devices, followed by re-fillable devices (12%).

Reason for Use: Of those who reported *any* e-cigarette use and responded in 2026 (n=261), 36% reported that they used e-cigarettes as a smoking cessation tool (37% in 2025), however given differences in how the question was asked in 2026, significance testing was not undertaken. Other common reasons for use included 'addicted to nicotine' (32%), 'enjoyment of vaping/using e-cigarettes' (31%) and 'taste better than regular cigarettes' (21%).

Nicotine Pouches

Recent Use (past 6 months): Four per cent of the sample reported recent use of nicotine pouches in 2026, stable relative to 2025 (2%; $p=0.063$) (Figure 32).

Frequency of Use: Participants reported using nicotine pouches on a median of seven days (IQR=2-20; n=37) in the six months preceding interview, stable relative to 2025 (7 days; IQR=4-30; n=21; $p=0.553$).

Steroids

Recent Use (past 6 months): Reports of recent use of non-prescribed steroids have remained consistently low (between <1% and 3%) since monitoring commenced in 2010. Few participants (n≤5) reported recent use in 2026 (1% in 2025; $p=0.137$) (Figure 32).

GHB/GBL/1,4-BD

Recent Use (past 6 months): In 2026, 16% of the sample reported recent use of GHB/GBL/1,4-BD, stable relative to 2025 (17%; $p=0.437$) (Figure 32).

Frequency of Use: Participants reported use of GHB/GBL/1,4-BD on a median of five days in the preceding six months (IQR=2-16; n=140), stable relative to 2025 (5 days; IQR=2-20; n=147; $p=0.780$).

Recent Injecting Use: Of those who reported recent use (n=141), few participants (n≤5) reported injecting as a route of administration (n≤5 in 2025; $p=0.723$).

Forms Used: Among participants who had recently consumed GHB and commented (n=141), the majority reported using GHB (66%), with smaller proportions reporting use of GBL (10%) or 1,4-BD (9%). One fifth (22%) reported that they did not know what form they had used. This question was not asked in 2025.


Unisom

Unisom SleepGels is a Schedule 3 medicine containing diphenhydramine that is available over-the-counter from a pharmacist for use as an antihistamine or temporary sleep aid. It comes in a gel capsule formulation intended for oral use. There have been [reports](#) of injecting use in Australia, raising concern of attendant injecting-related injuries.

Recent Use (past 6 months): In 2026, 3% of the sample reported use of Unisom in the six months preceding interview (2% in 2025; $p=0.198$) (Figure 32).

Recent Injecting Use: Of those who had recently used Unisom and commented ($n=21$), 91% reported injecting as a route of administration (79% in 2025; $p=0.346$).

Figure 32: Past six month use of licit and other drugs, nationally, 2007-2026



Note. Data from 2022 onwards refers to illicit e-cigarettes only. This was defined as non-prescribed use in 2022-2024 and as use of e-cigarette products obtained from non-pharmacy locations in 2025-2026. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Drug-Related Harms and Other Behaviours

Participants were asked about various drug-related harms and other behaviours, including polysubstance use and bingeing, non-fatal overdose, AUDIT-C, naloxone awareness and uptake, equipment access and injecting behaviours, injection-related injuries and diseases, drug treatment, opioid and methamphetamine dependence, bloodborne virus testing and treatment, sexual health behaviours, mental health and psychological distress, health service access, driving under the influence of drugs, drug checking and crime. It should be noted that the following data refer to participants' understanding of these behaviours (e.g., may not represent medical diagnoses in the case of reporting on health conditions).

9

Polysubstance use
use of two or more drugs
the day before interview



63%

Non-fatal overdose
(past 12 months)

19%

Obtained Naloxone

Lifetime: 65%
past 12 months: 56%

Injection-related issues
(past 6 months)



28%

Mental health issues
(past 6 months)



53%

Drug treatment
(current)

37%

Health service access
(past 6 months)

87%

(for AOD support)

Hepatitis C test
(past 12 months)



55%

Driving under influence
(past 6 months)

8%
(among recent drivers)



77%

Drug checking
(past 12 months)

12%

Arrest
(past 12 months)

21%

Modes of purchasing
(past 12 months)


84%
(face-to-face)

Polysubstance Use

In 2026, the majority of the sample (92%) reported using one or more drugs (including alcohol and prescription medications, but excluding tobacco and e-cigarettes) on the day preceding interview. Of those who reported using one or more drugs and commented (n=833), the most commonly used substances were stimulants (54%), opioids (48%), cannabis (48%), alcohol (19%) and benzodiazepines (13%).

Sixty-three per cent of the sample reported using two or more drugs (including alcohol and prescription medications, but excluding tobacco and e-cigarettes) on the day preceding interview. Sixteen per cent reported concurrent use of cannabis and stimulants, and 10% reported concurrent use of opioids and cannabis, on the day preceding interview (Figure 33). Eighteen per cent of respondents reported using stimulants alone, 15% reported using opioids alone, and 8% reported using cannabis alone.

Figure 33: Use of opioids, stimulants, benzodiazepines and cannabis on the day preceding interview and most common drug pattern profiles, nationally, 2026




Note. % calculated out of total IDRS 2026 sample. The horizontal bars represent the per cent of participants who reported use of each drug class on the day preceding interview; the vertical columns represent the per cent of participants who used the combination of drug classes represented by the blue circles. Participants who did not report use of any of the four drug classes depicted are not shown in the figure but are counted in the denominator. 'Stimulants' includes methamphetamine, cocaine, ecstasy and pharmaceutical stimulants. 'Opioids' includes heroin, methadone, morphine, oxycodone, buprenorphine, buprenorphine-suboxone, fentanyl, other pharmaceutical opioids (codeine, tapentadol, tramadol, etc). Use of benzodiazepines, opioids and stimulants could be prescribed or non-prescribed use. Y axis reduced to 20% to improve visibility of trends.

Binge Drug Use

Participants were asked whether they had used any drug/s for 48 hours or more continuously without sleep (i.e., binged) in the six months preceding interview. The per cent of the sample who have reported bingeing has remained stable over time. In 2026, three fifths (60%) of the sample had binged on one or more drugs in the preceding six months, stable from 2025 (60%; $p=0.879$) (Figure 34).

Figure 34: Past six month use of drugs for 48 hours or more continuously without sleep ('binge'), nationally, 2018-2026



Note. Participants were first asked about bingeing in 2018. Statistical significance for 2025 versus 2026 is presented in figure; $*p<0.050$; $**p<0.010$; $***p<0.001$. Please refer to Table 1 for a guide to tables and figures.

Overdose Events

Non-Fatal Overdose

There have been some changes in the way questions about overdose have been asked over the years, which may account for some variation in estimates.

From 2019 onwards, participants were asked about their past 12-month experience of overdose where symptoms aligned with the examples provided and effects were outside their normal experience, or they felt professional assistance may have been helpful. We specifically asked about:

- **Opioid overdose** (e.g., reduced level of consciousness, respiratory depression, turning blue, collapsing and being unable to be roused). Participants who reported this experience were asked to identify all opioids involved in such events in the past 12 months;
- **Non-opioid overdose** (e.g., nausea, vomiting, chest pain, tremors, increased body temperature, increased heart rate, seizure, extreme paranoia, extreme anxiety, panic, extreme agitation, hallucinations). Drugs other than opioids were split into the following:
 - **Stimulant overdose:** Stimulant drugs include ecstasy, methamphetamine, cocaine, MDA, methylone, mephedrone, pharmaceutical stimulants and stimulant NPS (e.g., MDPV, Alpha PVP); and
 - **Other drug overdose:** 'Other drugs' include (but are not limited to) alcohol, cannabis, GHB/GBL/1,4-BD, amyl nitrite/alkyl nitrite, benzodiazepines and LSD.

It is important to note that overdose episodes reported across the drug types may not be unique given high rates of polysubstance use amongst the sample.

Each year we compute the total per cent of participants who have experienced any past 12-month overdose event by looking for any endorsement across the drug types queried (see Table 17).

Despite some fluctuation, the per cent reporting **any past 12 month non-fatal overdose** has remained relatively stable over the past two decades. In 2026, one fifth (19%) of the sample reported any past 12-month non-fatal overdose, unchanged relative to 2025 (19%) (Figure 35). Non-fatal overdose in the past 12-months remained stable in all capital city samples in 2026, relative to 2025.

Twelve per cent of the sample reported a **non-fatal overdose following opioid use** in the past 12 months (12% in 2025; $p=0.715$), whilst 6% reported a **non-fatal overdose following stimulant use** in the past 12 months (5% in 2025; $p=0.755$) (Table 17). The most commonly cited substance involved in past year non-fatal overdose was heroin (10%; 10% in 2025; $p=0.937$).


Participants who had overdosed on an opioid ($n=103$) had done so on a median of one occasion (IQR=1-2) in the last 12 months. Among those who had overdosed on an opioid in the past year and commented ($n=102$), 52% reported being administered naloxone, 34% reported that an ambulance had attended their most recent overdose, 28% were admitted to an emergency department and 15% reported receiving oxygen. One quarter (25%) reported not receiving any treatment. The most

commonly cited other drugs involved in participants' most recent opioid overdose were methamphetamine crystal (29%), benzodiazepines (23%), cannabis (22%) and alcohol (17%).

Among those who reported a non-fatal overdose in the past year and responded (n=72), 22% reported that they thought an opioid may have been present without their knowledge.

Please contact the Drug Trends team (drugtrends@unsw.edu.au) to request further findings regarding non-fatal overdose in the IDRS sample.

Figure 35: Past 12-month any non-fatal overdose, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 17: Past 12-month non-fatal overdose by drug type, nationally, 2025-2026, and by capital city, 2026

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=865	N=902	N=151	N=100	N=149	N=102	N=100	N=100	N=100	N=100
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
% Any opioid	N=856 12	N=893 12	N=146 14	N=100 19	N=149 16	N=102 -	N=97 6	N=100 11	N=99 -	N=100 14
% Heroin overdose	N=854 10	N=893 10	N=146 12	N=100 18	N=149 16	N=102 -	N=97 -	N=100 11	N=99 -	N=100 11
% Methadone overdose	N=854 -	N=893 -	N=146 -	N=100 0	N=149 0	N=102 -	N=97 0	N=100 0	N=99 0	N=100 -
% Morphine overdose	N=854 -	N=893 -	N=146 0	N=100 0	N=149 0	N=102 -	N=97 0	N=100 0	N=99 0	N=100 -
% Oxycodone overdose	N=854 1	N=893 -	N=146 0	N=100 0	N=149 0	N=102 0	N=97 0	N=100 0	N=99 0	N=100 -
% Stimulant overdose	N=856 5	N=890 6	N=144 -	N=100 14	N=149 -	N=102 -	N=98 8	N=100 -	N=99 -	N=99 14
% Other overdose	N=856 5	N=890 7	N=144 -	N=100 14	N=149 5	N=102 -	N=98 8	N=100 7	N=99 -	N=99 11
% Any drug overdose	N=859 19	N=894 19	N=146 19	N=100 35	N=149 18	N=102 8	N=98 22	N=100 20	N=99 8	N=100 27

Note. Participants reported on whether they had overdosed following use of the specific substances; other substances may have been involved on the occasion(s) that participants refer to. N is the number who responded (denominator). Statistical significance for 2025 versus 2026 is presented in table for national estimates; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Alcohol Use Disorders Identification Test-Concise (AUDIT-C)

The Alcohol Use Disorders Identification Test ([AUDIT](#)) was designed by the World Health Organization (WHO) as a brief screening scale to identify individuals with problematic alcohol use in the past 12 months. The AUDIT-C is a modified version of the 10 question AUDIT instrument, comprising three questions and is scored on a scale of 0-12.

In 2026, the mean score on the AUDIT-C for the total sample (including participants who had not consumed alcohol in the past 12 months) was 2.8 (SD 3.8), a significant decrease relative to 2025 (2.9; SD 3.7; $p=0.005$).

AUDIT-C scores of ≥ 4 (men) and ≥ 3 (women) are likely to indicate hazardous drinking, and potentially, alcohol dependence (Table 18). In 2026, one third (34%) of male participants had obtained a score of four or more (35% in 2025; $p=0.661$), and one third (33%) of female participants had obtained a score of three or more (39% in 2025; $p=0.155$), indicative of hazardous use. Both percentages for male and females were the lowest since monitoring commenced in 2010.

Table 18: AUDIT-C total scores and per cent of participants scoring above recommended levels, nationally, 2010-2026

	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026
	Men																
Mean AUDIT-C score (SD)	4.1 (3.9)	4.3 (3.9)	4.1 (3.9)	4.3 (4.0)	4.5 (4.0)	4.1 (3.9)	4.1 (3.9)	4.6 (3.6)	3.5 (3.6)	3.7 (3.8)	6.4 (3.5)	6.8 (3.4)	3.5 (4.0)	3.4 (3.8)	3.0 (3.9)	2.9 (3.6)	2.9 (3.9)
Score of ≥4 (%)	50	50	48	50	53	49	46	53	41	43	71	81	41	42	36	35	34
	Women																
Mean AUDIT-C score (SD)	3.2 (3.4)	3.6 (3.7)	3.5 (3.5)	3.6 (3.6)	3.9 (3.8)	3.4 (3.8)	3.4 (3.6)	4.0 (3.5)	3.5 (3.7)	2.7 (3.4)	5.3 (3.4)	5.7 (3.5)	3.2 (3.8)	3.0 (3.8)	2.7 (3.8)	2.8 (3.7)	2.5 (3.7)
Score of ≥3 (%)	45	51	50	49	53	44	47	54	49	36	74	77	44	39	35	39	33

Note. Monitoring of AUDIT commenced in 2010. Computed from the entire sample regardless of whether they had consumed alcohol in the past twelve months. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Naloxone Program and Distribution

Naloxone is a short-acting opioid antagonist that has been used for over 40 years to reverse the effects of opioids. In 2012, a take-home naloxone program commenced in the ACT (followed by NSW, VIC, and WA) through which naloxone was made available to peers and family members of people who inject drugs for the reversal of opioid overdose. In early 2016, the Australian Therapeutic Goods Administration (TGA) placed 'naloxone when used for the treatment of opioid overdose' on a dual listing of Schedule 3 and Schedule 4, meaning naloxone could be purchased OTC at pharmacies without a prescription, and at a reduced cost via prescription. From 1 December 2020 to 30 June 2022, under the take home naloxone pilot program, naloxone was made available free of charge and without a prescription in NSW, SA and WA. Following the evaluation of this pilot, the Australian Government announced that a national take home naloxone program was to be implemented in all Australian states and territories from 1 July 2022. Furthermore, naloxone nasal spray (Nyxoid) is now available in Australia as a PBS-listing, which is expected to increase use of naloxone in the community.

Awareness of Naloxone: From 2013-2026, at least four in five participants reported awareness of naloxone each year. In 2026, 84% of the sample reported that they had heard of naloxone, stable relative to 2025 (86%; $p=0.286$) (Figure 36). There was large variation across capital city samples, however, ranging from 42% among Darwin participants to 97% among Canberra and Melbourne participants (Table 19).

Awareness of Take-Home Naloxone: In 2026, four fifths (79%) of participants had heard about take-home naloxone, stable relative to 2025 (80%; $p=0.546$) (Figure 36). In 2026, 3% of the sample reported having heard of paid access (2% in 2025; $p=0.416$), and 78% of participants reported having heard of free access (79% in 2025; $p=0.854$). In 2026, knowledge regarding the take-home naloxone program was highest among the Melbourne (96%) and Canberra (95%) samples (Table 19).

Obtained Naloxone: In 2026, two thirds (65%) of the sample reported having obtained naloxone at least once in their lifetime, unchanged relative to 2025 (65%), with 56% having done so in the past year (55% in 2025; $p=0.774$) (Figure 37). Among participants who had ever obtained naloxone and responded ($n=572$), the majority reported most recently accessing naloxone from a Needle and Syringe Program (NSP) (64%; 63% in-person and $n\leq 5$ via post), followed by a pharmacy (10%), health service (7%) and the medically supervised injecting centre/room (7%). The majority (97%) reported that they did not have to pay the last time they obtained naloxone.

In 2026, of those who reported obtaining naloxone at least once in their lifetime ($n=576$), a median of two naloxone kits were stored away at the time of interview (IQR=1-3). Among those with at least one naloxone kit stored away and who responded ($n=392$), 96% had at least one kit within its expiration date, 7% had at least one kit that was expired, and 3% had both expired and non-expired kits.


Three per cent of the sample reported that they had tried to obtain naloxone in their lifetime but had been unsuccessful (3% in 2025), with 1% reporting this had occurred in the past year (1% in 2025; $p=0.507$). An additional 30% of participants reported that they had never tried to obtain naloxone

(29%; $p=0.675$) (note: a small per cent of participants reported never trying to obtain naloxone despite having obtained it in their lifetime – this could reflect that they had been given naloxone, but never actively sought it out). Out of those who had trouble obtaining naloxone or had never tried to obtain naloxone ($n=271$), the reasons included: 'don't use opioids' (18%), 'don't consider myself/my peers at risk of overdose' (15%) and 'not available in the pharmacy' (6%).

Of those who had ever obtained naloxone, had used opioids in the past month and responded ($n=489$), 53% reported that they 'always' had naloxone on hand when using opioids, followed by 19% reporting 'often', 15% 'sometimes', 5% 'rarely' and 8% 'never'.


Use of Naloxone to Reverse Overdose: In 2026, of those who responded ($n=891$), one third (36%) of the sample reported that they had resuscitated someone using naloxone at least once in their lifetime (32% in 2025; $p=0.065$), with 21% having done so in the past year. Six per cent reported that they had been resuscitated by a peer using naloxone in the past year, stable relative to 2025 (8%; $p=0.259$).

Figure 36: Lifetime awareness of naloxone and naloxone take-home programs, nationally, 2013-2026



Note. ^Wording of this question changed from 'Have you heard about take home naloxone programs' (after receiving a blurb about what these programs entailed: 2013-2022) to 'Are you aware that naloxone is available for people to take home' in 2023. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 37: Past 12 month obtainment of naloxone, nationally, 2020-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 19: Lifetime awareness of naloxone, and past year access, by capital city, 2026

	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=147	N=100	N=149	N=102	N=78	N=98	N=98	N=99
% Heard of naloxone	93	97	97	91	80	80	42	84
% Heard of take-home naloxone	88	95	96	85	64	76	39	78
% Obtained naloxone in past year	65	77	68	49	47	54	12	59

Note. N is the number who responded (denominator). Please refer to Table 1 for a guide to table/figure notes.

Equipment Access and Injecting Behaviours

Equipment Access

In 2026, participants reported obtaining a median of 100 new needle and syringes (IQR=30-200; 100 in 2025; IQR=25-200; $p=0.125$) and providing a median of 10 needles and syringes to others (IQR=0-50; 10 in 2025; IQR=0-50; $p=0.635$) in the past month. Participants reported having a median of 15 needles and syringes currently 'stored away' (IQR=2-80; 15 in 2025; IQR=2-58; $p=0.519$).

Fourteen per cent of the sample reported difficulties obtaining new needles and syringes in the past month (11% in 2025; $p=0.083$), 4% reported difficulties accessing filters (4% in 2025; $p=0.800$) and 10% reported difficulties obtaining sterile water (9% in 2025; $p=0.374$) (Table 20). The majority of participants reported that they obtained needles from an NSP (85%; 84% in 2025; $p=0.504$) in the past month, followed by a NSP vending machine (20%; 22% in 2025; $p=0.508$) and a friend/partner (13%; 13% in 2025; $p=0.880$) (Table 20).

Injecting Behaviours

In 2026, participants reported injecting on a median of 30 occasions in the past month (IQR=12-70), a significant change relative to 2025 (30 occasions; IQR=12-60; $p=0.017$). In 2026, 5% of the sample reported receptive sharing (5% in 2025) and 8% reported distributive sharing (8% in 2025; $p=0.857$) in the past month. The per cent who had shared injecting equipment other than syringes (e.g., spoons, tourniquet, water, and filters) in the month preceding interview more than halved between 2000 (51%) and 2011 (25%) and has remained relatively stable since (notwithstanding a sharp decline in 2019). In 2026, one fifth (20%) reported sharing other equipment (19% in 2025; $p=0.813$) (Table 21 and Figure 38). The per cent of the sample who reported re-using their own needles in the past month has also declined over time, with one third (33%) reporting reuse in 2026 (31% in 2025; $p=0.534$) (Table 21 and Figure 38). In 2026, 53% of the sample reported re-using other injecting equipment in the past month (55% in 2025; $p=0.495$) (Table 21), most commonly spoons and mixing containers (34%; 35% in 2025; $p=0.576$) and tourniquets (34%; 35% in 2025; $p=0.765$).

Thirty-six per cent of the 2026 sample reported that they had injected someone else in the past month, stable relative to 2025 (35%; $p=0.509$) and one fifth (20%) had been injected by someone else (21% in 2025; $p=0.591$) (Table 21).

The location of last injection remained stable between 2026 and 2025 ($p=0.500$). Consistent with previous years, most participants (80%) reported that they had last injected in a private home (80% in 2025) (Table 21). Eleven per cent of Melbourne participants and 5% of Sydney participants reported last injecting at the Medically Supervised Injecting Room or Centre (Table 21). Forty-five per cent of the national sample reported injecting alone on the last occasion of injecting (46% 2025; $p=0.629$).


The site of last injection remained stable between 2026 and 2025 ($p=0.808$). Most reported injecting in their arm on the last occasion of injecting (78%; 76% in 2025) followed by injecting in their hand/wrist (10%; 10% in 2025) (Table 21).

Table 20: Injecting equipment access in past month, nationally, 2025-2026, and by capital city, 2026

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=865	N=902	N=151	N=100	N=149	N=102	N=100	N=100	N=100	N=100
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
% Location of needle/syringe access past month	n=845	n=871	n=148	n=97	n=148	n=101	n=93	n=90	n=99	n=95
NSP	84	85	81	85	74	91	87	78	93	97
NSP vending machine	22	20	53	6	30	-	34	0	12	-
Posted from NSP	1	-	0	0	-	0	0	-	0	0
Chemist	11	9	7	9	-	18	10	9	-	19
Friend/Partner	13	13	7	10	11	13	23	22	8	11
Dealer	3	5*	5	-	-	-	16	-	-	-
Hospital	2	2	7	-	-	0	-	0	-	-
Outreach/peer worker	2	3	-	-	13	0	-	-	0	0
Medically supervised injecting Centre/Room	4	5	9	0	19	0	0	0	0	0
Other	3	3	-	23	-	0	-	-	0	0
% Difficulties accessing filters^ in the past month	4	4	11	-	-	-	-	-	-	-
% Difficulties accessing needles/syringes in past month	11	14	23	11	13	19	10	8	14	6
% Difficulties accessing sterile water in the past month	9	10	12	6	18	17	-	-	11	9
% Equipment used past month	n=860	n=897	n=151	n=99	n=149	n=102	n=98	n=99	n=99	n=100
Needle and syringe (e.g., 0.5mL, 1mL)	95	91**	93	81	91	90	93	97	92	90
Syringe or barrel (e.g., 3mL, 5mL, 10mL, 20mL, 50mL)	18	17	17	24	9	21	15	-	18	31
Spoons/mixing containers	65	62	92	84	87	29	47	59	12	55
Tourniquet	56	55	68	51	64	34	54	55	49	51
Swabs	85	83	91	82	89	63	90	80	82	82
Water	90	87	94	86	88	79	96	88	85	78
Any filters	63	60	81	76	80	31	41	60	29	59

Note. ^Filters included wheel filters, Sterifilt basic filters, Sterifilt plus filters and commercial cotton filters (e.g., Stericups). Statistical significance for 2025 versus 2026 is presented in figure for national estimates; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 38: Borrowing and lending of needles and sharing of injecting equipment in the past month, nationally, 2007-2026



Note. Borrowed (receptive): used a needle after someone else. Lent (distributive): somebody else used a needle after them. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 21: Injecting behaviours in the past month, and location and site of last injection, nationally, 2025-2026, and by capital city, 2026

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=865	N=902	N=151	N=100	N=149	N=102	N=100	N=100	N=100	N=100
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
% Injecting behaviours past month										
Borrowed a needle	n=854 5	n=890 5	n=146 5	n=98 10	n=149 4	n=101 7	n=98 -	n=99 -	n=99 7	n=100 -
Lent a needle	n=850 8	n=887 8	n=145 11	n=97 9	n=148 8	n=102 -	n=98 -	n=99 11	n=98 8	n=100 8
Shared any injecting equipment ^	n=860 19	n=896 20	n=150 18	n=98 14	n=149 34	n=102 10	n=98 18	n=99 30	n=100 15	n=100 13
Reused own needle	n=852 31	n=888 33	n=146 37	n=99 41	n=149 38	n=99 31	n=97 25	n=99 25	n=99 34	n=100 25
Reused any other equipment	n=857 55	n=888 53	n=149 56	n=97 52	n=149 67	n=98 33	n=98 64	n=98 51	n=99 55	n=100 42
Injected partner/friend ~	n=855 35	n=891 36	n=146 42	n=99 29	n=149 38	n=101 37	n=98 40	n=99 29	n=99 35	n=100 35
Somebody else injected them ~	n=854 21	n=891 20	n=146 21	n=99 13	n=149 19	n=101 19	n=98 28	n=99 16	n=99 26	n=100 19
% Location of last injecting use	n=849	n=891	n=146	n=99	n=149	n=102	n=98	n=98	n=99	n=100
Private home	80	80	75	89	65	80	88	83	83	90
Car	3	2	-	-	-	-	-	-	-	-
Street/car park/beach	8	7	11	-	13	8	-	-	7	-
Public toilet	6	5	-	-	7	11	-	9	-	-
Medically supervised injecting Centre/Room	3	3	5	0	11	0	0	0	0	0
Prison	0	0	0	0	0	0	0	0	0	0
Stairwell	-	-	-	-	0	0	0	0	-	0
Other	-	1	4	0	-	0	0	0	-	-
% Last injection site	n=855	n=890	n=145	n=99	n=149	n=102	n=98	n=98	n=99	n=100
Arm	76	78	74	86	70	84	88	78	78	68
Leg	6	5	5	-	5	7	0	8	6	-
Hand/wrist	10	10	8	6	15	-	7	10	14	15
Foot	2	1	-	-	-	-	-	-	-	-
Groin	1	1	-	-	-	0	0	0	0	-
Neck	5	5	9	-	7	-	-	-	-	7
Other	1	1	0	0	-	-	-	0	0	-

Note. Borrowed (receptive): used a needle after someone else. Lent (distributive): somebody else used a needle after them. ^Includes spoons, water, tourniquets and filters; excludes needles/syringes. ~With a new or used needle. N is the number who responded (denominator). Statistical significance for 2025 versus 2026 is presented in table for national estimates; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Self-Reported Injection-Related Injuries and Diseases

In 2026, 28% of the sample reported having an injection-related health issue in the month preceding interview, stable relative to 2025 (30%; $p=0.269$) (Table 22). The most common injection-related health issue reported by participants was any infection/abscess (12%; 14% in 2025; $p=0.156$), primarily skin abscess or cellulitis (10%; 12% in 2025; $p=0.096$), followed by any nerve damage (9%; 12% in 2025; $p=0.093$).

Table 22: Injection-related issues in the past month, nationally, 2025-2026, and by capital city, 2026

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=865	N=892	N=150	N=99	N=149	N=101	N=97	N=98	N=99	N=99
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
% Artery Injection	7	6	7	9	7	-	7	-	-	7
% Any nerve damage	12	9	10	11	15	-	13	-	-	15
% Any thrombosis	6	6	8	8	5	-	14	-	-	-
Blood clot	5	5	5	7	-	-	13	-	-	-
Deep vein thrombosis	2	2	5	-	-	-	-	0	0	-
% Any infection/abscess	14	12	21	19	15	6	7	-	-	10
Skin abscess	12	10	16	18	12	6	6	-	-	8
Other serious infection (e.g., sepsis, osteomyelitis)	3	2	4	-	-	0	-	0	0	-
Endocarditis	1	1	5	-	-	0	0	0	-	0
% Dirty hit	9	8	8	7	8	-	6	12	-	15
% Any injection related problem	30	28	35	37	33	16	32	19	12	33

Note. Statistical significance for 2025 versus 2026 is presented in table for national estimates; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Drug Treatment

In 2026, 37% of the sample reported that they were in any drug treatment for their substance use, stable from 38% in 2025 ($p=0.587$), with the most common treatment being methadone (21%; 20% in 2025; $p=0.591$), followed by buprenorphine depot injection (9%; 8% in 2025; $p=0.930$) (Table 23). Among those who reported oral methadone or buprenorphine treatment and commented ($n=215$), 76% reported receiving takeaway doses of methadone/buprenorphine (73% in 2025; $p=0.506$).

One tenth (11%) of participants reported having tried to access treatment in the past six months but being unable to, stable relative to 2025 (10%; $p=0.702$). Among those who had tried to access treatment but been unable to, most reported that, on the most recent occasion, they had been seeking treatment for their methamphetamine (55%; 55% in 2025) or heroin (35%; 33% in 2025; $p=0.875$) use. The most common service that participants had tried to access, but were unable to, were rehabilitation/therapeutic community (39%; 44% in 2025; $p=0.550$) and detoxification (26%; 19% in 2025; $p=0.289$), with the most common reasons being 'too hard to get into treatment (e.g., no places available, long waiting list)' (45%; 55% in 2025; $p=0.236$) and 'lack of support' (16%; 17% in 2025; $p=0.841$).

Table 23: Any current drug treatment, nationally, 2025-2026, and by capital city, 2026

	National		Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=865	N=902	N=151	N=100	N=149	N=102	N=100	N=100	N=100	N=100
	2025	2026	2026	2026	2026	2026	2026	2026	2026	2026
% Any current drug treatment	38	37	46	42	53	25	20	48	7	42
Methadone	20	21	26	26	30	6	16	32	-	20
Buprenorphine	2	1	-	0	0	-	0	0	0	-
Buprenorphine-naloxone	4	2*	-	-	-	-	-	-	-	-
Buprenorphine depot injection	8	9	15	7	17	-	-	10	-	-
Drug counselling	5	4	-	7	-	9	-	-	0	10
Other	3	4	-	8	-	7	-	0	-	11

Note. Statistical significance for 2025 versus 2026 is presented in table for national estimates; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Opioid and Methamphetamine Dependence

From 2015, participants were asked questions from the Severity of Dependence Scale (SDS) adapted to investigate opioid and methamphetamine dependence. The SDS is a five-item tool designed to screen for potential dependence on a variety of drugs. The SDS focuses on the psychological aspects of dependence, including impaired control of drug use, preoccupation with use, and anxiety about use. A total score was created by summing responses to each of the five questions. Possible scores range from 0 to 15.

To assess methamphetamine dependence in the past six months, a [cut-off value of four](#) was used, as this has been found to be a good balance between sensitivity and specificity for identifying dependent methamphetamine use. No validated cut-off for opioid dependence exists; however, researchers typically use a [cut-off value of five](#) as an indicator of likely dependence.

Of those who had recently used an opioid and commented (n=505), the median SDS score was seven (IQR=3-10), with two thirds (66%) scoring five or above, indicating possible dependence, a significant increase relative to 2025 (59%; $p=0.030$) (Table 24). Fourteen per cent of participants obtained a score of zero on the opioid SDS (15% in 2025; $p=0.587$), indicating no symptoms of opioid dependence.

Of those who had recently used methamphetamine and commented (n=694), the median SDS score was three (IQR=0-7), with 47% scoring four or above, indicating possible dependence, a significant decrease relative to 2025 (54%; $p=0.014$) (Table 24). One quarter (26%) of participants obtained a score of zero on the methamphetamine SDS, indicative of no symptoms of methamphetamine dependence (22% in 2025; $p=0.078$).

Table 24: Total opioid and methamphetamine SDS scores and per cent of participants scoring above cut-off scores indicative of dependence, among those who reported past six month use, nationally, 2015-2026

	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026
Opioid	N=690	N=674	N=659	N=659	/	/	N=553	N=530	N=527	N=530	N=500	N=505
Median total score (IQR)	7 (4-10)	7 (4-10)	7 (3-10)	5 (1-9)	/	/	6 (2-9)	5 (2-9)	5 (2-9)	6 (2-9)	6 (2-9)	7 (3-10)
% score = 0	8	9	12	25	/	/	14	16	17	15	15	14
% score ≥ 5	72	73	67	57	/	/	57	58	57	59	59	66*
Methamphetamine	N=616	N=639	N=609	N=680	/	/	N=700	N=666	N=638	N=696	N=674	N=694
Median total score (IQR)	3 (0-6)	3 (0-7)	3 (0-6)	1 (0-6)	/	/	3 (0-7)	3 (0-6)	3 (1-7)	3 (1-7)	4 (1-8)	3 (0-7)
% score = 0	32	31	31	44	/	/	28	26	25	22	22	26
% score ≥ 4	43	46	45	36	/	/	47	47	47	49	54	47*

Note. Severity of Dependence scores calculated out of those who used opioids/methamphetamine recently (past 6 months). A cut-off score of ≥5 and ≥4 is used to indicate screening positive for potential opioid and methamphetamine dependence, respectively. Imputation used for missing scale scores. Statistical significance for 2025 versus 2026 is presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Bloodborne Virus Testing and Treatment

In 2026, fifty-five per cent of participants reported that they had received a hepatitis C virus (HCV) antibody test in the past year, stable relative to 2025 (51%; $p=0.148$). Two fifths (43%) of participants had received a PCR or RNA test in the past year, stable relative to 2025 (39%; $p=0.121$) and 3% reported having a current HCV infection (4% in 2025; $p=0.261$) (

Table 25). Six per cent of the sample reported that they had received HCV treatment in the past year, stable relative to 2025 (5%; $p=0.659$), of which the majority (68%; $n=30$) reported that their treatment had been successful (63% in 2025; $p=0.638$). Eighty-six per cent of these participants reported having been re-tested with a PCR/RNA test to determine whether they had acquired a new HCV infection (re-infection) after successful treatment (96% in 2025; $p=0.355$).

Amongst those who had undergone an HCV RNA test in the last year and commented ($n=317$), almost one third (31%) reported it took more than five days to receive a result (whether positive or negative) following the administration of the last HCV RNA test, followed by 36% reporting 1-5 days, and another 28% reporting 0-24 hours.

Eighty-four per cent of those that commented ($n=838$), reported having had a test for human immunodeficiency virus (HIV) in their lifetime, a significant increase relative to 2025 (79%; $n=815$; $p=0.030$), with 31% reporting that the test occurred within the six months preceding interview (30% in 2025; $p=0.668$). Four per cent of those having had a test for HIV in their lifetime, reported a positive diagnosis (4% in 2025) (

Table 25).

Table 25: HCV and HIV testing and treatment, nationally, 2018-2026

	2018	2019	2020	2021	2022	2023	2024	2025	2026
%	N=905	N=902	N=884	N=888	N=879	N=820	N=884	N=865	N=902
Past year Hepatitis C test									
Past year hepatitis C antibody test	N=861 60	N=876 54	N=861 31	N=868 44	N=846 43	N=785 52	N=847 53	N=831 51	N=860 55
Past year hepatitis C PCR or RNA test	N=794 45	N=817 44	N=831 36	N=839 40	N=803 37	N=751 44	N=833 44	N=802 39	N=797 43
Current hepatitis C status									
Currently have hepatitis C [^]	N=807 20	N=823 15	N=836 11	N=826 9	N=805 7	N=737 7	N=808 4	N=780 4	N=780 3
Past year treatment for hepatitis C									
Received treatment in past year	N=852 18	N=794 15	N=854 9	N=862 12	N=835 10	N=749 8	N=830 6	N=799 5	N=789 6
Most recent treatment was successful (among those who had received treatment in past year)	N=99 94	N=79 97	N=80 72	N=100 69	N=85 69	N=63 75	N=47 77	N=41 63	N=44 68
Re-tested with a PCR or RNA test to determine re-infection (among those who underwent successful treatment)	/	/	/	/	/	N=43 77	N=36 81	N=26 96	N=29 86
HIV test				N=864	N=823	N=789	N=835	N=815	N=838
HIV test in past 6 months	/	/	/	31	23	33	30	30	31
HIV test more than 6 months ago	/	/	/	53	55	51	50	49	53
HIV status				N=722	N=633	N=669	N=672	N=643	N=694
Lifetime HIV positive diagnosis (among those who received HIV test)	/	/	/	4	3	3	3	4	4

Note. [^]This includes people who had not been tested for HCV. N is the number who responded (denominator). Timeframes for HCV and HIV differ; i.e., HCV questions focus on current and past year; HIV questions focus on lifetime and past six months. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Sexual Health Behaviours

In 2026, forty-five per cent of the sample reported some form of sexual activity in the past four weeks, stable relative to 2025 (49%; $p=0.177$) (Table 26). Given the sensitive nature of these questions, participants were given the option of self-completing this section of the interview (if the interview was undertaken face-to-face).

Amongst those who reported engaging in sexual activity in the past four weeks and commented ($n=381$), participants reported a median of one partner (IQR=1-2; median of 1 partner in 2025; IQR=1-2; $p=0.345$). Eight per cent reported engaging in sexual activity in the past four weeks in exchange for money, drugs, or other goods and services (10% in 2025; $p=0.447$) (Table 26).

Of those who commented ($n=839$), almost one quarter (24%) reported having a sexual health check-up in the six months prior to interview (22% in 2025; $p=0.323$), whilst 67% had done so in their lifetime (64% in 2025; $p=0.261$). Three per cent of the sample reported that they had received a positive diagnosis for a sexually transmitted infection (STI) in the past six months, stable relative to 2025 (2%; $p=0.514$) and 18% had received a positive diagnosis in their lifetime (17% in 2025; $p=0.512$) (Table 26). The most common STI reported amongst participants who commented ($n=17$) was chlamydia (47%), followed by gonorrhoea (41%), with few participants ($n\leq 5$) reporting other STIs. Information about HIV testing is provided in

Table 25.

Table 26: Sexual health behaviours, nationally, 2022-2026

	2022	2023	2024	2025	2026
Of those who responded[#]:	n=809	n=788	n=837	n=805	n=844
% Any sexual activity in the past four weeks	48	47	44	49	45
Of those who reported any sexual activity in the past four weeks and responded[#]:	/	/	n=362	n=387	n=372
% Engaged in sexual activity in exchange for money, drugs or other goods or services	/	/	10	10	8
Of those who responded[#]:	n=813	n=786	n=831	n=808	n=839
% Had a sexual health check in the last six months	17	24	23	22	24
% Had a sexual health check in their lifetime	60	64	61	64	67
Of those who responded[#]:	n=813	n=784	n=830	n=805	n=832
% Diagnosed with a sexually transmitted infection in the last six months	1	2	2	2	3
% Diagnosed with a sexually transmitted infection in their lifetime	15	17	18	17	18


Note. [#] Due to the sensitive nature of these items, there is missing data for some participants who chose not to respond. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Mental Health, Psychological Distress (K10), and Disability

Mental Health

In 2026, 53% of the sample self-reported that they had experienced a mental health problem in the preceding six months, stable relative to 2025 (55%; $p = 0.316$). Amongst those who had experienced a mental health problem, the most commonly reported problems were depression (61%; 63% in 2025; $p = 0.482$), anxiety (55%; 53% in 2025; $p = 0.583$) and post-traumatic stress disorder (PTSD) (34%; 32% in 2025; $p = 0.518$). Fewer participants reported attention deficit hyperactivity disorder (ADHD) (15%), schizophrenia (14%) and manic depression/bipolar disorder (13%). Twenty-seven per cent of the total sample (51% of those who reported a mental health problem) had seen a mental health professional during the past six months, stable from 2025 (27%; $p = 0.868$; 49% of those who reported a mental health problem; $p = 0.644$) (Figure 39). Seventy per cent of those who reported having seen a health professional about a mental health problem had been prescribed medication for their mental health problem in the preceding six months (69% in 2025; $p = 0.914$).

Figure 39: Self-reported mental health problems and treatment seeking in the past six months, nationally, 2007-2026



Note. The combination of the per cent who report treatment seeking and no treatment is the per cent who reported experiencing a mental health problem in the past six months. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Psychological Distress (K10)


The [Kessler Psychological Distress Scale 10 \(K10\)](#) was administered to obtain a measure of psychological distress in the past four weeks. It is a 10-item standardised measure that has been found to have good psychometric properties and to identify clinical levels of psychological distress as measured by the Diagnostic and Statistical Manual of Mental Disorders, and the Structured Clinical Interview for DSM disorders.

The minimum score is 10 (indicating no distress) and the maximum is 50 (indicating very high psychological distress). Scores can be coded into four categories to describe degrees of distress: scores from 10–15 are considered to indicate 'low' psychological distress; scores between 16–21 indicate 'moderate' psychological distress; scores between 22–29 indicate 'high' psychological distress; and scores between 30–50 indicate 'very high' psychological distress. Among the general population, scores of 30 or more have been demonstrated to indicate a high likelihood of having a mental health problem, and possibly requiring clinical assistance.

Among those who responded in 2026 (n=847), the per cent of participants scoring in each of the four K10 categories remained stable relative to 2025 ($p=0.526$), with 36% having a score of 30 or more (34% in 2025) (Figure 40).

The [National Health Survey 2022-23](#) provides Australian population data for adult (≥ 18 years) K10 scores. IDRS participants in 2026 reported greater levels of 'high' and 'very high' distress compared to the general population (Figure 40).

Figure 40: K10 psychological distress scores, nationally, 2007-2026, and among the general population, 2022-2023



Note. Data from the National Health Survey are a national estimate from 2022-23 for adults 18 or older. Imputation used for missing scale scores (IDRS only). Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Disability

In 2026, 53% of the sample self-reported that they identified as a person living with a disability (52% in 2025; $p=0.708$). Among participants who identified as a person with a disability and who commented in 2026 ($n=457$), the most common disability types were mental health (57%) and physical disability (38%), followed by neurodivergence (11%) and brain injury (10%) (**Error! Reference source not found.**).

In 2026, 12% of the entire sample reported that they had received support from the National Disability Insurance Scheme (NDIS) in their lifetime (12% in 2025; $p=0.878$), with 10% having done so in the past six months.

Table 27: Nature of disability, among those who self-identified as a person living with a disability, nationally, 2026, and by capital city, 2026


	National	Syd	Can	Mel	Hob	Ade	Per	Dar	Bri/GC
	N=457	N=83	N=58	N=85	N=51	N=40	N=40	N=43	N=57
% Nature of disability									
Physical disability (e.g., injury, amputation, rheumatoid arthritis)	38	36	36	33	39	40	48	40	37
Vision or hearing impairment	5	-	-	-	-	-	-	-	12
Mental health (excluding substance use disorder)	57	55	71	55	61	53	50	47	60
Neurodivergence (e.g., ADHD, ASD, dyslexia or other learning disabilities)	11	11	17	14	-	-	-	14	12
Brain injury	10	10	10	20	-	-	0	14	-
Intellectual disability	3	-	-	-	0	-	0	-	0
Pain-related neurological disorder (e.g., Complex Regional Pain Syndrome, Fibromyalgia)	5	-	-	-	0	-	-	-	-
Other neurological disorder (e.g., chronic fatigue, epilepsy, MS, Alzheimer's, Tourettes, cerebral palsy, Sneddon syndrome)	7	-	10	7	-	-	-	-	-
Other	2	0	0	-	-	-	0	0	-

Health Service Access

In 2026, 87% of participants reported accessing any health service for alcohol and/or drug (AOD) support in the six months preceding interview, stable relative to 2025 (90%; $p=0.053$) (Figure 41). The most common services accessed by participants for AOD support in 2026 were a NSP (77%; 80% in 2025; $p=0.123$), a general practitioner (GP) (35%; 41% in 2025; $p=0.012$) and a pharmacy (28%; 32% in 2025; $p=0.063$) (Table 28).

Nearly all participants (93%) reported accessing any health service for any reason in the six months preceding interview in 2026, although this was a significant decrease relative to 2025 (96%; $p=0.022$) (Figure 41). The most common services accessed by participants for any reason in 2026 were a NSP (80%; 84% in 2025; $p=0.077$), a GP (63%; 68% in 2025; $p=0.021$) and a pharmacy (49%; 53% in 2025; $p=0.150$) (Table 28).

Figure 41: Health service access for alcohol and other drug reasons, and for any reason in the past six months, nationally, 2020-2026



Note. Questions regarding health service access for AOD support were first asked in 2018, however due to differences in response options between 2018 and 2021, data are only presented from 2022 onwards. Statistical significance for 2025 versus 2026 is presented in figure; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.

Table 28: Types of health services accessed for alcohol and other drug reasons and for any reason in the past six months, nationally, 2022-2026

	AOD support					Any reason				
	2022	2023	2024	2025	2026	2022	2023	2024	2025	2026
% accessing health services	N=877 82	N=820 89	N=884 84	N=864 90	N=900 87	N=879 91	N=820 96	N=884 96	N=864 96	N=901 93*
GP	34	37	34	41	35*	61	66	64	68	63*
<i>In-person</i>	/	/	/	40	33**	/	/	/	67	61*
<i>Telehealth</i>	/	/	/	6	5	/	/	/	10	10
Emergency department	10	13	12	11	10	22	28	25	23	24
Hospital admission (inpatient)	8	9	9	8	9	17	20	20	19	21
Medical tent (e.g., at a festival)	0	-	-	-	-	1	1	-	1	-
Drug and Alcohol counsellor	17	16	13	14	14	17	17	14	15	14
Hospital as an outpatient	3	5	5	3	3	9	12	10	8	8
Specialist doctor (not including a psychiatrist)	5	3	2	2	2	10	10	8	7	8
Dentist	5	4	4	3	1	16	13	13	12	11
Ambulance attendance	6	7	7	6	6	11	14	14	13	10
Pharmacy	/	/	24	32	28	/	/	46	53	49
Other health professional (e.g., physiotherapist)	2	2	2	2	2	8	11	8	8	6*
Psychiatrist	6	5	5	4	3	11	11	9	7	6
Psychologist	6	4	6	5	3	11	9	10	8	6
NSP	69	76	74	80	77	71	80	82	84	80
Peer based harm reduction service	10	10	7	5	9**	11	11	7	6	10*
Other harm reduction service	6	7	8	7	8	6	7	9	8	9

Note. Statistical significance for 2025 versus 2026 is presented in table; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.


Driving

Thirty-six per cent of participants had driven a car, motorcycle or other vehicle in the last six months in 2026, stable relative to 2025 (37%; $p=0.545$). Of those who had driven within the last six months and commented in 2026 ($n=306$), 8% ($n=24$) reported driving while over the perceived legal limit of alcohol, stable relative to 2025 (9%; $p=0.771$). Among those who had driven within the last six months and commented ($n=309$), 77% ($n=239$) reported driving within three hours of consuming an illicit or non-prescribed drug, also stable relative to 2025 (77%; $p=0.921$) (Figure 42).

Of those who had driven within three hours of consuming an illicit or non-prescribed drug in the last six months and responded ($n=239$), participants most commonly reported using methamphetamine crystal (61%) prior to driving, followed by heroin (40%) and cannabis (32%).

Of those who had driven in the six months prior to interview and commented ($n=314$), 22% ($n=70$) reported that they had been tested for drug driving by the police roadside drug testing service (18% in 2025; $p=0.198$), and 29% ($n=91$) reported that they had been breath tested for alcohol by the police roadside testing service (28% in 2025; $p=0.788$) (Figure 42). Among those who had had been tested for drug driving by the police roadside drug testing service ($n=70$), 37% reported that a drug/s had been detected (45% in 2025; $p=0.460$), mostly commonly methamphetamine (34%), followed by cannabis (15%). Among those who reported that cannabis had been detected ($n=10$), no participants reported that this had been prescribed to them.

Figure 42: Self-reported testing, and driving over the (perceived) legal limit for alcohol or within three hours following illicit drug use, among those who had driven in the last six months, nationally, 2007-2026




Note. Computed of those who had driven a vehicle in the past six months. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Drug Checking

Drug checking is a common strategy used to test the contents and purity of illicit drugs. At the time interviewing commenced in 2026, the only government-sanctioned drug checking services that had operated in Australia were in the ACT, QLD, VIC and NSW. In Canberra, ACT, drug checking was provided at the Groovin the Moo festival in 2018 and 2019, and a fixed-site drug checking service (CanTEST) has been operational since 17 July 2022. Queensland's first fixed-site drug checking service, CheQpoint, opened in Brisbane on 20 April 2024, and a second service opened in the Gold Coast in July 2024. Drug checking services were also provided at 3 festivals in 2024 - Rabbits Eat Lettuce and Wildlands (by Pill Testing Australia) and Earth Frequency (by CheQpoint) - and as part of the 2024 Qld Gov Schoolies Response (CheQpoint). However, all government funded services ceased in April 2025. In Victoria, drug checking was provided at 'up to' 10 festivals throughout 2024-2025 during an 18-month implementation trial, followed by the establishment of a fixed site service in August 2025. In March 2025, NSW commenced a 12-month trial of mobile drug checking at 'up to' 12 festivals. Following the trial, the NSW Government announced that drug checking would become a permanent service, with testing to be provided at up to 15 festivals annually from October 2026 onwards.

In 2026, 12% of participants reported that they or someone else had tested the contents and/or purity of their illicit drugs in Australia in the past year, stable relative to 2025 (15%; $p=0.114$) (Figure 43). Of those who reported that they or someone else had tested their illicit drugs in the past year in 2026 and responded ($n=109$), 59% reported using a personal testing kit, most commonly colorimetric or reagent test kits (28%), followed by testing strips (e.g., BTNX fentanyl strips or other immunoassay testing strips) (27%). Of those who reported that they or someone else had tested their illicit drugs in the past year ($n=109$), 49% reported that they had submitted drugs for testing at a drug checking service – most commonly at a fixed-site face-to-face drug checking service (e.g., a drop-in service in a central location) (48%) Few participants ($n\leq 5$) reported using an event-based face-to-face testing service (e.g., festival pill-testing service) or submitting samples via a postal/online testing service (e.g., Energy Control, Ecstasy Data).

Figure 43: Past year engagement in drug checking, nationally, 2019-2026



Note. Questions on drug checking commenced in 2019. From 2025, survey questions were separated into 'personal testing kits' and 'drug checking services' and focused on past year use only. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Experience of Crime and Engagement with the Criminal Justice System

Past month self-reported criminal activity has remained relatively stable over the past two decades, ranging between 36% and 45%. In 2026, 'any' past month crime remained stable relative to 2025 (43%; 44% in 2025; $p=0.585$) (Figure 44).


Property crime (29%) and selling drugs for cash profit (23%) remained the most common forms of criminal activity in 2026, stable relative to 2025 (29%; $p=0.909$ and 24%; $p=0.602$, respectively). Fewer participants reported past month violent crime (5%; 4% in 2025; $p=0.570$) or fraud (3%; 4% in 2025; $p=0.235$) (Figure 44). Being the victim of a crime involving violence (e.g., assault) in the month preceding interview remained stable in 2026, relative to 2025 (15%; 17% in 2025; $p=0.263$) (Figure 45).

Fifty-seven per cent of those that commented ($n=882$) reported a lifetime prison history in 2026, stable relative to 2025 (60%; $p=0.332$) (Figure 46).

One third (33%) of participants reported a drug-related encounter with police which did not result in charge or arrest, stable relative to 2025 (29%; $p=0.138$) (Figure 46). This predominantly comprised being stopped and searched (60%; 67% in 2025; $p=0.155$), followed by stopped and questioned (53%; 48% in 2025; $p=0.263$). Thirty-one per cent were asked to move along (30% in 2025), 29% were stopped and issued a court attendance notice (26% in 2025; $p=0.487$), and one fifth (22%) were stopped and issued a caution (24% in 2025; $p=0.530$).


In 2026, one fifth of participants (21%) reported past year arrest, stable from 2025 (22%; $p=0.553$) (Figure 46). Of those who had been arrested and commented in 2026 ($n=175$), the main reasons for arrest were use/possession of drugs (25%), followed by property crime (22%). In 2026, 10% of the sample had been convicted of a drug-related offence in the past year (9% in 2025; $p=0.866$), and 10% had been sentenced to a community corrections order (9% in 2025; $p=0.616$).

Figure 44: Self-reported criminal activity in the past month, nationally, 2007-2026




Note. 'Any crime' comprises the per cent who report any property crime, drug dealing, fraud and/or violent crime in the past month. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 45: Victim of crime involving violence in the past month, nationally, 2019-2026



Note. Questions regarding being the victim of a crime involving violence were first asked in 2019. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Figure 46: Lifetime incarceration, and past 12 month arrest and drug-related encounters with police that did not result in arrest, nationally, 2007-2026



Note. Statistical significance for 2025 versus 2026 is presented in figure; * $p < 0.050$; ** $p < 0.010$; *** $p < 0.001$. Please refer to Table 1 for a guide to table/figure notes.

Modes of Purchasing Illicit or Non-Prescribed Drugs

In interviewing and reporting, 'online sources' were defined as either surface or darknet marketplaces.

Purchasing Approaches

In 2026, the most popular means of arranging the purchase of illicit or non-prescribed drugs in the 12 months preceding interview was face-to-face (84%), a significant increase relative to 2025 (79%; $p=0.019$) (Table 29). This was followed by phone call (50%; 52% in 2025; $p=0.415$) and text messaging (32%; 30% in 2025; $p=0.404$). Twelve per cent reported using social networking or messaging applications (e.g., Facebook, Wickr, WhatsApp, Snapchat, Grindr, Tinder) (11% in 2025; $p=0.544$). It is important to re-iterate that this refers to people *arranging the purchase* of illicit or non-prescribed drugs. This captures participants who messaged friends or known dealers on Facebook Messenger or WhatsApp, for example, to organise the purchase of illicit or non-prescribed drugs, which may have then been picked up in person.

Table 29: Purchasing approaches in the past 12 months, nationally, 2022-2026

%	2022	2023	2024	2025	2026
% Purchasing approaches in the last 12 months ^{^#}	N=855	N=809	N=873	N=853	N=881
Face-to-face	74	79	85	79	84*
Surface web	1	1	1	1	-
Darknet market	2	1	2	1	1
Social networking or messaging applications [`]	11	11	13	11	12
Text messaging	37	34	35	30	32
Phone call	62	56	54	52	50
Grew/made my own	/	/	2	2	2
Other	/	/	-	-	-

Note. [^] participants could endorse multiple responses. [#]This refers to people *arranging the purchase* of illicit or non-prescribed drugs. [`] This captures participants who messaged friends or known dealers on Facebook Messenger or WhatsApp, for example, to organise the purchase of illicit or non-prescribed drugs, which may have then been picked up in person. Statistical significance for 2025 versus 2026 presented in table; * $p<0.050$; ** $p<0.010$; *** $p<0.001$. Please refer to Table 1 for a guide to table/figure notes.