

2022 SAMPLE CHARACTERISTICS



In 2022, 102 participants, recruited from Brisbane/Gold Coast, QLD were interviewed.



The median age in 2022 was 23, and 53% identified as male.



In the 2022 sample, 51% were enrolled students, 30% were employed full time and 11% were unemployed.



Ecstasy



Cocaine



Other stimulants

Participants were recruited on the basis that they had consumed ecstasy and/or other illicit stimulants at least monthly in the past 6 months.

PAST 6 MONTH USE OF SELECTED DRUGS

Ketamine


LSD

Hallucinogenic mushrooms

GHB/GBL/ 1,4-BD Amyl nitrite

Nitrous oxide (nangs)

Non-prescribed e-cigarettes




MENTAL HEALTH AND SEXUAL HEALTH BEHAVIOURS



MH issue



Seen a MH professional



Of those who had a mental health condition, the three most common mental health issues reported were anxiety (70%), depression (70%) and PTSD (14%).




Reported sexual activity



STI test

In the total sample, 88% reported sexual activity in the past 4 weeks, and 45% had a sexual health check in the past 6 months.



Sexual risk behaviours among those who reported any sexual activity in the past four weeks and were able to comment

OTHER RISK BEHAVIOURS



In the total sample, 75%

self-reported a mental health issue

and 50% had seen a mental health


professional in the past 6 months.

Among recent drivers, 55% reported driving a vehicle within 3 hours of consuming illicit drugs and 29% while over the legal limit of alcohol.





In the 2022 sample, 36% reported a non-fatal depressant overdose in the previous 12 months, relatively stable to 2021 (22%).




In the total sample, 82% reported concurrent use of two or more substances on the last occasion of ecstasy or related drug use.




stimulants and depressants Stimulants, depressants and cannabis

The most commonly used combinations of drug classes on the last occasion of ecstasy or related drug use.

ECSTASY



Past 6 month use of ecstasy capsules, crystal, pills, and powder in 2022.



Of those who had recently consumed ecstasy, 15% reported weekly or more frequent use.



2 Capsules 2 Pills

0.30 grams of crystal 0.30 grams of powder

Median amounts of ecstasy consumed in a 'typical' session using each form.




In 2022, more participants perceived the availability of capsules and crystal forms as 'difficult' or 'very difficult' relative to 2021.


METHAMPHETAMINE




Past 6 month use of any methamphetamine, crystal, powder and base in 2022.



Of those who had recently consumed methamphetamine, n≤5 reported weekly or more frequent use.




All participants who had recently used crystal smoked it. Of those who had recently used powder, 60% snorted it.




Of those who could comment, 88% perceived crystal methamphetamine to be 'easy' or 'very easy' to obtain.

COCAINE




Past 6 month use of any cocaine remained stable between 2021 and 2022.



Of those who had recently consumed cocaine, 16% reported weekly or more frequent use.



In 2022, the median price of a gram of cocaine remained stable at \$350.




Of those who could comment, 81% perceived cocaine to be 'easy' or 'very easy' to obtain.

CANNABIS AND/OR CANNABINOID REL







Past 6 month use of non-prescribed cannabis and/ or cannabinoid related products remained stable between 2021 and 2022.



Of those who had recently consumed non-prescribed cannabis and/or cannabinoid related products, 67% reported weekly or more frequent use.



Of participants who had consumed non-prescribed cannabis and/or cannabinoid related products in the last 6 months, 94% had smoked it.



Of those who could comment, 92% perceived hydro to be 'easy' or 'very easy' to obtain.