
2021 SAMPLE CHARACTERISTICS



In 2021, 100 people from Adelaide, SA, participated in EDRS interviews.



The median age in 2021 was 25 (IQR = 21 - 32), and 57% identified as male.



In the 2021 sample, 41% were enrolled students, 29% were unemployed, and 20% were employed full time.




Participants were recruited on the basis that they had consumed ecstasy or other illicit stimulants at least monthly in the past 6 months.

PAST 6 MONTH USE OF OTHER DRUGS




MENTAL HEALTH AND SEXUAL HEALTH BEHAVIOURS




34% Seen a MH professional

65% Anxiety PTSD


In the total sample, 57% self-reported a mental health issue and 34% had seen a mental health professional in the past 6 months.



Of those who commented, the top three most common mental health issues reported were anxiety (70%), depression (65%) and PTSD (25%).




In the total sample, 82% reported sexual activity in the past 4 weeks, and 33% had a sexual health check in the past 6 months.




Sexual risk behaviours among those who reported any sexual activity in the past four weeks (82%) and were able to comment.


OTHER RISK BEHAVIOURS



In the total sample, 38% reported driving a vehicle within 3 hours of consuming illicit drugs and 23% while over the legal limit of alcohol.



The most common drugs used prior to driving were cannabis (63%) and crystal methamphetamine (29%).



In the total sample, 86% reported concurrent use of two or more substances on the last occasion of ecstasy/stimulant use.




depressants


Stimulants & depressants & cannabis

In the total sample, 26% reported to have used stimulants and depressants on one occasion whereas 17% reported using stimulants, depressants and cannabis.


ECSTASY




Past 6 month use of ecstasy capsules, crystal, pills, and powder in 2021.



Of those who had recently consumed ecstasy, 17% used it weekly or more frequently.




Median amounts of ecstasy consumed in a 'typical' session using each form.




Of those who could comment 75% perceived ecstasy capsules to be 'easy' or 'very easy' to obtain.

METHAMPHETAMINE



Past 6 month use of any methamphetamine (33%), crystal (32%), powder (n≤5) and base (0%) in 2021.




Of those who had recently


consumed methamphetamine,

70% used it weekly or more

frequently.




94% of people who had recently used crystal smoked it. Of those who had recently used powder, 100% snorted it.



Of those who could comment 89% perceived crystal methamphetamine to be 'easy' or 'very easy' to obtain.


COCAINE




Past 6 month use of any cocaine remained stable from 2020 (69%) to 2021 (78%).



Of those who had consumed cocaine recently, 10% reported weekly or more frequent use.




Of those who had consumed cocaine in the last 6 months, all particiants had snorted it (99%).




Of those who could comment 80% perceived cocaine to be 'easy' or 'very easy' to obtain.

CANNABIS




Past 6 month use of any cannabis remained stable from 89% in 2020 to 84% in 2021



Of those who had consumed cannabis recently, 74% reported weekly or more frequent use.



Of people who had consumed cannabis in the last 6 months, 96% had smoked it



Of those who could comment 93% perceived hydro to be 'easy' or 'very easy' to obtain.