

The 2016 Sydney Gay Asian Men Survey is a cross sectional research project designed to provide an updated understanding of key HIV-related behaviours and practices amongst gay men of Asian background in Sydney. It also provides a comprehensive assessment of their ongoing HIV prevention needs.

This research was led and sponsored by the Sydney-based Culturally and Linguistically Diverse (CALD) Gay Men Action Group; a partnership between ACON, the Multicultural HIV and Hepatitis Service (MHAHS), South Eastern Sydney LHD, Sydney LHD, South Western Sydney LHD, Western Sydney LHD, Northern Sydney LHD and the Centre for Social Research in Health (CSRH) at UNSW.

The data presented here was collected from September 2015 and June 2016 at community and clinical venues across Sydney and also online.



PARTICIPANTS 468

Total number of participants



Chinese 39.7% Filipino 12.6%

Thai **12.4%** Indian **7.1%**

Other Asian Backgrounds 28.2%









GENDER 97.2% Male (cisgender) — 2.8% Male (transgender)



SEXUALITY 88.7% 9.2% 2.1% Other





RELATIONSHIP STATUS IN THE **LAST 6 MONTHS** 17.9% Regular male partner only (monogamous)

31.6% Regular and casual male partners

35% Casual male partners only

15.4% No sexual relationships



TOP 3

WHERE MEN **MEET SEX PARTNERS**

Mobile Apps

69.2% 47.6%

Internet

Gay Sauna





B DRUGS

MOST FREQUENTLY USED DRUGS IN THE LAST 6 MONTHS (%)

Alcohol Amyl Marijuana

71.6% 38.5% 14.7%

Viagra Ecstacy Crystal

14.3% 10.0% 14.7%

GHB 5.8% Steroids 1.3%



* ANAL

ANAL INTERCOURSE AMONG MEN WITH **CASUAL PARTNERS**



always use condoms



any condomless anal intercourse



STIs

71.6% of sample tested for any STI 21.5% in the last 12 months, of which

reported an STI diagnosis



HIV TESTING 92.7% of sample have ever tested for HIV



STATUS HIV STATUS AMONG PARTICIPANTS TESTED 86.5% 4.9% 8.5%

HIV- HIV+ Don't Know?

HIV NEGATIVE HIV NEGATIVE
PARTICIPANTS IN THE **LAST 12 MONTHS**

Got tested 86.1[%]



Got tested 3 or more times







knowledge of the existing availability of PEP, of which



had taken PEP in last 6 months





of HIV negative men had knowledge of the existing availability of PrEP, of which



had taken PrEP in last 6 months



TEST 43.9% 20.6% 18.4% 13.0% 1.5%

TOP 5 SOURCES FOR HIV & STI INFORMATION



in English

DISCRIMINATION

in the past 12 months

37.2° **25.4**°

Homophobia



For more information about the study and for a copy of the report please click here: https://csrh.arts.unsw.edu.au/research/publications/gcps/







