



**Bachelor of Science (BSc)**  
**BSc Advanced Science**  
**BSc Medicinal Chemistry**  
**Honours in Chemistry 2024**

Never Stand Still

Science

School of Chemistry






# WELCOME

The School of Chemistry at UNSW is one of the leading centres of chemistry research in Australia. Composed of over 30 well-funded **research teams**, we are located in the following buildings on lower campus: Dalton (F12), Chemical Sciences Building (F10NA and F10), Hilmer Building (E10) and the Science and Engineering Building (SEB). The School has state of the art research facilities that enable research spanning the entire breadth of chemistry. The UNSW Mark Wainwright Analytical Centre (MWAC) is co-located adjacent to the School of Chemistry (F10NA) and provides major research facilities that are unsurpassed internationally.

Research in the School of Chemistry can be classified in **four strategic areas**:

- Nanoscience
- Energy & Environment
- Medicinal Chemistry
- Catalysis & Industry



In each area our School has world-renowned scientists that make significant impact on international research, making an impact in areas diverse as medicine, the molecular sciences, chemical industry and materials science.

The School of Chemistry at UNSW has strong links to Australia's professional body for chemists, the Royal Australian Chemical Institute (**RACI**) and the International Union of Pure and Applied Chemistry (**IUPAC**). It also has close ties with the American Chemical Society (**ACS**). Several research team leaders hold senior positions in the RACI, and the NSW state branch is located in the School. Professor Sir Fraser Stoddart (2016 Nobel Laureate) has also commenced research activities within the School.

The School welcomes applicants for Honours from students throughout the world, acknowledging that the Honours year is an outstanding introduction to research. We are confident that the wide range of research undertaken in the School provides applicants with a rewarding Honours year.

Professor Leslie Field (Head of School)

Associate Professor Neeraj Sharma (Chemistry Honours Coordinator)

## OVERVIEW OF HONOURS

This booklet provides details for students interested in undertaking an Honours year with a major in chemistry in either the BSc or BSc Advanced programs or undertaking a BSc Medicinal Chemistry (3992). If you are an external applicant you would have an equivalent BSc.

Is the Honours year worth the extra year it takes? The answer is certainly “Yes!” for many people.

- The response from potential employers in industry and the public sector is that they will employ a good Honours graduate over someone with a pass degree.
- Honours is necessary for anyone contemplating postgraduate study in chemistry.
- Honours gives you “hands-on” experience in tackling projects, and provides a rewarding finishing quality to your education.
- Honours provides you with experience at managing your own project, independence and time management skills.
- Most important of all, the Honours year allows you to work closely with the staff in the School of Chemistry, and it transforms the University for you into a very human organisation that has people who support you.

## HONOURS IN THE SCHOOL OF CHEMISTRY


Every **term** the School of Chemistry offers Honours programs that are suitable for students enrolled in **Science, Advanced Science, Medicinal Chemistry and Environmental Science** or **with a related undergraduate Science degree**. School of Chemistry researchers also supervise or co-supervise the equivalent of a Honours project with colleagues in the Faculty of Science and in other Faculties which are typically made in a case by case basis.

***We welcome all Science graduates from all over the world to apply to our Honours programs. You can start in any term of the year.***

Within the School of Chemistry all programs involve a proportion of coursework. Assessment is based **entirely** on your record during the Honours year, during which students have the opportunity to demonstrate skills in both research and coursework.

In the UNSW3+ model, the streamlined access to Honours is shown in the figure below. Students enrol in 48 credit points by enrolling in CHEM4516 every term for full time enrolment or combinations of CHEM4516 and CHEM4508 for part time enrolment. Marks for all courses, your Honours year mark is provided at the end of

course. Feedback is provided on progress throughout the course by both the primary supervisor and other researchers within the School of Chemistry. Although Honours can be started in any term, we would encourage students to plan for a T1 or T3 start for Honours.



### Steps to apply

1. Choose 3-4 supervisors from this booklet.
2. Arrange online or in-person meetings with each supervisor and request an email stating that they are willing to supervise your honours project.
3. Log on to <https://www.science.unsw.edu.au/honours> to apply for Honours. You will need to attach the supervision email in this application and rank your preferences.
4. For external applicants, please apply for admission either before or with this process.

### Timing

Every term the Faculty of Science opens the application process (<https://www.science.unsw.edu.au/honours>) around week 5 for admission into the following term. This is open typically until week 10 or so. In week 11, this is provided to the School of Chemistry and we send out emails shortly thereafter. Honours commences on Day 1 of Week 1 of every Term.

### Preferences

We will endeavour to provide students with their first preference. In instances where multiple students have selected the same supervisor, the two students with the highest WAM will be allocated to this supervisor and the other students will be assigned to their next preference.

## WHAT DOES AN HONOURS YEAR ENTAIL?

The aim of the Honours year is to continue your development into a well rounded chemist and research scientist by exposing you to independent research, advanced courses in Chemistry and a broad range of fields in chemistry through your attendance at research seminars.

The **original research project** forms the main component of the Honours year. Staff members offer numerous projects in their field of research, some of which are suitable for Chemistry Honours. Students nominate supervisors in research areas that interest them. They then work on a project under the close supervision of an allocated supervisor for the duration of the Honours year. Student-supervisor allocations are made on the basis of student WAM and demand for specific supervisors. Every attempt is made to allocate students their preferred supervisor or supervisors. Supervisor selections - <https://www.science.unsw.edu.au/honours>

**Basic research skills** are developed to prepare you for your research project; areas covered include writing a research proposal, presentation skills, research ethics and modern occupational health and safety requirements.

**Advanced level lecture courses** are designed to extend your knowledge and broaden your understanding of key branches of chemistry. Honours students must undertake 3 short courses offered by the School of Chemistry. These advanced level courses are a combination of either theory, skills or both.

**Research seminars** are conducted throughout the year and are an important means of exposing Honours students to research conducted in the School and at national and international institutions. Attendance is compulsory for the Thursday 12pm seminars and unsatisfactory attendance will result in reduction of your Honours grade.




# ADMISSION

## ELIGIBILITY

For admission to Honours in the School of Chemistry, it is expected that a student will have achieved a credit average of WAM over 65 and have a BSc (or BSc Advanced) with major in chemistry. Students with qualifications in other disciplines may also be eligible for admission.

Students who have completed pass degree requirements at a University other than UNSW, or who have already graduated with a pass degree from UNSW or elsewhere, may be eligible to undertake an Honours year in the School of Chemistry. In such cases, please contact the Honours Coordinator ([neeraj.sharma@unsw.edu.au](mailto:neeraj.sharma@unsw.edu.au)) for clarification.

**In all cases, admission is subject to the formal approval of the Head of School.**



## HOW TO APPLY

- If you are eligible to enrol (see previous page), consult the School of Chemistry Research Booklet (<http://www.chemistry.unsw.edu.au/current-students/undergraduate/honours-research>) to determine which areas of research interest you and discuss these with the relevant academic member of staff. **It is strongly recommended that you talk to all prospective supervisors in your area of interest (i.e., at least 3 potential supervisors).**
- A good working relationship with your supervisor is paramount for the success of your Honours year. The choice of project is also important, and you are advised to obtain as much information as possible before making your decision. Find out what exactly is involved and what would be expected of you if you were to undertake a particular project.
- Honours deadlines for supervisor selection can be found via the on-line portal (<https://www.science.unsw.edu.au/honours-apply>). Before the due date for each term you are required to submit this form with at least **three** supervisor preferences. Rank your order of preference. You must have spoken to and been given permission by at least three supervisors prior to nominating them (**note:** a cross-check will be made with all nominated academics).
- The School will contact you via email to advise the outcome of your application for Honours.

### For further information and assistance, contact:

Honours Coordinator, Associate Professor Neeraj Sharma:

[neeraj.sharma@unsw.edu.au](mailto:neeraj.sharma@unsw.edu.au)





# ASSESSMENT

The Honours degree is graded into Class 1, Class 2 Division 1, Class 2 Division 2, Class 3 or Honours may not be awarded (see below for grade boundaries).

Assessment is on the basis of performance in the various components of the course, including the research project and the formal course work. Note that in order to be awarded Honours, you **must** achieve a satisfactory performance (>50%) in each component (coursework **and** research).

**Course Work (10%):** You are required to take three different short courses during your Honours year. Details are available from the Honours Coordinator (A/Prof Neeraj Sharma).

**Research Component (90%):** The research project is the distinctive feature of the Honours year. It is the major undertaking of the year and is both the most challenging and rewarding aspect of Honours.

Students work on original research projects conceived and overseen by a member of staff. While you will be instructed by your supervisor on the nature of the project and will be given guidance in how to conduct the project, it is expected that you will perform all experimental work independently. You will be expected to prepare and analyse experimental results and work with your supervisor to identify and overcome any problems. At the completion of the year you will present a thesis detailing the background, aims, experimental procedures, results and future directions of your project.

**Research Proposal and Introductory Viva-voce:** To assist in the preparation of your thesis and to allow your writing style to be improved, you will also be required to submit a research report describing your chosen research project and a review of pertinent literature. You will also do a short presentation on your project and will be asked questions by a panel of experts. This will be in the first term of your Honours year and written feedback will be provided for both aspects. This part is worth 10% of your mark.

**Thesis Submission:** You will write your thesis and submit it during your third term in Honours with the dates made available on your first term. Your thesis is a significant portion of your Honours mark and should be read and edited by numerous people including your supervisor.

You are required to submit a final version of your thesis that incorporates the Thesis Committee's comments to your supervisor and the School for completion of Honours. Failure to do so will result in your Honours grade being withheld. The thesis accounts for 50% of your mark.

**Honours Seminar:** You will be expected to present your work at two seminars during your Honours year; one will be a short overview of your project in your first term which

will be graded within your **introductory viva-voce**; and a longer seminar detailing your results to be presented after submitting your thesis; this seminar is typically 15 minutes long plus up to 10 minutes for questions. It comprises 10% of your mark.

**Oral thesis defence (viva-voce):** The thesis defence will be shortly after you have submitted your Honours thesis and presented your final seminar and will be a closed examination of around 20-30 minutes with a panel of experts. It will comprise 20% of your mark.

**Attendance at School Research Seminars:** Attendance at School Research Seminars is compulsory.

### **HONOURS DEGREE GRADE BOUNDARIES (%)**

Class 1	85+
Class 2 Division 1	75 – 84
Class 2 Division 2	65 – 74
Class 3	50 – 64



# **HONOURS SUPERVISORS**



## A/PROF. GRAHAM BALL

Level 2, Dalton Building (F12)  
T: 9385 4720 E: [g.ball@unsw.edu.au](mailto:g.ball@unsw.edu.au)

### NMR SPECTROSCOPY AND COMPUTATIONAL CHEMISTRY: APPLICATIONS TO ORGANOMETALLIC AND BIOLOGICAL CHEMISTRY

Our research focuses on applying NMR spectroscopy to shed light on important chemical problems, often in the areas of organometallic and biological chemistry. NMR spectroscopy is probably the most powerful technique available to the chemist and the Mark Wainwright Analytical Centre is bristling with state-of-the-art instruments eagerly awaiting **YOU** to run experiments that push the boundaries!

Our experimental work is complemented and enriched by using computational techniques. We model small chemical systems with *ab initio* and DFT methods and biomolecular systems with molecular mechanics and QM/MM methods. This is a superb way to get detailed information about your molecules and their reactivity without all the risk assessments!

#### (a) Short-lived metal complexes and reactive intermediates


We use photochemistry in combination with *in situ* NMR at low temperatures to study molecules that have fleeting existence at room temperature. With this strategy, we have observed several types of alkane complex<sup>1,2,3</sup> including the JACS cover opposite<sup>1</sup> and even complexes where xenon acts as a ligand.<sup>4</sup> Alkanes contain no lone pairs for binding to the metal centre. Instead, they bind using the electrons in the C-H sigma bond. This is why they are poor ligands and their complexes are so short-lived (~100 ms maximum lifetime at 25 °C).



#### (i) Alkanes: Binding and Beyond (in collaboration with Prof. Les Field)

Chemists around the globe have been working on ways of converting relatively unreactive alkanes found in petroleum into useful compounds using process known as C-H activation. Alkane complexes are key short-lived intermediates in the activation process.

Current projects are aimed at answering questions such as: Can we make more stable alkane complexes? Can we do chemistry with the alkanes when they are bound? When bound to a cationic metal centre, the alkane should be activated towards conversion into molecules with functional groups, which would be revolutionary new chemistry! We have recently achieved some exciting results making the most stable alkane complexes observed to date using Os compounds and there is much scope to extend this chemistry to similar Fe and Ru complexes and explore the chemistry of the bound alkane.



## (ii) Computational design of new exotic molecules: alkane and noble gas complexes


We employ computational methods (DFT, *ab initio*) to aid the design and understanding of these fascinating compounds. Can we design then observe complexes with ligands that bind even more weakly than alkanes e.g. Xe and Kr even? For example, the recently observed cationic alkane complexes shown above were designed computationally prior to observation.<sup>2</sup>

Projects in these areas can be primarily synthetically based (making new alkane complex precursors), NMR spectroscopy based (observing the new complexes and their reactions) or computationally based (designing new compounds and predicting their reactivity). The 3 components can be blended to suit the interests of students tackling the project.

- 1 Young, R.D.; Lawes, D.J.; Hill, A.F.; Ball, G.E. *J. Am. Chem. Soc.*, **2012**, *134*, 8294.
- 2 Yau, H.M.; McKay, A.I.; Hesse, H.; Xu, R.; He, M.; Holt, C.E.; Ball, G.E. *J. Am. Chem. Soc.* **2016**, *138*, 281.
- 3 Young, R.D.; Hill, A.F.; Hillier, W.; Ball, G.E. *J. Am. Chem. Soc.*, **2011**, *133*, 13806.
- 4 Ball, G.E.; Darwish, T.A.; Geftakis, S.; George, M.W.; Lawes, D.J.; Portius, P.; Rourke, J.P. *Proc. Natl. Acad. Sci. USA.*, **2005**, *102*, 1853.

## (b) Anti-cancer drug-DNA interactions (in collaboration with A/Prof Larry Wakelin, A/Prof Luke Hunter and Dr Don Thomas, NMR Facility)


DNA presents one of the most logical and practical targets for anti-cancer therapeutics. We are investigating the binding of several mono and bis-intercalating molecules that show promise as next generation anti-cancer drugs and also the binding of clinically established drugs such as mitoxantone. The solution structures of the DNA-ligand adducts are obtained via a suite of 2D NMR techniques coupled with NOE-constrained molecular dynamics simulations employing the AMBER forcefield. Our recent results have lead to a re-evaluation of how these bis-intercalators interact with DNA.<sup>5, 6</sup>



The project involves a fusion of NMR spectroscopy and molecular modelling, at the molecular mechanics or QM/MM level. The project can be tailored to focus solely on NMR studies, solely molecular modelling or a balanced amount of both. We have a number of drugs synthesised that are ready for investigation.

5. Serobian, A.; Pracey, C. P.; Thomas, D. S.; Denny, W. A.; Ball, G. E.; Wakelin, L. P. G. *J. Mol. Recognit.* **2020**, *33*, e2843.
6. Rowell, K.N.; Thomas, D.S.; Ball, G.E.; Wakelin, L.P.G. *Biopolymers.* **2021**, *112*, e23409.

## (c) New methods for measuring X-H bond lengths using NMR spectroscopy



Unlike organic chemistry, where bonds involving hydrogen atoms have predictable lengths, inorganic chemistry is awash with compounds where X-H bond lengths vary significantly.

We are using various NMR techniques, including the little-known dynamic frequency shift (left) to use NMR to measure bond lengths in inorganic systems. We have recently published a study involving dihydrogen complexes,<sup>7</sup> which contain stretched H-H bonds and there is scope to extend this methodology to measure other, stretched bond lengths such as C-H, B-H and N-H.

7. Gilbert-Wilson, R.; Das, B.; Mizdrak, D.; Field, L.D.; Ball, G.E. *Inorg. Chem.* **2020**, *59*, 15570.



## A/PROF. JON BEVES

Level 2, Dalton Building (F12)

T: 9385 4673 E: [j.beves@unsw.edu.au](mailto:j.beves@unsw.edu.au) | [bevesgroup.wordpress.com](http://bevesgroup.wordpress.com)

### SUPRAMOLECULAR AND COORDINATION CHEMISTRY

Our research involves using the weak interaction *between* molecules to control their function, with a particular focus on using *visible light* to change the properties of colourful molecules. All projects involve some synthesis, and usually NMR spectroscopy to study structure and properties.



It would be great to work with Honours students on the following projects:

#### (a) Photo-driven molecular machines

(collaborations with Prof. Tim Schmidt, UNSW, Prof. Pall Thordarson, UNSW, Prof. Dean Astumian, University of Maine, Prof. Ayusman Sen, Penn State)

We are designing and synthesizing small molecules capable of performing tasks such as controlled motion or selective binding. A particular goal is to control the diffusion of molecules so we can *direct their movement using light* (e.g. with an LED torch), which would offer the potential for applications ranging from pollution remediation to control over biological function.

**Skills:** organic synthesis, NMR spectroscopy, absorption/emission spectroscopy, kinetics...

**Relevant publications:** *J. Am. Chem. Soc.* **2020**, *142*, 20014–20020; *J. Phys. Chem. Lett.* **2021**, *12*, 1236–1243.

#### (b) Molecular photoswitches


(collaboration with Prof. Joakim Andreasson, Chalmers Institute of Technology)


Some types of organic molecules can be isomerised between two forms using light. These two forms typically have very different properties, such as polarity,  $pK_a$  and reactivity. We are looking to use visible light switchable molecules to control molecular reactions, such as driving pH changes or switching ON/OFF catalytic activities.

**Skills:** organic synthesis, NMR spectroscopy, absorption/emission spectroscopy, kinetics,...

**Relevant publications:**

*J. Am. Chem. Soc.* **2021**, *143*, 20758–20768; *Chem. Commun.* **2022**, *58*, 5610–5613; *ChemPhotoChem*, **2020**, *4*, 407–412; *Chem.-Eur. J.* **2020**, *26*, 1103–1110.





### (c) Photoredox catalysis

(collaboration with A/Prof. Evan Moore, UQ, A/Prof. Alex Bissember, UTas)


The use of visible light to catalyse reactions, generally using transition metal complexes, has developed into a major area of research in the past decade. Despite the practical use of common photoredox catalysts in organic synthesis, their performance and the mechanisms of the reactions they catalyse remain very poorly understood. This project will take a “coordination chemistry” approach to the problem, characterising essential redox and photophysical properties of viable photoredox catalysts and correlating these values with synthetic performance.

**Skills:** organic and inorganic synthesis, NMR (including photo-NMR), cyclic voltammetry, X-ray crystallography, photophysical measurements,...

**Relevant publications:** *Angew. Chem. Int. Ed.*, **2020**, 59, 9522-9526; *Inorg. Chem.* **2020**, 59, 9135-9142 *Inorg. Chem.* **2018**, 57, 8476-8486; *Inorg. Chem.*, **2016**, 55, 12737-12751.


### (d) Self-assembly of functional structures

Using appropriately designed molecular components, large and symmetrical structures can be formed by “self-assembly” upon simple mixing of the different components. The resulting structures can exhibit remarkable properties, especially when constructed using transition metal complexes. This project will build useful redox- and photo-properties into these ordered structures, either as components (as in the two examples shown) or by the encapsulating metal complexes within organic hosts to tune their properties.




**Skills:** organic and inorganic synthesis, NMR, mass spectrometry, cyclic voltammetry, X-ray crystallography, photophysical measurements,...

**Relevant publications:** *Angew. Chem. Int. Ed.*, **2022**, in press; *Chem. Eur. J.* **2022**, 28, e2021044; *Chem.- Eur. J.* **2020**, 26, 1103-1110. *Chem.- Eur. J.* **2019**, 25, 5708-5718; *Org. Lett.*, **2017**, 19, 4034-4037.





### (e) ...other projects tailored to your interests!

Our other interests cover areas ranging from NMR to microfluidics to polymer chemistry!







## DR. JEFFREY BLACK

Office: Level 1, Dalton Building (Rm 133)

Email: [Jeffrey.black@unsw.edu.au](mailto:Jeffrey.black@unsw.edu.au)

### CHEMISTRY EDUCATION AND ToF-SIMS

---

The Chemical Education group at UNSW is focused on improving the learning experience and outcomes of students studying chemistry in a scholarly manner, contributing to the global chemistry and science education communities.

While our primary focus is on tertiary chemistry students, it may be applicable to other subjects, especially other sciences as well as secondary education or education in general. As such we welcome potential students with diverse backgrounds in science and/or education.

My personal interest is in the use of technology to assist with learning and assessment including as preparation for other activities, complementing learning activities, standalone learning activities, and improving assessment and feedback practices with the use of technology. Additionally, I have a passion for lab work and am eager to explore the development of new laboratory experiments to enhance student learning while providing an authentic experience.

Projects may involve intervention design and evaluation; engaging with chemistry students through learning activities such as labs, workshops and tutorials; interviews and questionnaires/surveys of students and staff; artifact/data collection; as well as both qualitative and quantitative analysis of data.

#### **Digital Assessment (in collaboration with Dr Siobhán Wills)**

With technology becoming more widely used in all aspects of life, the use of pen and paper based assessment appears very much out of place. However the shift to digital assessment brings along its own opportunities and challenges. Some important questions are how students perceive digital assessment and if any groups are advantaged or disadvantaged by this shift.

Projects in this area may include new digital assessment techniques and tools; tools to prepare students for digital assessment, or evaluation of student interactions, performance and perceptions of digital assessment.

#### **Technologically assisted assessment (in collaboration with Dr Siobhán Wills)**

Even with assessment remaining as pen and paper, there is opportunity for assessment to be enhanced with technology, such as measures to prevent collusion, increasing efficiency of marking, providing students with better feedback, or the use of technology to assist students in completing the assessment.

Projects in this area may include designing new tools or evaluating existing tools to determine their effectiveness of preventing and detecting collusion, the efficiency of the assessment preparation and marking process, the usefulness of feedback provided to students, and any student or staff perceptions on the process.

## Virtual labs for increased preparation and performance

Coming into a chemistry lab for the first time can be a daunting experience, with lots of equipment and glassware that may not have been seen before, in a new space, with new techniques needing to be performed, along with potentially hazardous chemicals. This places a significant cognitive load on students and may make their performance in the lab suffer. By providing virtual lab experiences students can be safely introduced to the lab environment in advance of their laboratory class, making the space a more familiar environment when it comes time for their lab class. It can also allow them to interact with the lab, equipment and glassware to practice techniques in a safe environment to focus more on the experiment and understanding the chemical principles during their laboratory classes.

Projects in this area may include the development of virtual lab tours or virtual lab experiments; and the evaluation of student interaction with these tools and what impact it has had on the learning experience and performance of the student.


## Other Chemistry Education projects of your choosing.

I am happy to entertain any other ideas you may have for chemical education projects. If you have any ideas or questions, feel free to speak to to see what is possible. Collaborations between academics researching Chemical Education at UNSW (or more broadly) for new projects are also possible.

## ToF-SIMS for analysis of tribological surfaces. (collaborators include Drs Jonathan Palmer, Chris Marjo, A/Prof Jason Harper and Prof. Chris Tierney, UNSW; Prof. Larry Scott, Boston College; Prof.'s Sergei Glavatskih and Mark Rutland, KTH, Stockholm)

My other passion is research in sustainable materials and processes for a sustainable future with a current focus on developing an understanding of how lubricants are transformed during use with the aim of producing better, and longer lasting lubricants.

Time of Flight – Secondary Ion Mass Spectrometry (ToF-SIMS) allows probing the 3D distribution of chemical species in a solid surface. This allows developing an understanding of the breakdown and tribofilm formation pathways of lubricants, including how different species can influence further breakdown and formation; and how this process evolves over time.



ToF-SIMS analysis showing the breakdown products of an orthoborate ionic liquid in a wear scar after a lubrication test.



## A/PROF. ALEX DONALD

Level 6, SEB (E8)

T: 9385 8827 E: [w.donald@unsw.edu.au](mailto:w.donald@unsw.edu.au)

### FUNDAMENTAL & APPLIED MASS SPECTROMETRY


---

Mass spectrometry is a core enabling technology that is used in many emerging and existing scientific fields. Dr. Alex Donald and his team are developing and applying experimental methodologies in mass spectrometry with a focus on problems in chemistry and biochemistry. We are looking for students who are interested in developing a valuable skillset in mass spectrometry and allied topics.

#### (a) Rapid, ultra-sensitive protein structure elucidation by mass spectrometry

Potential drugs, pesticides, and antibiotics often fail because they bind to many proteins, leading to off-target side effects and safety issues. Pesticides and antibiotics can fail because of resistance resulting from changes to binding sites. This project will develop a method for rapidly discovering classes of molecules that bind to unique sites on proteins. This will provide scientists with novel starting points for designing new bioactive molecules aimed at improving effectiveness, safety, and preventing resistance.

The development of new pharmaceuticals is frequently delayed by the time and resources required to identify the sites that new chemical entities bind to protein targets. A recent breakthrough discovery in our laboratory has resulted in the ability to completely characterise large protein sequences directly from single mass spectra. This project aims to leverage this breakthrough by developing a rapid new approach for revealing ligand-protein binding sites using whole-protein mass spectrometry. The success of this project will enable novel sites of interactions between molecules and protein targets to be discovered rapidly and with high sensitivity. This will allow the efficient design of next-generation classes of bioactive molecules.




#### Where does the drug bind?

Methods for rapidly pinpointing where small molecules bind to druggable targets are urgently needed for discovering the next generation classes of bioactive molecules.

#### (b) Single-cell chemical analysis by mass spectrometry

We are interested in answering the fundamental question of what makes a cancer cell a cancer cell? Why are some cells drug-resistant while others are susceptible? Why do some metastasize while others do not? Not every cell was created equal. Individual cells within a population can be as dissimilar as the members of human families. Thus, we need to be able to perform chemical analysis on the contents of single cells, which requires the development of powerful analytical methods that have unprecedented sensitivity and selectivity.

For single-cell chemical analysis, mass spectrometry is one of the most promising analytical techniques because it enables many different types of molecules to be rapidly detected and identified nearly



**Single-cell mass spectrometry: What makes a cancer cell a cancer cell?** Powerful analytical methods must be developed to enable the contents of single cells to be identified and quantified with unprecedented sensitivity and selectivity.

simultaneously from exceedingly small sample volumes. However, matrix ion suppression is a key challenge that hinders the ability of scientists to detect the vast majority of metabolites and biomolecules in human cells. Recently, we have developed a novel, surface-selective ionization approach that enables trace chemicals to be rapidly detected from complex mixtures with minimal ion suppression using mass spectrometry.

In this project, you will take this research to the next level by fabricating novel surface-enhanced microprobes to sample and analyse the contents of single cells by a range of mass spectrometry techniques to target important disease biomarkers. The success of this project will provide a rapid, high-throughput platform to characterise a wide variety of important biomarkers expressed uniquely in each cell, with the goal of understanding how cellular heterogeneity leads to disease states and drug resistance.

### (c) Cancer breathalyser

Imagine a breathalyser test that can sniff out cancer and other diseases. The ultimate goal would be a personalised and highly accurate warning system for diagnosing disease in the earliest possible stages to maximise the possibility of recovery. This will require (i) high sensitivity, (ii) reliable detection, (iii) rapid sampling, and (iv) selective detection of many different types of molecules that are indicative of disease.

We have recently developed a compact ionisation method, called “surface enhanced ionisation,” that can be used to directly ionise analytes from highly complex chemical mixtures without sample preparation for rapid detection by mass spectrometry. This is important because it eliminates chromatographic instrumentation which will significantly improve the performance of portable handheld mass spectrometers by (i) reducing size and power requirements and (ii) increasing sensitivity and tolerance for complex mixtures.

In this project, you will use surface enhanced ionisation mass spectrometry to rapidly detect volatile organic molecules in breath and saliva that are “signatures” for lung and breast cancer with ultrahigh sensitivity. This project is part of a longer-term thrust towards developing a high performance portable, handheld, and personal mass spectrometer for monitoring/detecting disease and detecting harmful substances in your vicinity.



**Portable mass spectrometer for personal chemical analysis:** The ultimate device for preventative medicine? New and improved field deployable ionisation methods are urgently required to enable complex mixtures to be directly analysed (e.g., for early detection of cancer and other diseases). *J. Am. Soc. Mass Spectrom.* **2008**, *19*, 1442-48.



## DR. WESLEY DOSE

Level 2, Dalton (F12)

E: [wesley.dose@unsw.edu.au](mailto:wesley.dose@unsw.edu.au)

### MATERIALS FOR ENERGY STORAGE

We are in the midst of a battery revolution! Battery markets are growing rapidly and are set to expand even more quickly over the coming decade as we transition to cleaner energy sources and pursue net zero emissions. My research group focuses on the most pressing scientific challenges facing current and future rechargeable batteries, encompassing lithium-ion and beyond lithium-ion chemistries.

Why do batteries “die”? How can we increase their storage capacity? Can we charge in under 10 min? How do we make batteries more sustainable? Can batteries be recycled? What will the materials of the future be? Your research will seek to answer one, or more, of these questions.

The research activities in my group span several areas of chemistry, with projects available in inorganic/solid state (electrode materials), organic (liquid electrolyte), and physical (reactions at an electrode interface) chemistry. Students in the group will gain expertise in electrochemistry and a range of characterisation techniques (NMR spectroscopy, mass spectrometry, X-ray/neutron diffraction, X-ray absorption spectroscopy, electron microscopy, etc.). If you would like to learn more, feel free to contact me via email or on Twitter @DoseChemistry.

**It would be great to work with Honours students on the following projects:**

**(a) Advanced materials for lithium-ion batteries**

Lithium (Li)-ion batteries are currently the leading energy storage chemistry for use in electric vehicles (EVs) and stationary storage applications (e.g., residential- and grid-scale storage). Projects in this area will focus on understanding the chemistry of next-generation cathodes and novel electrolytes in full cells.

*Low or no cobalt (Co) positive electrode materials*

Cathode materials with low or no Co are promising next-generation materials to achieve low cost, high energy density, and more sustainable Li-ion batteries. However, these materials suffer from rapid performance fading issues that presently limit their lifetime. This project will investigate the degradation mechanisms for cathode materials from one of the following classes of materials: Ni-rich  $\text{LiNi}_x\text{Mn}_y\text{Co}_z\text{O}_2$  (NMC), Li- and Mn-rich, disordered rock-salt, and disordered spinel.

*New Li-ion battery electrolytes*

Our recent findings show that conventional Li-ion battery electrolytes are incompatible with next-generation high-energy cathodes, in particular at high voltage (<https://doi.org/10.1021/acsenergylett.2c01722>). This project will seek to understand the degradation of conventional electrolytes and develop new electrolytes that are mutually compatible with next-generation cathodes and next-generation anodes, while also having appropriate physical properties (i.e., viscosity, conductivity).




## (b) Beyond lithium-ion batteries

Concerns about the future cost of lithium, due to increasing demand, are driving battery research into new chemistries beyond Li-ion. Sodium (Na)-ion batteries are a front-runner because of the relative abundance of raw materials, their environmental-friendliness, safety, and low price (<https://doi.org/10.1039/C7EE02995K>). These projects will tackle the most pressing issues limiting the storage capacity and long-term performance of Na-ion batteries.

### *Sodium ion inventory*

The low sodium content of best in class Na-ion cathode materials and irreversible sodium trapping reactions at the anode severely limit the energy density of current Na-ion batteries (<https://doi.org/10.1016/j.coelec.2021.100827>). In this project we will discover new sodium source materials that increase the sodium inventory, and hence energy density, of the full cell. Further, we will elucidate the underlying mechanisms leading to device improvement.



### *New Na-ion battery electrolytes*


Innovation in electrolytes for Na-ion batteries is at present quite limited. This project will untangle the reaction mechanisms causing electrolyte degradation and design new electrolyte solutions with a particular emphasis on high-voltage stability.

### *Fast charging*

Fast charge of batteries (e.g., charge in <10 min) is a critical challenge that must be addressed to ensure widespread adoption of batteries into certain markets, such as EVs. Present-day high-energy batteries are unable to achieve this without significant degradation of the battery performance. In this project, we will explore the main limitations for fast charge in the context of Na-ion full cells. We will use an approach combining cell construction, various testing conditions, and extensive materials characterisation.

## (c) Recycling of lithium-ion batteries

The number of large Li-ion battery modules, as in EVs and residential- and grid-scale storage, reaching the end-of-life (EOL) is currently small. But as these markets grow this will soon exceed several thousand per year. This necessitates the development of safe, economic, and environmentally sound processes to reuse or recycle the materials contained within Li-ion batteries, many of which are considered critical and/or high value resources. This project seeks to explore new physical, chemical, and electrochemical processes for reuse, recycle, and up-cycle of materials extracted from EOL Li-ion batteries.





## DR. ALBERT FAHRENBACH

Level 7, Hilmer (E10), 719


E: [a.fahrenbach@unsw.edu.au](mailto:a.fahrenbach@unsw.edu.au); W: [fahrenbachresearch.com](http://fahrenbachresearch.com)

Director of the Australian Centre for Astrobiology

### ORIGINS OF LIFE ORGANIC CHEMISTRY

Ever thought about how life got started? Are you interested in astrobiology? Then consider doing your honours with my group!


**My group seeks to develop experimental and theoretical models for understanding how “life-like” behaviours can emerge from simple chemical systems. We are particularly interested in developing and understanding autocatalytic reaction networks that can be exploited for the construction of a simple protocell “from scratch”.**



It would be great to work with Honours students on the following projects:

#### (a) Optimising Autocatalytic Syntheses for RNA Precursors

(Co-supervisor: Associate Professor Giancarlo Pascali, School of Chemistry)



Autocatalysis is an essential feature of life. The cell utilizes a variety of different autocatalytic reactions to carry out metabolism. Autocatalysis affords not only dynamic “life-like” behaviours, but also provides a mechanism for reproduction at the molecular level. In fact, you might consider the entire process of cellular reproduction a

complex suite of coupled autocatalytic reactions.


While the cell is a master of autocatalysis, we still don't understand very well how to utilize autocatalysis in synthetic contexts. The aim of this project is to optimise an autocatalytic reaction network, known as the formose reaction, for production of RNA precursors using a continuous stirred tank reactor (CSTR).

Unlike your typical organic reaction, which takes place inside of a round-bottom flask and so must eventually come to equilibrium, a CSTR is an open system, continually supplied with fresh reagents pumped into the reactor, while products are likewise continually removed at the same rate. To a first approximation, a CSTR allows one to carry out synthesis more like the cell does, which constantly takes in nutrients from the environment while expelling waste. More than just fundamental science, we expect this type of autocatalysis carried out within a CSTR to lead to novel methodologies for production of commercially important synthetic compounds.

## (b) Autocatalytic Reaction Networks in Vesicles for Emergent “Life-Like” Behaviours

(Co-supervisor: Dr Anna Wang, School of Chemistry)

The cell is a master of autocatalysis, but we still don't understand the fundamentals of autocatalytic reactions when carried out in cell-sized compartments. This project aims to demonstrate “life-like” behaviours made possible when autocatalytic reactions are carried out within the confines of a protocell – a model for a simple cell based on lipid vesicles.




One of the cell's most fundamental life-like features is bistability. Bistability refers to the fact that many metabolic processes can be switched “on” or “off” (hence are bistable) depending on a variety of factors such as nutrient supplies in the environment. Under the right circumstances, autocatalytic reactions afford mechanisms for achieving bistability, being able to turn themselves on and off in response to exposure to appropriate chemical stimuli. In this project, we aim to show that the autocatalytic formose reaction when carried out inside of a lipid vesicle can be turned on and off by different chemical stimuli, molecules which can be considered as environmental “nutrients”. Showing this sort of bistability in a protocell vesicle will reveal how even relatively simple molecular systems can lead to life-like emergent behaviours that allow a protocell to respond to its environment “intelligently”.

## (c) Autocatalysis Starting from Atmospheric Chemistry

(Co-supervisor: Dr Chris Hansen, School of Chemistry)

The autocatalytic formose reaction depends on the constant supply of formaldehyde (CH<sub>2</sub>O), but how could formaldehyde have been made available on early Earth during the age of prebiotic chemistry around 4 billion years ago? This project aims to demonstrate that formaldehyde could have been produced atmospherically by the photochemical reaction of CO<sub>2</sub> and H<sub>2</sub>O, which then “rained out” onto the early Earth's surface.



Given the lack of an ozone layer, the flux of UV-light penetrating the early troposphere especially in the 200–350 nm region is suspected to have been much greater in comparison to today. Moreover, the partial pressure of CO<sub>2</sub> is thought to have been as high as ~100 bar. In this project, we aim to model experimentally the atmospheric synthesis of formaldehyde using a custom-built UV reactor that takes into account these early Earth atmospheric conditions. We expect to see the accumulation of formaldehyde by passing the gaseous reaction mixture through an aqueous solution, mimicking the early Earth's surface. Demonstrating accumulation of formaldehyde will support the idea the autocatalytic formose reaction could have occurred on early Earth some 4 billion years ago.





## PROF. LES FIELD

Level 10, Chemical Sciences Building (F10)

T: 9385 2700 E: [l.field@unsw.edu.au](mailto:l.field@unsw.edu.au)



### SYNTHETIC ORGANOMETALLIC CHEMISTRY

- Research in the Field group is centred around synthetic organometallic chemistry:
  - Development of organometallic catalysts that are able to activate small molecules (such as  $N_2$ ,  $CO_2$ ,  $CH_4$  etc), and to functionalise organic hydrocarbons ( $CH_4$ , ethylene, acetylene etc) to make value-added products and perform specific organic transformations.
  - Development of organometallic polymers for application in areas such as molecular conductors, molecular semiconductors and molecular electronics.
- Skills you will learn in the Field group:
  - Organic & organometallic synthesis; manipulation of air and moisture sensitive compounds.
  - Structure elucidation and determination of reaction mechanisms.
  - Heteronuclear NMR spectroscopy ( $^{31}P$ ,  $^{15}N$ ,  $^{29}Si$ ,  $^{19}F$ ), 2D NMR spectroscopy, IR spectroscopy, electrochemistry and X-Ray diffraction.

It would be great to work with Honours students on the following projects:

#### (a) Organometallic Polymers

Organometallic compounds containing complexed metals linked by bridging groups have many potential applications in materials science. We are particularly interested in the use of unsaturated organic groups (e.g. alkynes and arenes) as the bridging units and we are developing new methods for forming metal complexes where the metal centres are bridged by organic acetylides. Acetylide-bridged organometallic complexes show interesting electrochemical behaviour, and electronic communication between the two metal centres is often observed.




The majority of alkyne-bridged organometallic polymers use linear aromatic spacer units as the bridge between metal centres. We are interested in introducing bridges based on naphthalenes and other aromatic systems as well extending the oligomers to 2- and 3-dimensional networks.

#### (b) The Organometallic Chemistry of Carbon Dioxide



Carbon dioxide reacts with many organometallic compounds to give products in which the  $CO_2$  is incorporated into the metal complex. We are exploring new ways to trap and capture  $CO_2$  and new alternate uses for this wasted and environmentally dangerous compound.

There have been reports in the literature of *catalytic* activation of CO<sub>2</sub> to yield formic acid H-COOH (by hydrogenation), acrylic acids RCH=CH-COOH (by reaction of CO<sub>2</sub> with ethylene and terminal alkenes), propiolic acids RC≡CCOOH (by reaction of CO<sub>2</sub> with acetylene and terminal alkynes), and carbonates (by reaction of CO<sub>2</sub> with epoxides). We are exploring the ability of new iron(II) and ruthenium(II) phosphine complexes that we can prepare in the lab to catalytically activate CO<sub>2</sub> and to produce “value-added” compounds using CO<sub>2</sub> as a 1-carbon starting material.



**(c) Metal-to-metal communication through cross-conjugated frameworks (with Dr Martin Peeks)**

Quinones are a class of organic compounds which have a rich redox-chemistry, and which are heavily used as oxidizing agents both by chemists and in biology. Metallaquinones are analogues of quinones where one or both of the oxygen atoms are replaced by metals. This project involves synthesising new bi-metallic or polymetallic quinonoid compounds and examining the redox chemistry and metal-to-metal electronic communication in this unusual class of molecules. The results provide fundamental insight into the nature of electronic communication and could underpin the design of the next generation of advanced materials.



**(d) Alkane binding to metals (with Associate Professor Graham Ball)**

Alkanes are amongst the most stable and unreactive classes of organic compounds, and we have been studying the binding of alkanes to metals. Binding an alkane is the first step to activation of a C-H bond so that it can react with other reagents to make a substituted alkane. We are studying the key short-lived intermediates where one of the C-H bonds in an intact alkane molecule is bound to a metal.

Current projects are aimed at trying to make more stable alkane complexes and then doing chemistry on the alkanes when they are bound. We have been working with cationic metal complexes of Re and Os and we need to extend the study to complexes of Ru and Fe. The approach so far has been to generate a reactive metal species at low temperature by splitting off carbon monoxide from a metal carbonyl compound with UV light, then letting the metal react with an alkane. We characterise the compounds using NMR spectroscopy.

**ToF-SIMS analysis showing the breakdown products of an orthoborate ionic liquid in a wear scar after a lubrication test.**

**Selected publications from the group:**

1. Dinuclear Acetylide-bridged Ruthenium(II) Complexes with Non-aromatic Spacers. Surabhi Naik, Synøve Ø. Scootwell, Hsiu L. Li, Chanel F. Leong, Deanna M. D'Alessandro, Leslie D. Field, *Dalton Transactions*, 2020, 49, 2687–2695. DOI: 10.1039/C9DT04856A.
2. Fe(0)-Mediated Reductive Disproportionation of CO<sub>2</sub>. Peter M. Jurd, Hsiu L. Li, Mohan Bhadbhade, Leslie D. Field, *Organometallics*, 2020, 39, 2011–2018. Published online at dx.doi.org/10.1021/acs.organomet.0c00175.



## SCIENTIA PROF. J. JUSTIN GOODING

Level 7, Hilmer Building (E10)

T: 9385 5384 E: [justin.gooding@unsw.edu.au](mailto:justin.gooding@unsw.edu.au)


Australian Centre for NanoMedicine

### SMART MATERIALS AND SURFACES

Our research group specializes in using self assembled monolayer or other surface modification technique to provide surfaces with unique functionality. The surfaces are the base upon which we build functional devices from nanscale component including polymer, protein, nanoparticles, and porous material. The three major programs in which these surfaces are applied are, biomaterials, biosensor, and drug delivery. The multidisciplinary nature of our research means we need people with interest in medicinal chemistry, surface chemistry, polymer chemistry, nanotechnology or analytical chemistry. All new members of the group will be looked after by a post-doctoral fellows as well as Prof. Gooding. Specific projects are:


#### Digital assays - Sensitive Biosensors for the Digital Age (in collaboration with Professor Richard Tilley)

The detection of disease biomarkers (such as proteins, DNA fragments and RNAs) in biological fluid is essential for the early detection of diseases. One of the primary challenges is the low concentration (typically in the femtomolar range) of the biomarkers. We are looking into new approaches to construct digital biosensors based on plasmonic nanoparticles. With the help



of a dark-field optical microscope, we can look at the scattering arising from individual nanoparticles. The wide field nature of this measurement allows for the simultaneous characterization of thousand nanoparticles. When a biochemical sensing reaction is performed, the optical signature of the nanoparticle is altered thereby leading to change in the colour of the nanoparticle. By setting a threshold, we digitalize the data to 0 (unreacted) and 1 (reacted) nanoparticles. Our aim is to push this approach for the detection of individual biomarkers on individual nanoparticles.


#### Detection of Single Biomolecules using Magnetic Nanoparticles and Nanopore Sensors



A typical biosensors detects many molecules to give the concentration of species. Nanopores, which are commonly proposed for DNA sequencing, can detect single molecules and give concentration of species by counting many single molecules. This avoids the need for calibration however, detection limits are not as low as one expects because of the time taken for the molecules to find the nanopores. We have solved this problem by developing a new type of nanopore, referred to as a nanopore blockade sensor. In this system, antibody magnetic nanoparticles capture the analyte of interest and bring it to the


nanopore. The nanopore modified nanoporesarticle then blocks the nanopore to give a single molecule measurement. An additional benefit is the nanopore blockade sensors can operate in complex biological fluids. This project will involve developing the next generation of this exciting single molecule sensor.

### 3D printing of cells for improved tumour models and drug assays (in collaboration with Australian Centre for NanoMedicine)



Our current understanding of cancerous tumours is heavily based on in vivo experiments in animals or in vitro experiments on tissue culture plates. To date, few techniques exist that can satisfactorily recreate the tumour environment in vitro in 3-Dimensions. Such models would allow biologists to better understand the effect of spatial organisation of biomolecules on cell behaviour. Of particular interest are molecules that trigger cancer cell metastasis, or invasion, to other parts of the body. In our lab we are developing materials that can recreate the 3D tumour environment, made from polymers that provide a matrix for cells to attach to (see figure). In the proposed project, the polymers will be modified to include a peptide (protein-based) crosslink that stabilises the structure. Such protein-based regions are susceptible to degradation by specific types of enzymes (proteases) released by cancer cells when they invade surrounding tissue. The new materials developed in this project will be used as an extracellular matrix for the 3D printing of cells in collaboration with a 3D printing start-up company.

### The synthesis of electrocatalysts for fuels cells that mimic enzyme structure (in collaboration with Professor Richard Tilley)



Electrocatalysts are important in applications as broad as fuel cells to sensors to production of fine chemicals. There are however a clear differences between a man made metallic electrocatalyst and a biological catalyst (an enzyme). In man made catalyst the catalytic sites are on the surface of the particle and the entire particle is conducting. However recent work in *Science* suggests catalytic sites in depressions may in fact be more active. In depressions or clefts are where most catalytic sites are located in enzymes. In this way the catalytic site is separated from the reactant solution which allows the chemical environment to be different from the bulk solution and the site to be protected from other species in solution. In this project we will synthesize catalytic nanoparticles for the oxygen reduction reaction that mimic enzyme structure by having the catalytic sites buried inside the particle but accessible via a small channel. Hence this work will focus on making core-shell nanoparticles, electron microscopy characterisation and performing electrocatalytic experiments with them.



## DR CHRISTOPHER HANSEN

Level 6, Science and Engineering Building (E8)

T: 9065 3085 E: [christopher.hansen@unsw.edu.au](mailto:christopher.hansen@unsw.edu.au)

### The true impact of fluorinated compounds in the atmosphere

Use lasers to learn about the chemical reactions that occur after gas-phase compounds absorb light. I am concerned about the true environmental fate of anthropogenic fluorinated compounds and interested in the molecular environment of interstellar space. Use fundamental physical chemistry/chemical physics to address problems in atmospheric science and astrochemistry!

#### It would be great to work with Honours students on the following projects:


Hydrofluorocarbons (HFCs) are the replacements to the chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs). They have no ozone depletion potential yet still present an enormous risk to the environment as powerful global warming agents. These HFCs have a high infrared activity and long atmospheric lifetimes (decades to centuries) leading to global warming potentials (GWPs) up to 10s of 1000s of times worse than CO<sub>2</sub>. An important aim of my research is to improve the underpinning science that is incorporated into atmospheric chemistry models so that humankind can understand the environmental risk of new compounds before they are emitted in large quantities.

#### (a) UV Photochemistry of Fluorocarbonyls (with Prof. Scott Kable)

Hypothesis: The GWP of a molecule's decomposition products needs to be considered when evaluating its GWP. Particularly for short-lived compounds celebrated as low GWP replacements for hydrofluorocarbons.


Current HFC replacements incorporate reactive chemical subunits (*e.g.* double bonds) that reduce their atmospheric lifetime to weeks. However, the most likely fluorine-containing end-products have a higher risk to the atmosphere than the compounds being replaced. This project aims to identify these products to assess the true atmospheric risk for emission of new fluorine-containing compounds.

Recent results from my group (in collaboration with Prof. Scott Kable's group) have revealed that the decomposition product of an important next generation refrigerant (HFO-1234ze or 1,3,3,3-tetrafluoropropene), with a GWP of zero, is removed from the atmosphere via photolysis to yield a significant quantity of the worst of the HFCs *i.e.* fluoroform (CHF<sub>3</sub>) with a global warming potential ~12 000 times worse than CO<sub>2</sub>.




**What you will do:** Velocity-mapped ion imaging, Fourier-transform infrared (FT-IR) spectroscopy, chirped pulse Fourier-transform microwave spectroscopy, computational chemistry.

**(b) Interstellar Molecules (with Dr Chris Hansen)**



The *Molecular Photonics Laboratories* house sophisticated lasers and equipment with which we can discover new transient chemical species of importance in the gas phase chemistries of our atmosphere and the interstellar medium.



As stars die, they eject complex organic molecules into the interstellar medium, where they live out millennia before being incorporated into new stars and planetary systems. These organic molecules are the seeds of life, but, as yet, we do not know the chemical make-up of the interstellar medium from which planetary systems are formed. Using a star as a lamp, we can peer into this medium using telescopes by observing molecular absorption spectra. However, despite there being hundreds of nibbles taken out of the visible stellar spectra of stars occluded by diffuse clouds, only a few molecules have been unambiguously detected by their visible spectra. The unidentified features are known as the *diffuse interstellar bands* (DIBs), and are the longest standing mystery in astrophysical spectroscopy. In this project, we will develop techniques to capture the spectra of isolated, never-seen-before aromatic cations and radicals (which are the leading candidates for carrying the DIBs), and (hopefully) solve this long-standing problem.

**What you will do:** Laser spectroscopy, electric discharges, vacuum techniques, quantum chemistry



## A/PROF. JASON HARPER

Level 2, Dalton Building (F12)

T: 9385 4692 E: [j.harper@unsw.edu.au](mailto:j.harper@unsw.edu.au)


[www.jasonbharper.com](http://www.jasonbharper.com)

### MECHANISTIC AND PHYSICAL ORGANIC CHEMISTRY

Our research is focussed on understanding how organic processes happen and what affects reaction outcomes. Particularly, this work encompasses examining how structural features in both the reagents themselves and the solvent used can change how a reaction proceeds. This knowledge can then be applied to a range of fields, including bioorganic, synthetic, analytical and environmental chemistry. Being particularly interdisciplinary, there is extensive opportunity for collaboration and projects are currently underway in catalysis, reaction kinetics, synthesis and molecular dynamics simulations.


- a) **Ionic liquid effects on organic reactions: understanding solvation, designing better solvents and getting the reaction outcomes you want.<sup>1</sup>** (collaborators include Dr Ron Haines & Prof. Stuart Prescott, UNSW; Prof.'s Anna Croft, University of Loughborough; Christof Jäger, AstraZeneca, Sweden; Prof. Bill Price, WSU; Prof. Tam Greaves, RMIT)

Ionic liquids are salts that melt below 100°C. They have the potential to replace volatile organic solvents but outcomes of reactions in ionic liquids are often different to those in traditional molecular solvents. The aim of this project is to understand the nature of solvation in these systems – the interactions between a solute and the ions of the ionic liquid – through analysis of reaction outcomes, measurements of solution properties (such as diffusion) and molecular dynamics simulations. The result would be to extend the understanding of these solvent effects we have developed and to use this knowledge to control reaction outcome.



A molecular dynamics simulation showing the organisation of the **cation** and **anion** of an ionic liquid around **pyridine**.


The project would involve kinetic analyses using NMR spectroscopy to monitor the progress of reactions, along with synthetic organic and analytical chemistry. Importantly, it can be readily tailored to either the physical and analytical aspects, with the opportunity to focus on methods to measure interactions and molecular dynamics simulations, or the more synthetic aspects, by focussing on designing new ionic liquids, increasing reaction yield and optimising isolation. Either way, you will be designing solvents to get the reaction outcome you want!



A series of <sup>1</sup>H NMR spectra showing the progress of a reaction, particularly the consumption of the starting material (◆) and formation of the product (◆).

## b) Catalysis using *N*-heterocyclic carbenes: understanding structure/activity relationships<sup>2</sup>


*N*-Heterocyclic carbenes, have significant roles in both organo- and organometallic catalysis, however some carbenes are effective for some processes but not for others; the origin of this is not well understood. This project aims to relate the structure



and chemical properties of carbenes to catalytic efficacy; particularly the effects of changing steric and electronic properties will be assessed. Along with making the precursors to the carbenes, this project involves the opportunity to utilise various characterisation techniques (such as measuring acidity of parent cations to generating electronic probes based on Pd and Se) along with evaluation of catalytic systems; the latter can vary from screening of catalysts to detailed kinetic analyses. The ultimate goal is to be able to rationally choose an NHC catalyst for a given process.

## c) Broader applications of physical organic chemistry.<sup>3</sup> (collaborators include Drs Jeffrey Black, Jonathan Palmer, Chris Marjo and Prof. Chris Tierney, UNSW; Prof. Larry Scott, Boston College; Prof.'s Sergei Glavatskih and Mark Rutland, KTH, Stockholm)

The understanding developed above can be applied broadly – from understanding lubrication mechanisms to develop new compounds for mechanical engineering, through the synthesis of carbon nanostructures, to the preparation of samples to evaluate ancient climates. These projects focus on the ability to transfer understanding from one context to another and the skill sets required vary dramatically between projects. However, they all would suit someone with an interest in combining chemistry with an outside discipline as there will be opportunities to work closely with collaborators in different fields. Ultimately, these projects seek to expand the impact of the knowledge gained through our fundamental research.



ToF-SIMS analysis showing the breakdown products of an orthoborate ionic liquid in a wear scar after a lubrication test.

For more information, visit the group website at [www.jasonbharper.com](http://www.jasonbharper.com)

For recent examples of our work in the above areas see:

1. A. Y. Hsieh *et al.*, *RSC Adv.* **2023**, *13*, 21036; D. C. Morris *et al.* *ChemPlusChem* **2023**, e202300015; *Phys. Chem. Chem. Phys.* **2021**, *23*, 9878; M. D. Coney *et al.*, *J. Org. Chem.* **2022**, *3*, 1767; A. Gilbert *et al.*, *J. Phys. Org. Chem.* **2021**, *34*, e3217; *Org. Biomol. Chem.* **2020**, *1*, 5442; **2019**, *17*, 675 & 9336; J. B. Harper *et al.*, *Phys. Chem. Chem. Phys.* **2021**, *23*, 2742 & **2020**, *22*, 23009; K. T.-C. Liu *et al.*, *Org. Biomol. Chem.* **2020**, *18*, 7388; K. S. Schaffarczyk McHale *et al.*, *ChemPlusChem*, **2019**, *84*, 465, 534 & 9243, **2018**, *83*, 1162. For a review see *Adv. Phys. Org. Chem.* **2018**, *52*, 49.
2. C. Barnett *et al.*, *ChemistrySelect*, **2022**, *7*, e202104348; *Eur. J. Inorg. Chem.* **2021**, *47*, 4954; *Chem. Method.* **2021**, *1*, 374; N. Konstandaras *et al.*, *Org. Biomol. Chem.* **2020**, *18*, 66 & 1910; M. H. Dunn *et al.*, *J. Org. Chem.* **2017**, *82*, 7324.
3. J. J. Black *et al.*, *Sci. Rep.* **2022**, *11*, 24021; P. Rohmann *et al.*, *Tribol. Int.* **2023**, *181*, 108263; **2021**, *161*, 107075; S. A. P. Blake *et al.*, *Dendrochronologia* **2020**, *60*, 125644; X. Zheng *et al.*, *Mires and Peat*, **2019**, *24*, 30; S. R. D. George *et al.*, *Polycycl. Arom. Compd.* **2016**, *36*, 897; *Org. Biomol. Chem.* **2015**, *13*, 9035 & 10745.





## DR. JUNMING HO

Level 2, Dalton Building (F12), Room 220


E: [junming.ho@unsw.edu.au](mailto:junming.ho@unsw.edu.au)

### COMPUTATIONAL CHEMISTRY AND BIOMOLECULAR SIMULATIONS

We develop and apply computational chemistry methods to elucidate the mechanisms underlying many processes in synthesis and in biochemical systems (<https://sites.google.com/view/mmg-unsw/home>). This enables us to design improved chemical reagents, drug molecules or enzymes that our experimental colleagues can test or implement in practical applications. Topics of particular interest include, but are not limited to drug design, solvent effects and supramolecular chemistry. We work closely with experimental groups (here at UNSW and from overseas) so projects can be tailored to include an experimental component if desired. The following outlines several representative projects but feel free to get in touch to discuss your interests. No background beyond 2<sup>nd</sup> year physical chemistry is assumed.

#### (a) Anionophores as novel anti-cancer agents

Anionophores are molecules that bind anions, most commonly through hydrogen bonding. Recent studies have revealed that these molecules can also perturb the ionic gradient in cells by transporting anions across cell membranes thereby leading to cell death (see for example, *Nature Chemistry* **2017**, 9, 667). To further develop their potential as anti-cancer agents, we would like to simulate the binding and transport process for several families of anionophores. In this project, you will learn how to carry out




electronic structure calculations and classical molecular dynamics simulations to determine the free energy barriers for the transport of a ligand across the cell membrane. This project will help establish the molecular pre-requisites for anion transport that will facilitate the design of more effective anion transporters.

#### (b) Understanding isoform selectivity of carbonic anhydrase inhibitors [In collaboration with A/Prof Alex Donald (UNSW) and Prof Claudiu Supuran (Florence, Italy)]

Many enzymes exist in different forms that perform different biological functions. Selective targeting of specific isoforms is crucial for drug potency and minimising side-effects. The carbonic anhydrase enzyme that perform important regulatory functions and has 12 known isoforms in human; two of which are associated with tumour progression (CA XI and CA XII).

In this project, we will develop highly accurate multi-scale computational models to elucidate the molecular interactions that give rise to selectivity towards different isoforms of carbonic anhydrase. This insight will be crucial for the development of next generation CA inhibitors with greater selectivity. (above: crystal structure of a CA inhibitor in the enzyme active site).



The student will learn advanced techniques such as quantum mechanics/molecular mechanics (QM/MM) methods developed in the group to study enzymes and large macromolecules.<sup>1,2</sup>

### (c) Accelerating quantum chemistry calculations




One of the key achievements in modern quantum chemistry is the development of highly accurate methods that can rival experiments. However, the computational cost (CPU hours, memory requirements) is so high that many of these methods are limited to model systems composed of several tens of atoms. This project aims to develop an approximation based on the many-body expansion method (see equation below) to accelerate the calculations of interaction energies between molecules with their environments such as solvents, enzyme active site and surfaces.

Preliminary work in the group has indicated that this approach can lead to a **20-fold reduction** in computing time (e.g. months to days).<sup>3</sup> This project is ideal for a student interested in quantitative and data analysis and willing to pick up some computer programming.

$$E = \sum_{i=1}^N E_i + \sum_{ij} \Delta E_{ij} + \sum_{ijk} \Delta E_{ijk} + \dots$$

### (d) Computer-aided design of fluorinated bioactive molecules [with A/Prof Luke Hunter]

Fluorination is often used by medicinal chemists to impart structural rigidity and/or tune the lipophilicity of drug molecules. This project will use computational techniques (e.g. docking, quantum chemical and molecular dynamics simulations) to determine the 3D shapes, logP values and protein-binding ability of



a variety of fluorinated bioactive molecules.<sup>4</sup> Examples of medically-relevant targets that are currently of interest within the Hunter group are shown on the left. There will also be an opportunity in this project to validate some of the computational predictions through synthesis.

### Selected references

1. Chen, J.; Harper, J.; **Ho, J.** [Improving the accuracy of QM/MM models with polarised fragment charges. \*J. Chem. Theory and Comput.\* \*\*2022\*\*, in press.](#)
2. Chen, J.; Kato, J.<sup>#</sup>; Harper, J.; Shao, Y.; **Ho, J.** [On the accuracy of QM/MM models. A systematic study of intramolecular proton transfer reactions of amino acids in water. \*J. Phys. Chem. B\* \*\*2021\*\*, \*125\*, 9304.](#)
3. Chen, J.; Chan, B.; Shao, Y.; **Ho, J.** [How accurate of approximate methods at modelling solute-solvent interactions in solvated clusters? \*Phys. Chem. Chem. Phys.\* \*\*2020\*\*, \*22\*, 3855.](#)
4. Wu, Y. M.<sup>#</sup>; Salas, Y.; Leung, Y. C.; Hunter, L.; **Ho, J.** [Predicting octanol-water partition coefficients of fluorinated drug-like molecules; a combined experimental and theoretical study. \*Aust. J. Chem.\* \*\*2020\*\*, \*73\*, 677.](#)

**# = former honours student in the group**



## A/PROF. LUKE HUNTER

Level 2, Dalton Building (F12)

E: [l.hunter@unsw.edu.au](mailto:l.hunter@unsw.edu.au)

### MEDICINAL ORGANIC CHEMISTRY


---

In my lab, we seek to make molecules that can treat disease. Our work relies on synthetic organic chemistry as the foundational activity, but we also employ a variety of other techniques such as molecular modelling, docking, NMR-based conformational analysis, solid-phase peptide synthesis, and many types of bioassays. Much of our work is highly collaborative in nature, and my students frequently spend time in other labs across UNSW as part of their studies. The broad project areas described below are constantly evolving, and I hope that the descriptions will serve as the *starting point* for a conversation with you about an ideal project that best suits your interests.

**(a) “Molecular origami”: using fluorine to control the shapes of bioactive molecules**

(in collaboration with Dr. Junming Ho; Dr. Angela Finch [SOMS]; Dr. Nicola Smith [SOMS])


Fluorine is a small atom that packs a big punch. When incorporated into an organic molecule, fluorine can have a dramatic impact on molecular properties such as  $pK_a$ , metabolic stability, 3D conformation, and binding affinity for protein targets. We like to take conformationally flexible lead compounds, and decorate them with carefully designed patterns of fluorine atoms.<sup>[1–6]</sup> This can pre-organise the molecule into the target-binding conformation, thereby enhancing the biological potency and selectivity. In this project, we will apply this concept to the antidepressant drug, imipramine.



**(b) “Molecular Velcro”: targeting DNA to treat brain cancer**

(in collaboration with A/Prof. Larry Wakelin; A/Prof. Graham Ball; Prof. Martina Stenzel; Prof. Bill Denny [Auckland]; Dr. Euphemia Leung [Auckland])


Cancer is a common disease that kills 1 in 3 of us in the Western world. Chemotherapy is the principal treatment for metastatic cancer, but its effectiveness is limited by the resistance that tumour cells can develop to many conventional drugs. We are developing new drugs that will bind to DNA and weld the two strands together in a way that is difficult for tumour cells to repair. This will give potent anticancer activity, with a slower development of drug resistance.



**(c) A “molecular high-altitude chamber”:** activating the hypoxia response to treat stroke

(in collaboration with Dr. Nicole Jones [SOMS]; Prof. Christopher Schofield [Oxford])

Stroke is a leading cause of death and disability in Australia, and the treatment options are extremely limited. We are pursuing a new approach. We're developing drugs that activate nerve cells' natural hypoxia protective mechanisms, which will put nerve cells into damage-control mode after a stroke.<sup>[7]</sup> The key is a molecular-level understanding of the proteins that naturally activate this hypoxia response.



**(d) A “molecular production line”:** new ways to synthesise <sup>18</sup>F-labelled compounds

(in collaboration with A/Prof. Giancarlo Pascali; Dr. Ben Fraser [ANSTO])

<sup>18</sup>F-Labelled compounds are useful tools for PET imaging. We're pursuing efficient new methods for synthesising such compounds, including the use of flow chemistry, electrochemistry and photochemistry. We're also seeking to broaden the variety of <sup>18</sup>F-labelled compounds that are available in the clinic. For example, the pentafluorosulfanyl (SF<sub>5</sub>) group can be considered as a “super CF<sub>3</sub> group,” and it promises to deliver valuable future opportunities in medicinal chemistry and imaging applications.<sup>[8]</sup>



- 
- [1] Y. Lizarme-Salas, A. D. Ariawan, R. Ratnayake, H. Luesch, A. Finch, L. Hunter, “Vicinal difluorination as a C=C surrogate: an analog of piperine with enhanced solubility, photostability, and acetylcholinesterase inhibitory activity,” *Beilstein J. Chem.* **2020**, *16*, 2663.
- [2] A. Lawer, L. Hunter, “Controlling  $\gamma$ -peptide helicity with stereoselective fluorination,” *Eur. J. Org. Chem.* **2021**, 1184.
- [3] A. D. Ariawan, F. Mansour, N. Richardson, M. Bhadbhade, J. Ho, L. Hunter, “The effect of vicinal difluorination on the conformation and potency of histone deacetylase inhibitors,” *Molecules* **2021**, *26*, 3974.
- [4] S. Chen, Y. Ruan, J.L. Lu, L. Hunter, X. G. Hu, “Diastereoselective synthesis and conformational analysis of 4, 5-difluoropipercolic acids,” *Org. Biomol. Chem.* **2020**, *18*, 8192.
- [5] C. Au, C. Gonzalez, Y. C. Leung, F. Mansour, J. Trinh, Z. Wang, X. G. Hu, R. Griffith, E. Pasquier, L. Hunter, “Tuning the properties of a cyclic RGD-containing tetrapeptide through backbone fluorination,” *Org. Biomol. Chem.* **2019**, *17*, 664.
- [6] A. Lawer, J. Nesvaderani, G. M. Marcolin, L. Hunter, “Synthesis and biochemical characterisation of fluorinated analogues of pepstatin A and grassystatin A,” *Tetrahedron* **2018**, *74*, 1278.
- [7] N. L. Richardson, L. J. O'Malley, D. Weissberger, A. Tumber, C. J. Schofield, R. Griffith, N. M. Jones, L. Hunter, “Discovery of neuroprotective agents that inhibit human prolyl hydroxylase PHD2,” *Bioorg. Med. Chem.* **2021**, *38*, 116115.
- [8] G. Surjadinata, L. Hunter, L. Matesic, G. Pascali, “Analytical-scale synthesis of aryl-SF<sub>4</sub>Cl via flow microfluidic technology,” *J. Flow Chem.* **2021**, *11*, 107.



## DR SINA JAMALI

Level 7, Hilmer Building (E 10), 746


E: [s.jamali@unsw.edu.au](mailto:s.jamali@unsw.edu.au)


### Electrochemistry of living cells and materials

My research is centred around advancing electrochemical techniques toward enabling smarter design of materials as well as finding new ways to communicate with cells. Cells communicate via electrochemical mechanisms. Our ability to use technology to talk to cells in their native “electrochemical” language offers potential for new understanding and treatment of disease. Our ability to record and analyse very small currents and voltages that are linked to chemical reactions allow us to get ever closer to the fundamental functions of materials and cells. We exploit electrochemistry to look into the interaction between materials, physiological environment and cells to help with more advanced design of biosensors and biomaterials. New members will work within the Smart Materials and Surfaces research group led by Scientia Prof. Justin Gooding. Following projects are the specific areas of interest while other projects of mutual interest will also be considered. It would be great to work with Honours students on the following projects:

#### (a) Light activated electrochemistry (in collaboration with Professor Justin Gooding)

Localising electrochemistry to small regions on a large surface is of particular interest among recent advances in electrochemistry since it enables screening of single cells in a non-invasive manner. This allows cells response to drugs to be studied individually and also opens a new window to advancement of bioelectronics. Among the current approaches to localised electrochemistry, the Light Activated Electrochemistry (LAE) technique, that has been developed to great depth by the Smart Materials and Surfaces group, offers unique advantages. The activation of specific areas of a surface by light is based on light exciting photoelectrons from the valence band to the conduction band of a semi-conducting silicon electrode that is in depletion. The technique has been utilised for the detection of ion release from single cells, combined with fluorescence microscopy to show how the cells respond to drug treatment. Further work in this area will involve using LAE platform to record intrinsic noise that is generated by cells for a multimodal characterization as well as utilising the platform for modulation of cell's functions through application of external signals.





### (b) Biodegradation of implant materials

Bioresorbable metallic implants have introduced a new paradigm in regenerative surgery treatment, in particular as alternative to conventional cardiovascular stent and bone fixation materials. For the implant to be successful, it requires to degrade and dissolve in the surrounding media gradually as the supported tissue heals. The degradation rate needs to be fine-tuned to match the rate at which the tissue regains mechanical stability.

New projects in this area will investigate electrochemical techniques and sensors to monitor the onset and kinetic of degradation and the various physico-chemical processes taking place in real time. Surface modifications are often performed on the biomedical implants to improve chemical/electrochemical stability, wear resistance, surface texture and biocompatibility. Of particular interest is electrochemical noise method to enable non-intrusive and real time assessment of kinetic and thermodynamic of electrochemical reactions taking place at the surface of implant material. For realistic simulation of the degradation properties, a range of environmental parameters, e.g. load and cycles, ions/electrolyte, proteins, cell adhesion, mobility, temperature etc, will be taken into account to more closely mimic a biological environment.

### (c) Electrochemical noise as a platform for sensing

Current electrochemical sensors rely on three main protocols for data acquisition namely, amperometry, potentiometry and impedimetry. Electrochemical noise, unlike the three conventional techniques, relies on the naturally occurring chemical processes for data acquisition without the application of any external bias voltage or current and yet it yields all three amperometric, potentiometric and impedimetric information in a single measurement. This brings about new capabilities in biosensors. The intrinsic noise is originated from any physical and/or chemical phenomenon that changes the equilibrium state at the electrode/electrolyte interface and can cause fluctuation of current and potential. For example, adsorption of charged particles onto the electrode surface changes the electrical state at the electrical double layer that is manifested as a fluctuation in potential and can be experimentally measured. In the absence of external bias current and voltage, the electrode surface will re-establish equilibrium to balance out the extra charge and this will cause a fluctuation in current. This current fluctuation can also be measured. The aim for new projects in this space is to explore the implementation of electrochemical noise principle in developing biosensors with multimodal sensing capabilities.

I also welcome discussions on your project ideas around the electrochemistry and biomaterials. Feel free to send me an email or just drop by my office in Hilmer 746.

Interested students can find more context in the following references or by contacting Dr Jamali directly

Y. Yang, F.M. Mansfeld, M. Kavallaris, K. Gaus, R.D. Tilley, J.J. Gooding, Monitoring the heterogeneity in single cell responses to drugs using electrochemical impedance and electrochemical noise, *Chem. Sci.* 12 (2021) 2558–2566.

Y.B. Vogel, J.J. Gooding, S. Ciampi, Light-addressable electrochemistry at semiconductor electrodes: Redox imaging, mask-free lithography and spatially resolved chemical and biological sensing, *Chem. Soc. Rev.* 48 (2019) 3723–3739.

Kirkland NT, Birbilis N, Staiger MP, Assessing the corrosion of biodegradable magnesium implants: A critical review of current methodologies and their limitations, *Acta Biomater.* 8 (2012) 925–936.

L. Li, M. Zhang, Y. Li, J. Zhao, L. Qin, Y. Lai, Corrosion and biocompatibility improvement of magnesium-based alloys as bone implant materials: A review, *Regen. Biomater.* 4 (2017) 129–137.

S.S. Jamali, S.E. Moulton, D.E. Tallman, M. Forsyth, J. Weber, G.G. Wallace, Applications of scanning electrochemical microscopy (SECM) for local characterization of AZ31 surface during corrosion in a buffered media, *Corros. Sci.* 86 (2014) 93–100.

S.S. Jamali, D.J. Mills, A critical review on electrochemical noise measurement as a tool for evaluation of organic coatings, *Prog. Org. Coatings.* 95 (2016) 13–17.



## PROF. SCOTT KABLE

Level 1, Dalton Building (F12)

T: 9385 4713 E: [s.kable@unsw.edu.au](mailto:s.kable@unsw.edu.au)

### LASER PROBES OF CHEMICAL REACTIONS

- Use lasers to initiate photochemical reactions of relevance to atmospheric chemistry;
- Discover new chemical reaction mechanisms that cannot be explained by current theories;
- Discover new radicals using laser spectroscopy.

**It would be great to work with Honours students on the following projects:**

#### (a) The Atmosphere is on Fire!

**(Collaborators: Meredith Jordan, Sydney U.; Jenny Fisher, U. W/gong; Chris Hansen, UNSW)**

The poor state of our atmosphere is one of the most pressing issues facing society today. Everyone knows about the challenges of climate change. But did you realise that more people meet premature deaths from poor air quality than from either cancer or heart disease?



The chemical complexity of the atmosphere is extreme. More than 1 million organic molecules are suspected to be in the air. Add to the mix, solid and liquid aerosols, sunlight, and a range of pressure and temperature and you might understand the challenge in creating a model of our atmosphere that is accurate and predictive.

Fundamentally, models are only as good as the underlying chemistry that they contain.


Our contribution in this area is in the discovery of new chemical mechanisms. Not just a new reaction, but new classes of reaction that are relevant across large domains of atmospheric science. Our latest project is built on our discovery in the past 2 years of light-induced combustion reactions. Organic molecules react with  $O_2$  in a combustion environment because of the high temperature. This is an equilibrium environment where molecules are characterized by a temperature. But we discovered that organic molecules can absorb light and undergo combustion reactions in the atmosphere. Sunlight is acting like the match to induce these new reactions.

Your project can be experimental, computational, or modelling-based or any mix of the three. In the lab, you can use laser-based techniques to characterize light-initiated combustion in one target molecule.

a)



Computationally, you can determine the critical reaction pathways and critical energies for a number of target molecules and predict which will react and which will not. Using sophisticated atmospheric models, you can predict the impact of these new reactions on our understanding of atmospheric processes (see figure at left).




**(b) Weird chemistry – reactions that just don't go where they should. (Collaborators: Meredith Jordan, Sydney U., David Osborn, Sandia National Labs, USA)**

Since the 1930's, the concept of a transition state (TS) has formed the bedrock of chemical reaction theory. When the activation energy is very near the TS energy, the reaction becomes very slow and other unsuspected processes become competitive, even dominant. Over the past few years we have identified new chemical pathways never previously described.

*The "Roaming" reaction:* When a reaction is initiated near the energetic threshold, the products barely have enough energy to escape each other's influence. Here, they "roam" around each other and re-collide, forming new, unexpected products. Roaming has been described as the most important new fundamental reaction class discovered in the past 20 years and new aspects of how roaming works are still being discovered.

This project will explore quantum resonances in roaming. We are trying to learn how quantum aspects, such as interference and resonance, influence roaming outcomes. The project would suit a student with a strong background in physics and can be experimental or computational (or both) in nature. For a longer description of the chemical physics of this reaction have a look at this [video](#):



**(c) Radicals in the atmosphere, combustion and space (Collaborator: Tim Schmidt, UNSW)**

Free radicals are key intermediates in all complex chemical environments. OH radical attack is the first step in the "processing" of nearly all atmospheric compounds. Radicals are found all through the interstellar medium and propagate flame chemistry. Of course you cannot buy a bottle of radicals from Aldrich (!) so you have to make them *in situ* and study them before they react with anything.

This is an inherently spectroscopic project where radicals are made in a vacuum using a variety of methods in our lab. They are characterised by a suite of spectroscopic techniques to determine their structure and chemical properties. Many times, you would be "seeing" a chemical species never seen before.

This project will involve the formation, measurement and characterization of a radical, chosen depending on your interest (space, combustion, atmosphere). A variety of laser spectroscopy techniques will be used to measure its properties. In concert with computational methods the structure of the radical can be worked out in fine detail.

<sup>1</sup> [https://www.dropbox.com/s/ai9y1vyti3no8b9/Science\\_marketing\\_compressed2.mp4?dl=0](https://www.dropbox.com/s/ai9y1vyti3no8b9/Science_marketing_compressed2.mp4?dl=0)





**A/PROF. KRIS KILIAN**

Level 7, Hilmer Building (E10)

T: 9385 4666 E: [k.kilian@unsw.edu.au](mailto:k.kilian@unsw.edu.au)

## **BIOINSPIRED MATERIALS, TISSUE ENGINEERING, MECHANOCHEMISTRY**

Inspired by biological materials, we integrate nano- and micro- fabrication techniques with synthetic chemistry to mimic the physical and chemical properties of the cell and tissue microenvironment. Much of our work is motivated by a dynamic model of the microenvironment where the interplay between chemical cues (extracellular matrix composition), physical cues (geometry, mechanics and topography) and biological cues (paracrine and juxtacrine signals) guides mechanochemical signalling to influence cellular identity, fate and function. Our broad aims are to:

- 1) Develop model synthetic platforms for cell biology research and high-throughput drug development.
- 2) Use the output from 1 to design clinically relevant biomaterials that direct a functional outcome (e.g. synthetic organoids, model tumours, tissue repair and replacement).

Our work is necessarily interdisciplinary; honours students will gain practical experience in synthetic chemistry, materials fabrication (bioprinting, lithography), and cell and molecular biology techniques.

**It would be great to work with Honours students on the following projects:**

### **(a) Directing the chemistry/architecture of 3D extruded soft biomaterials**

3D printing of cells and tissues is limited by issues with complex bioink formulation, segregation of different cell types, cell viability during prolonged printing, and difficulty recreating complex architectures observed in nature. New methodologies to quickly fabricate cell-laden tissue structures with well-defined segregated populations has the potential to be transformational to tissue engineering. We are exploring the extrusion of multiple hydrogel materials of tissue-mimetic composition (Fig. 1; *Advanced Materials* 2015). By incorporating chemical handles in the polymers, microfluidics will be employed to establish gradients of multiple cell binding ligands. We aim to develop co-culture formulations for translation to a 3D printer to direct write the cell-laden extruded hydrogels within a 3D bulk poly(ethylene glycol) hydrogel.




Fig. 1. Extrusion of cell-laden chemically modified alginate (*Adv. Mater.*, 2015)

### **(c) Ceramic Omnidirectional Bioprinting in Cell-laden Suspensions (COBICS)**

The integration of hierarchical structure, chemistry, and functional activity is important for building bone mimics for tissue engineering. Bone is a highly mineralized tissue with an organic matrix containing bone residing cells. Inspired by bone biomineralization, we have developed a novel apatite-transforming ink that can be




Fig. 2. Ceramic Omnidirectional Bioprinting in Cell Suspensions (Romanazzo et al., *Adv. Funct. Mater.*, 2021)

printed into a supportive microgel matrix with living cells (Figure 2; Adv. Funct. Mater. 2021). Using this technique, complex bone-mimicked constructs are made at room temperature without requiring invasive chemicals or high temperatures. This new strategy for fabrication of synthetic bone has scope for creating custom microenvironments for disease modeling and 3D printing bone directly into a patient. We currently have projects exploring new ink formulations to modify the inorganic and organic part to improve printability and healing.

#### (d) Synthetic tumours for cancer nanomedicine development

Our interests in cellular “plasticity” has led us to cancer, where we believe progression and metastasis is a consequence of dynamic interactions in the tumour microenvironment that promote intravasation, extravasation and colonization. We microengineered small populations of melanoma cells across hydrogels and were able to uncover an intriguing role for geometry at the perimeter of these micro-tumors in orchestrating the activation of a cancer stem cell (CSC) state (Figure 3; Nature Materials 2016). This is important because these CSC-like cells are believed to be the root cause of recurrence and metastasis, the primary causes of suffering in cancer. Our vision for the future of this work is the integration of our model systems into autonomous tissue-mimetic architectures, for therapeutic development on patient derived cells. We have several new directions in need of students including: *new hydrogel chemistry and fabrication techniques, exploring spatiotemporal uptake of nanoparticles*, integration of multiple different cell types.




Fig. 3. Interfacial curvature will guide the activation of a stem-like state (Lee et al., *Nat. Mater.*, 2016)

#### Bringing mechanochemical activity to hydrogels

Hydrogels in tissue are viscoelastic materials that are continuously remodelled, and undergo dynamic changes in chemistry. Recreating dynamic chemistry in the laboratory most often involves incorporation of stimuli-responsive motifs, or secondary polymerization routines. We are investigating chemical linkages in hydrogels that are dynamic in response to stimuli including: temperature, pH, enzymatic activity and force. We are particularly interested in approaches where the chemistry can be modulated through applied compression or tension. Recently, we synthesised mechanophores that are “flex-activated” and demonstrated how




Fig. 4. Compression or tension triggers molecule release in double network hydrogels (Jayathilaka et al., *Chem. Commun.* 2021)

compression and tension will trigger a retro Diels-Alder reaction to stimulate molecule release double network hydrogels (Fig. 4; *Chem. Commun.* 2021). We are looking for honours students interested in synthetic chemistry and polymer science to build the next generations of molecule releasing hydrogels for use as dynamic coatings and scaffolds for biotechnology and tissue engineering.

Junmin Lee, Meredith N. Silberstein, Amr A. Abdeen, Sang Yup Kim, and Kristopher A. Kilian, Mechanochemical functionalization of disulfide linked hydrogels, *Materials Horizons*, 2016, 3, 447-451  
 Joshua M. Grolman, Douglas Zhang, Andrew M. Smith, Jeffrey S. Moore, and Kristopher A. Kilian, Rapid 3D extrusion of synthetic tumor microenvironments, *Advanced Materials*, 2015, 27 (37), 5512-5517  
 Amr A. Abdeen, Junmin Lee, N. Ashwin Bharadwaj, Randy H. Ewoldt, and Kristopher A. Kilian, Magnetoactive hydrogels for temporal modulation of stem cell activity, *Advanced Healthcare Materials*, 2016, 5 (19), 2536-2544.  
 Junmin Lee, Amr A. Abdeen, Kathryn L. Wycislo, Timothy M. Fan, and Kristopher A. Kilian, Interfacial geometry dictates cancer cell tumorigenicity, *Nature Materials*, 2016, 15, 856-862.



## DR. DONG JUN KIM

Level 7, Room 718, Science and Engineering Building (SEB, E8)

E: [dongjun.kim@unsw.edu.au](mailto:dongjun.kim@unsw.edu.au)

### SUPRAMOLECULAR ENERGY MATERIALS CHEMISTRY

We are a young research group which focuses on developing next-generation energy storages and supramolecular chemistry system. Our research approach is based on combining synthetic chemistry, electrochemistry, and materials science principles to develop advanced energy storage devices, in particular, rechargeable batteries. Additionally, we expect to conduct interdisciplinary research and establish collaborations with other research groups. Please feel free to contact me if you need any further information.


**It would be great to work with Honours students on the following projects:**

#### (a) Designing rechargeable Al-ion batteries

Aluminium is the third most abundant element in the Earth's crust. It has one of the highest theoretical volumetric capacity ( $8056 \text{ mAh mL}^{-3}$ ) on account of its multiple redox states. Therefore, developing rechargeable batteries utilising aluminium offers a golden opportunity for delivering a high energy to cost per price. The development of Al-ion batteries has not reached a stage yet. It has proved difficult to design an electrode material that can reversibly intercalate Al-ions, because the multivalent nature of aluminium is accompanied by significant structural changes, resulting in a rapid capacity fading.

Recently, we demonstrated one of the first rechargeable Al-ion batteries. Our approach was the utilisation of the triangular macrocyclic compound, which form layered superstructures resulting in the reversible insertion and extraction of an aluminium complex. This architecture exhibits an outstanding electrochemical performance along with superior cycle life.

The overarching goal of this Honour project is unlocking the full potential of rechargeable Al-ion batteries, by combining synthetic chemistry and battery engineering. Based on the large selection and synthetic versatility of various organic molecules, the redox-active compounds based rechargeable Al-ion batteries could provide a promising starting point for developing affordable large-scale energy storage applications.




**Figure 1.** Graphical representation of the macrocyclic building blocks into nano-channels.

**(b) Designing Molecular Dual Pump** (in collaboration with Prof. Sir Fraser Stoddart in Northwestern University)

Artificial molecular machines have received an increasing amount of attention over the past few decades. They have the unique ability to generate directional motion of components within their molecules by energy inputs or external stimuli. In our group, we have developed chemically- and electrochemically-driven molecular pumps in order to trap cyclobis(paraquat-p-phenylene) (CBPQT<sup>4+</sup>) rings on a collecting chain. A dual molecular pump can generate unidirectional motion along the dumbbell component using chemical reagents or electricity without accumulating waste products. By attaching a steric stopper at the end of the dual pump, the dumbbell will contain two collecting chains, making it possible to synthesize a [3]rotaxane sequentially.

This dumbbell consists of two pumps joined in series in a head-to-tail fashion with the first collecting chain located in the middle of them. It can be synthesized from the components that have already employed in the Stoddart Group. The second collecting chain is terminated by a bulky stopper. The target molecule will be produced using a click reaction.

Artificial molecular machines can be powered by chemical redox reactions where Zn (reductant) and NOPF<sub>6</sub> (oxidant) are used alternately. This in-series molecular dual pump can also be operated simply by the oscillation of two constant potentials (−0.7 V for reduction and 1.4 V for oxidation) in a controlled electrochemically powered process. The dumbbell contains two collecting chains which can accommodate at least two CBPQT<sup>4+</sup> rings. Heterotopic co-constitutional isomers of the [3]rotaxane could be generated by using CBPQT<sup>4+</sup> and a substituted CBPQT<sup>4+</sup> ring. By manipulating the pumping conditions with free and substituted CBPQT<sup>4+</sup> rings in the bulk solution, two different rings will be installed onto the dumbbell sequentially from the head to the tail.



**Figure 2.** Structure the molecular dual pump and pumping rings onto the collecting chain chemically and electrochemically.



## DR. SARA KYNE

Level 1, Dalton Building (F12)

E: [s.kyne@unsw.edu.au](mailto:s.kyne@unsw.edu.au)

### SCIENCE EDUCATION RESEARCH and SUSTAINABLE CHEMISTRY SYNTHESIS

---

We are a new group that joined UNSW in 2023. Please feel free to contact me to discuss your own research interests and potential projects in more detail.

- Our science education research encompasses inclusive learning environments, data analytics for improving student experiences and systems thinking for context-based learning
- Our laboratory-based research centres on catalysis and green chemistry, including reaction design and mechanistic understanding
- Students with chemistry, education, social science and other backgrounds are welcome, as we believe diversity is key to tackling interdisciplinary challenges
- We have ongoing close collaborations with the UK, Europe, the Philippines and across Australia

**It would be great to work with Honours students on the following projects:**

#### **(a) Promoting inclusive learning environments**

(in collaboration with the University of the Philippines Open University, The Philippines)

There is increasing diversity of learners in modern education with different backgrounds, and we are exploring effective ways to create inclusive learning environments for all students. The aim of this research is to benefit students through their access to equitable, quality science education. We are designing proactive and flexible teaching approaches and learning resources, guided by the Universal Design for Learning framework.<sup>1</sup> This framework promotes equity by intentionally incorporating teaching and learning strategies that support marginalised students, at the same time benefiting all students. To inform and evaluate our approaches, we are harnessing big data and the emerging fields of data analytics. Data science can be used to establish clear metrics to evaluate and improve inclusivity in learning environments.



#### **(b) Sustainability and green chemistry through systems thinking and context-based learning**

(in collaboration with Dr. Martin Peeks, UNSW)

We aim to future-focus the curriculum, to prepare undergraduate students to tackle global challenges confronting modern society. To achieve this, we are designing and implementing context-based learning activities that link fundamental chemistry concepts with modern society using a systems thinking approach. Our goal is for students to use chemistry to change their individual actions and develop multidisciplinary solutions to sustainability's "wicked problems". In addition to measures such as academic performance, student satisfaction and




**THE GLOBAL GOALS**  
For Sustainable Development

engagement, the impact of these activities will be evaluated based on shifts in students' motivation and changes in perceptions towards sustainability as influenced by the enhanced curriculum.

### (c) Authentic assessment practices for learning

(in collaboration with Dr. Siobhán Wills, UNSW and the University of Sydney)


In recent years, many universities in Australia and worldwide have seen major changes to their teaching and learning delivery. This includes assessment strategies, including summative final examinations. Together with the University of Sydney, we are undertaking a multi-institution exploration of first-year undergraduate examinations in STEM subjects comparing end-of-semester examinations since 2019 to determine the pedagogical changes that have occurred. Outcomes from our data analysis will inform best practices that balance university and student expectations to deliver authentic assessment experiences. Our research will enable us to make meaningful recommendations for best practice in Australian STEM subjects for summative examinations, including design that considers both technological as well as pedagogical aspects required to deliver effective assessments.



### (d) Iron catalysed sustainable reactions


(in collaboration with Prof. Les Field, UNSW and the University of Cambridge, UK)

First row transition metals have found applications as homogenous and heterogeneous catalysts that offer improved sustainability compared with precious metal catalysts. Iron can act as a powerful redox active catalyst for both single- and two- electron transfer processes, opening up a wide range of potential reactivities.<sup>2</sup> We are designing new iron-based synthetic methods for applications including polymerisation and depolymerisation, intra- and intermolecular radical reactions, and main-group bond formations. Our research investigates the mechanism of the catalytic cycles, aiming to identify key reaction intermediates and understand chemical, physical and electronic properties that impact catalysis.<sup>3</sup>




### (e) Sustainable photoredox catalysis

We are developing site-selective catalytic methods to transform simple sugars. These reactions use photoredox catalysts and visible light to promote carbon-carbon bond formation. Our research is investigating reaction mechanisms and how various reactive intermediates and metal species interact. This is important to inform the design of new, site-selective synthetic reactions of more complex sugars, ultimately upgrading waste biomass into high-value chemicals.



1. C. T. Reyes, G. A. Lawrie, C. D. Thompson and S. H. Kyne, "Every little thing that could possibly be provided helps": Analysis of online first-year chemistry resources using the universal design for learning framework. *Chem. Educ. Res. Pract.*, 2022, 23, 385–407. <https://doi.org/10.1039/D1RP00171J>
2. S. H. Kyne, G. Lefèvre, C. Ollivier, M. Petit, V.-A. R. Cladera and L. Fensterbank, Iron and cobalt catalysis: new perspectives in synthetic radical chemistry, *Chem. Soc. Rev.*, 2020, 49, 8501–8542. <https://doi.org/10.1039/D0CS00969E>
3. M. A. Farcaş-Johnson, S. H. Kyne and R. L. Webster, Dehydrocoupling polymerization: Poly(silyl ether) synthesis by using an iron  $\beta$ -diketiminate catalyst. *Chemistry – A European Journal*, 2022, 28, e202201642. <https://doi.org/10.1002/chem.202201642>





## PROF. NARESH KUMAR

Level 7, Science and Engineering Building (E8)

T: 9385 4698 E: [n.kumar@unsw.edu.au](mailto:n.kumar@unsw.edu.au)


### SYNTHETIC ORGANIC AND MEDICINAL CHEMISTRY

The main focus of the research undertaken in my group is the discovery and development of novel bioactive molecules. Naturally produced chemicals are of fundamental importance in biological systems. Such chemicals are used to mediate interactions across all levels of biological hierarchy. Very often such diverse molecules are produced only in minute quantities. New or innovative organic syntheses not only provide access to sufficient quantities of these molecules but also their analogues. The access to various structurally-related analogues allows full assessment of their biological activity and mode of action, and offers opportunities to develop new therapeutic leads. The research is multi-disciplinary in nature and involves a combination of synthetic organic chemistry, molecular modelling and biological screening.

#### (a) DESIGN AND SYNTHESIS OF NOVEL ANTIMICROBIAL AGENTS


##### Quorum Sensing Inhibitors

The emergence of multi-drug resistance in common human pathogens has highlighted the need to develop novel classes of antimicrobials for the treatment of human disease. A number of projects are available in this area focussing on a combination of organic synthesis, molecular modelling, and *in vitro* and *in vivo* antimicrobial screening. This project will develop novel antagonists of bacterial signalling pathways, which inhibit the regulatory quorum sensing communication pathways of bacteria, and will model the receptor-ligand interaction using the X-ray crystal structures of bacterial signal receptors e.g. *Pseudomonas* quinolone system (PQS).



##### New scaffolds for antimicrobial discovery

The majority of conventional antibiotics used today share a common feature in that they act on specific molecular targets. Having very well-defined targets, these drugs act with a high degree of selectivity, minimizing unwanted side effects. However, a major limitation of antibiotics targeting a single receptor is the ease with which resistance can be developed. The central aim of this project is to design novel small molecular antimicrobial peptide (SMAMP) mimics based on biphenyl scaffolds, which disrupt the normal functioning of the membranes of the bacterial cell, and as a consequence allow the development of antimicrobial agents with enhanced activity and the ability to bypass resistance mechanisms used by bacteria against other antibiotic types.



### Inhibitors of Bacterial Transcription Initiation

(in collaboration with A/Prof. Renate Griffith, UNSW and Prof. Peter Lewis, University of Newcastle)


The enzyme RNA polymerase (RNAP) that transcribes DNA into RNA is highly conserved across species. However, the factors that regulate the activity of RNAP are target-specific. Therefore, the unique interaction of sigma factors with RNAP in bacteria represents an ideal target for the development of small molecules that can specifically inhibit this interaction<sup>3</sup>. In this project new molecules that target these essential protein-protein interactions will be rationally designed and synthesized, and evaluated for their antimicrobial efficacy. These new small molecules would represent lead compounds for the development of new antibiotics.



### **(b) DEVELOPING ANTICANCER COMPOUNDS THAT ACTIVATE GLUCOSE OXIDATION**

(in collaboration with Dr Frances Byrne and A/Prof Kyle Hoehn, BABS, UNSW)


Cancer is a major burden of disease, affecting the lives of tens of millions on a global scale. A hallmark feature of nearly all cancer cells is their altered metabolism of glucose compared to non-cancerous cells. Relative to most normal cells, cancer cells use a greater proportion of incoming glucose for non-oxidative purposes including the production of building blocks for cell



division (lipid, DNA and protein), rather than oxidative pathways that produce carbon dioxide (CO<sub>2</sub>) in mitochondria. The goal of this proposal is to develop anticancer molecules that change cancer cell glucose metabolism to be more like that of non-cancerous cells. We have identified a small molecule that increases glucose oxidation and selectively kills cancer cells in vitro and in mice. The aim of this project is to generate new derivatives with enhanced activity and drug-like properties. The new compounds will be evaluated for anticancer activity in various cancer cell lines.

### **(c) DESIGN AND SYNTHESIS OF NOVEL HETEROCYCLIC SYSTEMS**

Flavones and isoflavones are two structurally related large and diverse groups of natural compounds with broad spectra of biological activities including antioxidant, anticancer, antiviral and anti-inflammatory properties. They are recognized as “privileged” medicinal chemistry molecular frameworks because they are commonly found in biologically active compounds that show drug-like characteristics. Rottlerin is a flavonoid isolated from the fruits of a medicinal plant, *Mallotus philippensis*. Our group has reported the successful synthesis of rottlerin via the acid-catalyzed reaction of 5,7,8-trimethoxyflavene. A number of projects are available in this area focussing on the design and synthesis of new azaflavone analogues of flavones and isoflavones in which the ring oxygen atoms are replaced by a nitrogen atom.







## DR. MARTINA LESSIO

Office: Room 124, Dalton Building

E: [martina.lessio@unsw.edu.au](mailto:martina.lessio@unsw.edu.au)

### COMPUTATIONAL MATERIALS SCIENCE AND CHEMISTRY FOR SUSTAINABILITY APPLICATIONS

Computer simulations are an essential tool to make high-impact discoveries in fields that are crucial to our sustainable future. In general, these types of simulations allow us to calculate properties of molecules and materials at the atomic scale, which can be too difficult to be measured by experiments. This information can be used to unravel the fundamental chemistry features of a system responsible for promising experimental observations and thus rationally guide experimental efforts towards optimizing those features for the application of interest.


**Research in my group focuses on using computer simulations to tackle a variety of sustainability issues, including the development of new water purification and plastic recycling technologies.** Additionally, I am eager to explore new application areas for computational chemistry, such as art conservation (see example project on the next page). Working on these projects will allow you to acquire/strengthen knowledge and skills in a variety of fields in chemistry, physics, and computer programming. Furthermore, most of the projects involve close collaboration with groups at UNSW and overseas (United States and Europe). Please don't hesitate to contact me to discuss possible projects in more details and/or your research interests. No prior knowledge of programming or computational chemistry is required.



#### Some of the projects currently available are:

##### (a) Computational Design of Metal-Organic Frameworks for Clean Water Harvesting and Water Purification


Access to clean water has been recognized as an essential human right by the United Nations. However, water contamination issues still exist and often render drinking water unsafe even in well developed countries. Developing cost-effective and efficient materials for clean water harvesting and polluted water treatment is necessary to ensure access to clean water for all. Metal-organic frameworks (MOFs) are promising materials for adsorption-based clean water technologies due to their extremely high surface area, the possibility to tune their selectivity by functionalizing their surface, and the



possibility to alter their pore size by choosing different building units. We use computational chemistry to aid the development of new materials based on MOFs for the adsorptive removal of heavy metals from water and for harvesting clean water from the air.

### (b) Plastic Waste Conversion into Useful Products using Transition-Metal Catalysts

The conversion of plastic waste into monomers and other useful chemicals is a promising avenue towards addressing the plastic waste issue and reducing the use of non-renewable resources to generate such products. Recent experimental studies have shown that transition metal catalysts can be used to perform this conversion at moderate temperatures and with good product control. This project uses computational tools to design improved catalysts.




Overall we expect the results of these projects to guide experimental efforts towards the synthesis of improved and more cost-effective catalysts for plastic waste conversion into useful products.

### (c) Computational Chemistry Meets Art Conservation: Design of Improved Surface Protective Treatments for Marble

Computer simulations are an established tool in the investigation of solid/liquid interfaces in many different fields ranging from materials science to biological applications. Solid/liquid interfaces are often the focus of art conservation efforts as solid artefacts are often exposed to harmful liquids. In spite of its great potential, the application of computational chemistry in the field of art conservation is still extremely limited.

In this project, we will use computational tools to investigate the chemical mechanism behind an innovative treatment for the protection of marble artefacts exposed to water. We will then use the acquired knowledge to develop improved protective treatments in close collaboration with conservation scientists at the University of Bologna, Italy.



Sassoni, E. *Materials* 2018, 11, 557



## DR. LAUREN MACREADIE

Dalton Building (F12)

E: [lauren.macreadie@sydney.edu.au](mailto:lauren.macreadie@sydney.edu.au)

### COORDINATION CHEMISTRY OF METAL-ORGANIC FRAMEWORKS

Metal-organic frameworks (MOFs) are porous coordination polymers built from the self-assembly of metal clusters and organic linkers. Through careful selection of these building blocks when designing our MOFs, materials with pre-determined properties can be synthesised. Our research area uniquely encompasses both organic and inorganic chemistry to develop interesting functional porous materials for catalytic hydrogen generation, gas storage and transport, negative thermal expansion and as MRI contrasting agents. Using diffraction methods from the Australian Synchrotron, we can piece together the structure-function relationships of these new materials and continually refine them for better performance.


**Skills acquired from all projects include:**

*X-ray single crystal and powder diffraction, synchrotron science, MOF synthesis, organic and inorganic synthesis and characterisation using NMR, photophysical and gas adsorption.... Among many others!*

**It would be fantastic to work with Honours students on the following projects:**

**(a) 3D-Linker MOFs for separations and storage**


Most MOFs are constructed using aromatic linkers, such as terephthalic acid ( $H_2bdc$ ), due to their low cost and well understood chemistry. Consequently, over 10,000 MOFs are made with only  $H_2bdc$ , giving a very poor representation of possible MOF environments. Our team works with rigid, 3D-linkers linkers such as cubane-1,4-dicarboxylic acid ( $H_2cdc$ ) and have discovered enormous potential in these systems. Due to the bulky nature of the cubane, more supramolecular interactions are possible between the host and guest systems. This project extends the investigation to other 3D-linkers which will exhibit exciting properties. This high impact project and involves multiple collaborations, and investigates different factors governing host-guest behaviours.



(Collaboration: Prof Omar Farha, Northwestern University, USA; and Dr Paul Savage, CSIRO)

**(b) Photoactive frameworks for water splitting or  $CO_2$  reduction**

Luminescent MOFs (LMOFs) are rapidly gaining interest due to their promise in a broad range of applications including chemical sensing, artificial photosynthetic catalysis and optoelectronics. Recently, we have found tuneable luminescence can be gained through modulation of linkers




with mixed functionalities and the incorporation of mixed metals (eg. Ru and Co). This project investigates increasing the luminescent lifetimes of phenanthroline based MOFs through varying the conjugation in the MOF linker. As an added bonus, MOFs constructed from these linkers lead to large pores which are ideal for gas storage.

(Collaboration: Prof Lyall Hanton, University of Otago, New Zealand)

### (c) Australian synchrotron high pressure X-ray diffraction to investigate negative thermal expansion (NTE) of 3D-Linker MOFs

Many materials exhibit positive thermal expansion with temperature. However, MOFs interestingly exhibit negative thermal expansion (NTE) – a phenomenon not often seen in materials. This is advantageous when trying to design zero thermal expansion materials which are highly sought after in industry. 3DL-MOFs exhibit enhanced NTE compared with aromatic frameworks due to the hyper-fast molecular rotor dynamics of the aliphatic cores. These rotor dynamics can be influenced by external pressure and temperature environments, further influencing the extent of NTE in these materials. This project will involve studying the response to external pressure of 3DL-MOFs through variable pressure and temperature X-ray diffraction studies run at the Australian Synchrotron using diamond anvil cells.


(Collaboration: Prof Cameron Kepert, University of Sydney; Prof. Stephen Moggach, UWA; Australian Synchrotron)



### (d) NanoMOFs as dual MRI contrasting agents/drug delivery agents

The highly porous nature of MOFs allows them to hold and deliver large payloads such as drugs and nutrients. Furthermore, exploiting the coordination polymer nature of MOFs means that a high amount of MRI active agent can be generated in a nanosized material, which can be tailored to target specific sites in the body. NanoMOFs exhibit numerous properties which make them ideal for biomedical applications. Their highly porous structures allow accommodation of high loadings of therapeutic and imaging agents and their controlled release, in addition to protection against enzymatic degradation.

This project investigates iron and gadolinium contrasting agents which can self-assemble to form NanoMOFs, and studies their controlled release at target sites in the body.



### (d) Other projects for your interest including agriculture remediation and CO<sub>2</sub> capture!

1. Macreadie, L.K., Mensforth, E.J., Babarao, R., Konstas, K., Telfer, S.G., Doherty, C.M., Tsanaktisidis, J., Batten, S.R., Hill, M.R. *Journal of the American Chemical Society*, 2019, 141, 3828.
2. Macreadie, L.K. Babarao, R., Setter, C.S., Lee, S.J., Qazvini, O.T., Seeber, A.J., Tsanaktisidis, J., Telfer, S.G., Batten, S.R., Hill, M.R. *Angewandte Chemie*, 2020, 59, 6090.
3. Macreadie, L.K., Qazvini, O.T., Babarao, R. *ACS Applied Materials & Interfaces*, 2021, 13, 26, 30885.



## DR. SHANNAN J. MAISEY

Level 1, Dalton Building (F12)

T: 02 9385 5967 E: [s.maisey@unsw.edu.au](mailto:s.maisey@unsw.edu.au)

### CHEMISTRY EDUCATION IN THE 21<sup>ST</sup> CENTURY

---

The UNSW chemical education group is interested in improving the learning outcomes and experience of Chemistry students and contributing to the chemical education research community. My personal research interests encompass how to integrate the global challenges facing science into chemistry curricula (systems thinking), the development and tracking of transferable skills of chemistry graduates and the role that technology must play on how we teach and interact with chemistry. Here are some of the projects available for honours/research project with me, though chemical education projects can be tailored and developed to suit you and your interests!

---

#### (a) Facing up to global challenges - Integrating systems thinking into Chemistry education

Keywords: Systems thinking, Global challenges, Mastery learning

The world is currently facing unprecedented challenges which are affecting all facets of life on earth. Climate change, Sustainability and the need for Renewable energy sources and storage are challenges which have chemistry at their core. A recent article in Nature Reviews Chemistry<sup>1</sup> served as a call to arms for chemistry educators to integrate systems thinking into chemistry curricula at all levels to empower our students with the knowledge and skills to face these challenges. Systems thinking is about putting chemical concepts into a real worlds context and showing how atoms and molecules (and the decisions we make with what to do with them) impact people's lives and our environment. There is very litter literature which describes systems thinking in chemistry which presents many exciting opportunities for your project from exploring the challenges of integrating it into chemistry curricula to finding out how student's viewpoints develop and change with a broader view of chemistry...it's an exciting time to be alive!


#### (b) UNSW Chemistry Graduates: Ready for Anything... But do they know that?

Keywords: Transferable skills, Work Integrated Learning, Micro-credentialing

Beyond an understanding of key concepts of chemical theory, Chemistry graduates require a unique set of transferable skills. UNSW Chemistry has recently introduced several exciting education developments designed to enhance the capabilities and skills of our graduates. We are interested in investigating the efficacy of these programs in the development of transferable skills as well as exploring how well our graduates can articulate their skills in a chemistry context (such as when applying for jobs or networking) and how we might develop an educational intervention to improve this.

(c) **There's an app for that! Pedagogical content knowledge in the age of technology**

Keywords: Digital Literacy, PCK, TPACK, Online learning, Blended Learning




Pedagogical content knowledge (PCK) theory recognizes that beyond the teacher's own understanding and knowledge of the content theory there is a surrounding body of knowledge to do with how students learn and process information specific to the theory being taught. In the age of technology, the way we interact with students has changed. Technological pedagogical content knowledge (TPACK) is the basis of effective teaching with technology, requiring an understanding of the representation of concepts using technologies; pedagogical techniques that use technologies in constructive ways to teach content; knowledge of what makes concepts difficult or easy to learn and how technology can help students overcome difficulties<sup>2</sup>.

How students engage with technology to learn is rapidly evolving. Like many other institutions, online learning is now one of our underpinning methods of teaching first year chemistry. What is not well understood is what strategies students are engaging to use and supplementing these materials to facilitate learning. Why are some formats preferred by students and how is this impacted by demographics? Is there potential to impact how effectively we can teach students chemistry by 'updating' our TPACK?

(d) **Research by Students: Developing an innovative program that facilitates high volume contributions to a newly designed urgently needed online spectroscopic database (collaboration with Laura McKemmish).**

Keywords: Spectroscopy, Django online Python databases, Citizen Science, Education/Outreach




This project has a bit of everything: programming, data science, spectroscopy and education.

This project enables high school and undergraduate students to contribute to an urgently needed online database, gaining valuable transferable skills, scientific knowledge and exposure to scientists and scientific research in a project linking research, teaching and outreach!

The Database: Update of 1979 Huber & Herzberg Constants of Diatomic Molecules, still cited once a day, into a modern online query-able database. This data is exceptionally useful in benchmarking quantum chemistry and predicting spectra for diatomics found across the universe for applications from monitoring to detection to creating the coldest molecules ever!

The Education Component: This 'research-in-schools' approach is part of a growing international movement including the US SEED program championed by UNSW staff member and Nobel Laureate Sir Fraser Stoddard. Here, we will investigate how to bring it to Australia, probably through the new "Science Extension" HSC course, through a thorough study of related approaches and interviews of high school teachers.

1. Mahaffy, P. G., Krief, A., Hopf, H., Mehta, G. & Matlin, S. A. Reorienting chemistry education through systems thinking. *Nature Reviews Chemistry* (2018). doi:10.1038/s41570-018-0126
2. Koehler, M. J., & Mishra, P. (2009). What is technological pedagogical content knowledge? *Contemporary Issues in Technology and Teacher Education*, 9(1), 60-70.



## DR. LAURA K. MCKEMMISH

Level 1, Dalton Building (F12)

E: [l.mckemmish@unsw.edu.au](mailto:l.mckemmish@unsw.edu.au)

### COMPUTATIONAL MOLECULAR SPECTROSCOPY FOR ASTROCHEMISTRY AND BEYOND

---

*Want to do research on a computer not in a lab? Feel constantly pulled between physics and chemistry? Love spectroscopy, quantum mechanics and energy levels? Or perhaps you want to utilise and strengthen your maths, programming and/or data science skills by exploring exciting molecular science applications from predicting spectroscopy to helping find aliens on exoplanets?*

**I am looking for keen students to undertake projects with customisable amounts of chemistry, physics, mathematics, programming, data science and education/outreach.**

During a research project with me, you can expect to develop and strengthen many key transferable and scientific skills such as Python, command line, power use of supercomputers and quantum chemistry programs, data science, data presentation, debugging and, perhaps most importantly, “Googling”.

---

*My major research focus is method development for and applications of computational molecular spectroscopy.*

#### **Looking for life and its molecular origin in space**

Keywords: *Computational Quantum Chemistry, Astronomy, Exoplanets, Spectroscopy, Supercomputers, Data Science, High Accuracy, High-throughput Calculations, Radio & Infrared Spectroscopy*

One of our group’s key motivations is to predict spectral data that is immediately useful, often for characterising unusual astrophysical environments including exoplanets and the interstellar medium. Sometimes this means very high accuracy  $\text{sub-cm}^{-1}$  predictions of rovibronic spectra of weird diatomics like TiO, using all the experimental data we can find. Other times, this means producing approximate data for thousands of molecules to identify strong absorbers and molecules that will be difficult to distinguish astrophysically.



The primary purpose of the data is to enable astronomers to confidently detect molecules in various astrophysical environments. The highest profile of these sought detections are of course biosignatures in the solar system (e.g. phosphine on Venus) and exoplanets. Almost as important are the searches for the origins of homochirality and life through searches for pre-biotic and chiral molecules in the interstellar medium.

On a more local level, this type of generated data is important for monitoring atmospheric composition and pollutants on local and global scales and in industrial plants. It can also be used to predict global warming potential of different molecular compounds (e.g. those proposed as replacements for CFCs).

## Machine Learning: Chemical Structure → Spectra

*Keywords:* Machine Learning, Data Science, Computational Quantum Chemistry, Supercomputers, High-throughput Calculations, Spectroscopy

Machine learning and “big data” science is starting to revolutionise many areas of chemistry, but one area hardly considered is spectroscopy. Can machine learning outperform quantum chemistry calculations in some (or all?) areas of modern computational molecular spectroscopy? The high-throughput data produced by my group provides a perfect training set for machine learning models to predict spectral properties from chemical structure without quantum chemistry, as a byproduct recreating organic chemistry infrared functional group tables.

## Rotational Spectroscopy of Pre-biotic Chiral Species: Experiment & Theory (collaboration with Chris Medcraft)

*Keywords:* Astrochemistry, Experimental Spectroscopy, Computational Quantum chemistry

Why pick? Do a project that combines experimental rotational spectroscopy with computational quantum chemistry predictions, focused on the rotational spectroscopy of a pre-biotic chiral molecule that may help tell scientists how life emerged. This project produces crucial high-accuracy astronomical data required for the upcoming Square Kilometre Array radio telescope and its precursors.

*Beyond this main body of work, other potential projects include:*

### Why is B3LYP/6-31G\* still so popular?

*Keywords:* Data extraction, Change theory, Computational chemistry, Qualitative research, Data analysis  
B3LYP/6-31G\* was the state of the art quantum chemistry method ... around the year 2000. Yet the widespread availability of better model chemistries (as benchmarked extensively), this older theory is still used extensively, especially for organic chemistry applications.

In this project, we will investigate the choices users make: what, how & why. This will be correlated to data on how method developers try to reach potential users. The data will be collected via interviews, surveys and parsing online data sources and analysed using the lens of change theory.

### Finding Illegal Drug Analogues using Cheminformatics (collaboration with Brynn Hibbert).


*Keywords:* Python, Application, Cheminformatics, Algorithm Design

Replacing a hydrogen with a fluorine atom often does little to affect the biological function of a molecule, so lawmakers need to ensure that molecules that are similar to illegal drugs are also illegal. But are the current laws too widespread – most critically, do they limit potential pharmaceutical medicines? In this project, you will enumerate illegal drug analogues and consider the implications of this law.

### Evaluating high-school outreach (collaboration with Shannan Maisey).

*Keywords:* Citizen Science, Education/Outreach/Teaching, Science Education, Evaluation

NSW Year 12 students have the opportunity to engage with a one-unit Science Extension course, where they pursue an independent research project ideally in collaboration with university researchers. At UNSW, we have developed SciX as a pathway to ensure equitable and widespread access to university research and researchers, and want your help in establishing and evaluating this programme's effect on the PhD student mentors, high school student researchers and other stakeholders.







## PROF. JONATHAN MORRIS

Level 2, Dalton Building (F12)

T: 9385 4733 E: [jonathan.morris@unsw.edu.au](mailto:jonathan.morris@unsw.edu.au)

### SYNTHETIC AND MEDICINAL CHEMISTRY


- The Morris group are focused on the development of organic molecules that can be used in biomedical research.
- They use their expertise in synthetic organic chemistry to access biologically active small molecules. Once an efficient strategy is developed, investigations into how these molecules interact with biological systems can be initiated.
- Being able to synthesise new small molecules in an efficient manner is critical and as such, the focus is on developing strategies to prepare these valuable materials and generate analogues that have improved potency and selectivity.
- Work on these areas leads to a number of collaborations with biomedical researchers where students can become involved in the understanding the biology.

**It would be great to work with Honours students on the following projects:**


#### (a) Modulation of RNA splicing using small molecule kinase inhibitors

The control of the fundamental biological process of alternative RNA splicing is an emerging method for treating diseases such as aged macular degeneration and cancer. It has been established that by controlling the phosphorylation of key proteins in the spliceosome, it is possible to alter RNA splicing and generate particular protein isoforms. The Morris group is actively engaged in the development of small molecules that can do this, and this is achieved by targeting the protein kinases that mediate the phosphorylation of the splicing factors.

Our recent publication (*ACS Chem. Biol.*, **2017**, *12*, 825) describes how we have developed a new class of kinases inhibitor that selectively inhibits the kinase SRPK1 and has led to the identification of a series of molecules that are currently being developed as an eye drop treatment for aged macular degeneration in collaboration with Exonate. The recent phase 1b/2 clinical trial has confirmed that this strategy is therapeutically viable. This class of inhibitors have also been used by our biological collaborators at the University of Nottingham in their investigations into cancer biology and pain modulation.



This work originated from earlier work on the synthesis of a natural product. Variolin B is a member of a unique class of marine alkaloids isolated from an extremely rare Antarctic sponge. It is no longer available from its natural source. The Morris group have devised a synthesis of variolin B that has restored access to the material and allowed further biological studies to be carried out. From this work it has been established that variolin B is a potent kinase inhibitor and represents an important scaffold for the further development of improved kinase inhibitors. A range of analogues have been developed that are more selective inhibitors of certain kinases, as well as have better properties such as solubility.




variolin B

Building on this work, we are engaged in the development of a number of different chemical scaffolds to inhibit the kinases that regulate alternative splicing. The aim is to develop selective inhibitors of the various RNA splicing kinases (the CLKs, DYRKs and SRPKs), with appropriate drug-like properties so they can be used as chemical probes to help understand the role these important kinases have on biological systems. A combination of synthesis and structure-based drug design is used to do this work, with students able to use Schrodinger and Cresset software to aid their design work.



**(b) Developing the AAL(S) Scaffold for Therapeutic Applications (with Prof Nigel Turner (Victor Chang CRI, UNSW Sydney), Prof Alaina Ammitt (UTS) and Dr Nikki Verrills (Newcastle))**

Ceramide synthase (CerS) and protein phosphatase 2A (PP2A) are two enzymes that play a critical role in the regulation of multiple cellular signalling processes. The malfunctioning of these two enzymes has been found to have implications in diseases such as cancer, diabetes, asthma and neurological diseases including Alzheimer's disease and stroke. Little is known about the biological mechanism of these enzymes and in particular, how they cause such diseases. To gain insight into these biological processes, the CerS and PP2A binders, FTY720 and AAL(S), will be used to explore the binding site of both enzymes and allow the identification of chemical probes which can be used to develop an understanding of the biological mechanisms of these complex diseases.



**(c) Designing drugs that cross the blood-brain barrier to treat childhood cancers (with Prof Matt Dun (Newcastle))**

Diffuse intrinsic pontine glioma (DIPG) is a highly aggressive brain tumour primarily affecting children. This disease is universally fatal, with critical location in brainstem ruling out surgical intervention. To date, there has been little progress made on developing therapeutics that target DIPG. One of the key challenges that arises is the inability of most small molecule drugs to cross the blood-brain barrier (BBB). Working with Prof Matt Dun (University of Newcastle) we are focussed on improving the brain-penetrant properties of potent anti-cancer drugs using a computer-aided multiparameter approach to predict which chemical changes will lead to improved permeability. Thus far we have demonstrated that simple chemical modifications can greatly improve BBB permeability while retaining anti-cancer activity. We are actively working on a series of chemical scaffolds that encompass a range of different synthetic approaches.



**(d) Disrupting the actin cytoskeleton to target breast cancer**

Actin filaments are the structures responsible for the shape and locomotion of cells. While cellular migration is essential for biological processes such as tissue repair, uncontrolled migration is implicated in cancer cell migration and metastasis. Early actin-targeting therapeutics have struggled with selectivity between healthy and cancerous cells leading to problems with toxicity. In this project, the Morris group are working in collaboration with Dr David Croucher and Dr Sharissa Latham (Garvan Institute of Medical Research) improving the drugs' potency and drug-like properties.



## DR. VINH NGUYEN

Level 2, Dalton Building (F12)


T: 9385 6167 E: [t.v.nguyen@unsw.edu.au](mailto:t.v.nguyen@unsw.edu.au)

### ORGANOCATALYSIS AND CHEMISTRY OF UNUSUAL MOLECULES

Nguyen's group has several Honours projects focusing on the development of novel organocatalytic systems or unusual molecules and applications of those in synthetic organic chemistry.

#### (a) Project NTV1 - Tropylium Ion as Chromophore for Organic Dyes

Tropylium ion is an unusual non-benzenoid aromatic system with  $6\pi$ -electron 7-carbon-ring structure.<sup>[1]</sup> Recent synthetic advances by our group have made this unique species much more accessible and understood, allowing us now to start to utilize it for a wide range of applications in organocatalytic chemistry<sup>[2-5]</sup> and photochemistry. This project will further investigate our recent findings that tropylium can be used as a versatile chromophore for a family of very interesting organic dyes and luminescent materials for *metal and pH sensing*. As some aspects of this project are confidential, students are encouraged to discuss with Vinh in person about this project.




#### (b) Project NTV2 - N-Heterocyclic Olefins as Novel Organocatalysts

Recently, N-Heterocyclic Olefins (NHOs, see scheme) have emerged as a new class of valuable reaction promoters with interesting action mechanisms. These compounds can be conveniently produced from commercially available precursors in one step. NHOs were originally targeted as a series of active agrochemicals in the 1970s, but they slowly revealed to be a far more interesting compound family. Due


to the donating ability of the two nitrogen atoms, the exocyclic C-C double bond is very electron-rich and strongly polarized. This interesting feature of NHOs offers multinucleophilic reactivity over the ketene amination frameworks.<sup>[6]</sup> Due to the strong nucleophilicity of the  $\alpha$ -carbon, NHOs can act as strong Lewis/Bronsted bases.<sup>[7-9]</sup> This project will focus on synthesizing a family of NHOs, estimating their basicity and applying them as organocatalysts to promote **environmentally friendly chemical processes**. Students are encouraged to discuss with Vinh in person about this project.

### N-Heterocyclic Olefins (NHOs)



### (c) Project NTV3 - Tropylium-Based Host-Guest (collaboration with A/Prof Pall Thordarson)

This project will explore the potential of tropylium-bearing systems in host-guest chemistry in **collaboration with A/Prof Pall Thordarson's group**. The electron-deficient nature of tropylium moiety makes it particularly attractive for the binding and sensing of small and medium-sized biologically important anions such as chloride, phosphate and carbonates. We propose the synthesis of tropylium-based macrocycles (see figure) as the starting point for this project, which will represent a new platform in supramolecular chemistry. Please also see Thordarson's Honours projects for more details.



### References

- [1] D. J. M. Lyons, R. D. Crocker, M. Blümel, T. V. Nguyen,\* *Angew. Chem. Int. Ed.* **2017**, 56, 1466-1484. <http://dx.doi.org/10.1002/anie.201605979>
- [2] T. V. Nguyen,\* A. Bekensir, *Org. Lett.* **2014**, 16, 1720-1723. <http://dx.doi.org/10.1021/ol5003972>
- [3] T. V. Nguyen,\* M. Hall, *Tetrahedron Lett.* **2014**, 55, 6895-6898. <http://dx.doi.org/10.1016/j.tetlet.2014.10.100>
- [4] T. V. Nguyen,\* D. J. M. Lyons, *Chem. Commun.* **2015**, 51, 3131-3134. <http://dx.doi.org/10.1039/C4CC09539A>
- [5] Demelza J. M. Lyons, Reece D. Crocker, Dieter Enders, Thanh V. Nguyen,\* *Green Chem.* **2017**, in press (DOI = 10.1039/C7GC01519D). <http://dx.doi.org/10.1039/C7GC01519D>
- [6] R. D. Crocker, T. V. Nguyen,\* *Chem. Eur. J.* **2016**, 22, 2208-2213. <http://dx.doi.org/10.1002/chem.201503575>
- [7] M. Blümel, J.-M. Noy, D. Enders, M. H. Stenzel, T. V. Nguyen,\* *Org. Lett.* **2016**, 18, 2208-2211. <http://dx.doi.org/10.1021/acs.orglett.6b00835>
- [8] M. Blümel, R. D. Crocker, J. B. Harper, D. Enders, T. V. Nguyen,\* *Chem. Commun.* **2016**, 52, 7958-7961. <http://dx.doi.org/10.1039/c6cc03771b>
- [9] Ugur Kaya, Uyen P. N. Tran, Dieter Enders, Junming Ho, Thanh V. Nguyen,\* *Org. Lett.* **2017**, 19, 1398-1401. <http://dx.doi.org/10.1021/acs.orglett.7b00306>




## A/PROF. GIANCARLO PASCALI

Level 1, Dalton Building (F12)

E: [g.pascali@unsw.edu.au](mailto:g.pascali@unsw.edu.au)

[gianp@ansto.gov.au](mailto:gianp@ansto.gov.au)

### INNOVATIONS IN RADIOPHARMACEUTICALS AND RADIOCHEMICAL TECHNOLOGIES



Credits: Dr. Mitch Klenner

Radiopharmaceuticals are becoming increasingly important for the management of many diseases, due to their extraordinary sensitivity and specificity. However, still many challenges are existent to deliver such high-value molecules, and they span from the initial molecular design, to efficient methods for radiolabelling and radioisotope purification, and to reliable process automation. Added to more standard radiopharmaceutical and radiolabelling development ideas, below are some examples of cross-disciplinary projects, that would anyway be tailored around student interests and desired skillset development.

#### (a) 3D-printed microfluidic reactors

(in collaboration with David Zahra, ANSTO)

Microfluidic systems have found multiple applications in the biochemical field, but are still not utilized extensively in the chemical reaction field. This is mostly due to the reduced availability of correct performance to be achieved for a chemical reaction (e.g. solvent resistance, pressure and temperature rating). However, new materials and advanced processes are now available for 3D-printing, and few works are demonstrating the feasibility of chemical reactions in microfluidic reactors manufactured in this way. In addition, it is demonstrated that radiolabelling reactions can happen with improved efficiency in the microfluidic environment; this fact, joined to the possibility of fast prototyping and single-use approaches (i.e. reduction of cross-contamination and radioactive waste), make 3D-printing of radiolabelling reactors a very attractive option.



Credits: David Zahra

This project will explore this possibility, mainly using non-radioactive assessment, but ultimately testing the best results in model radioactive scenarios. The knowledge acquired can be translated in the development of other Lab-on-Chip systems.


**Skills:** 3D-drawing software, 3D-printing equipment and process, functional test for microfluidic chips, instrumental analytical techniques (e.g. HPLC, GC), basics of CT imaging, basics of radiolabelling reactions.

**Key reference:** FOMSy: 3D-printed flexible open-source microfluidic system and flow synthesis of PET-tracer; *Journal of Flow Chemistry*, 13, 247–256 (2023).

## (b) Separation of radioisotopes by liquid/liquid flow extraction

(in collaboration with Elizabeth Tondl, ANSTO)

The production of radioisotopes is the first step towards manufacturing radiopharmaceuticals. In many cases, it requires bombardment of a defined target material to induce the nuclear transformation, the separation of the desired radioisotope from the bulk target material, and its formulation into a form usable for successive radiolabelling steps. There is an ongoing need of novel and more efficient methods to achieve such processes, linked to the increasing clinical demand of radiopharmaceuticals. Traditional separation methods were based on liquid/liquid extraction (LLE); however, current industrial (and also research) methods are based on resin-based separation, due to its ease of automation. On the other hand, this approach typically features variability of resin batches and packing, use of large volumes of eluents (that need dedicated radioactive waste storage), and challenges in scaling-up. In recent years, few systems have become available that allows LLE to be performed at the continuous flow regimen, and with capability to be easily integrated in automated systems. Therefore, there is a substantial potential to revisit the traditional LLE methods in radioisotope separations, possibly leading to processes with



Credits: Zaiput Flow Technologies

improved separation efficiency, better chemicals control, reduced waste and easy scale-up. This project will assess this concept using non-radioactive chemicals and conditions that mimic real-life concentrations, and the most performing systems will be tested with radioactive mixtures from real-life bombardments. The knowledge acquired can be translated to the integration of online LLE in flow chemistry processes for other synthetic chemistry applications.

**Skills:** *Hydrometallurgy basics, synthesis and/or purification of ligands, flow chemistry concepts, unbiased optimization methods (DoE, ML, AI, algorithms), radioisotope separation basics.*

**Key reference:** *Recovery of Gallium-68 and Zinc from HNO<sub>3</sub>-Based Solution by Liquid-Liquid Extraction with Arylamino Phosphonates; Molecules, 27, 8377 (2022).*

---



## **DR JOHN DOAN**

T: 9382 2200 E: [john.doan1@unsw.edu.au](mailto:john.doan1@unsw.edu.au)

### **Radiopharmaceutical Development**

---

Radiopharmaceutical science is a multidisciplinary field encompassing chemistry, physics and biology. It is the science of incorporating a suitable radionuclide into a pharmaceutical or other biologically active molecule in vivo physiological or biochemical processes. The resulting radiopharmaceuticals are used in the diagnostic imaging or therapy of patients with various diseases.

I have an interest in the development of radiopharmaceuticals with potential clinical applications in various fields including oncology and neurology. My role at the Department of Nuclear Medicine and PET, Prince of Wales Hospital is to provide the radiopharmaceutical clinical service for diagnosis of various diseases.

I have recently been appointed as a Conjoint Lecturer and a National Imaging Facility Fellow and I am seeking potential students to work on projects that could enhance the growing field of Radiopharmaceutical Sciences.

---



## DR. MARTIN D. PEEKS

Room 219, Level 2, Dalton Building (F12)

E: [m.peeks@unsw.edu.au](mailto:m.peeks@unsw.edu.au)


### SUPRAMOLECULAR & ORGANIC MATERIALS CHEMISTRY

Our research is concerned with understanding the nature of electronic communication and conjugation and using these principles to make interesting new molecules and assemblies. In doing this we have two real goals: making molecules that have useful properties, and those that help us learn something new and fundamental about chemistry. The overriding goal of all the projects is to give you the opportunity to **develop a broad set of research skills**: synthesis, computational chemistry, and in-depth analytical or photophysical studies, depending on your interests. There are many options for collaboration with other groups both at UNSW and overseas.

**It would be great to work with Honours students on the following projects:**

#### (a) Pushing the limits of $\pi$ -conjugation, aromaticity, and antiaromaticity

$\pi$ -conjugated molecules are like tiny little wires because they can delocalize electrons very effectively. Aromatic molecules are perhaps the archetypal  $\pi$ -conjugated molecules – things like benzene! They've been studied for more than 150 years, but much remains to be learned about aromatic and antiaromatic, as well as more unusual, molecules. For example, we recently reported the synthesis of the **largest known aromatic and antiaromatic molecules**.<sup>1</sup> In general we are interested in looking at new chemical structures which exhibit improved – or just unusual –  $\pi$ -electron delocalization and these are several projects available along these lines. We look at the effect of molecular structure on electronic delocalization and resulting properties like light emission and absorption (colour), wire-properties like electronic communication, and many more (see projects below).




Projects in this area span synthesis, analytical chemistry (NMR, optical spectroscopies), and computational chemistry – you can do whichever bits interest you most, or a bit of everything!

#### (b) Rational design of magneto-optic materials

All transparent materials exhibit an effect called *magneto-optic rotation*, or *Faraday rotation*. This effect is quite important: it's used in photonic devices to control the propagation of light on very fast timescales, and could be used in next-generation magnetic-field sensors. Such materials would be flexible and operative at room temperature: a far cry from the liquid-helium cooled (SQUID) detectors used currently.

Despite the Faraday effect's ubiquity, it's actually quite weak in most materials, except some ferrimagnetic garnet materials – or that was the prevailing wisdom. Recently it's been discovered that a range of organic materials, from polymers through to liquid crystals, exhibit extreme Faraday rotation.<sup>2</sup> So what? Well, the next step from this initial





discovery is to learn *how molecular structure controls* the Faraday rotation. With that knowledge, we will be able to logically design new materials with possible applications in healthcare, self-driving vehicles, and photonics/spintronics.



This project can be attacked in several directions: more synthetic or more supramolecular. You will have the opportunity to make new materials and measure their properties, either directly or in collaboration.

### (c) Molecules and assemblies for photon upconversion (with Prof. T. Schmidt)


The process of photon upconversion permits the conversion of low energy (red/near-infrared) light into higher energy light in the visible range. This process is important for two main applications: (1) enabling light-harvesting by photovoltaics across a wider spectral range; (2) powering photochemistry with low energy light, such as for in-vivo applications.

Photon upconversion requires the complex interplay of several different chromophores and their excited states. The relative arrangement of these chromophores in space, as well as their identities, is key for successful upconversion.<sup>4</sup>

The project will involve synthesising a series of organic and inorganic chromophores to systematically explore structure-property relationships. There is an opportunity to use computational chemistry to predict molecular properties, and to measure your new materials in collaboration with the Schmidt group.





Picture from Schulze, Lips, Schmidt. *Proc SPIE*, 10.1117/2.1201403.005390



### (d) Metal-to-metal communication through cross-conjugated frameworks (with Prof Les Field)

Quinones are a class of organic compounds which have a rich redox-chemistry, and which are heavily used as oxidizing agents both by chemists and in biology.

Metallaquinones are analogues of quinones where one or both of the oxygen atoms are replaced by metals. This project involves synthesising new bi-metallic or polymetallic quinonoid compounds and examining the redox chemistry and metal-to-metal electronic communication in this unusual class of molecules. The results provide fundamental insight into the nature of electronic communication and could underpin the design of the next generation of advanced materials.



### (e) Other projects

There are lots of other possible projects not listed here. If you're interested in our general area of research, or have your own ideas, please get in touch with Martin to discuss!

1. P. Wang *et al.* *JACS* **2018**, 6501; P. Wang *et al.* *JACS*, **2018**, 10881; 2. M. D. Peeks, T. D. W. Claridge, H. L. Anderson *Nature* **2017**, 541, 200; M. D. Peeks *et al.* *J. Phys. Chem. Lett.* **2019**, 2017; N. Toriumi *et al.* *JACS* **2015**, 82; 4. V. Gray *et al.* *Coord. Chem. Rev.* **2018**, 362, 54.



## DR NICOLE J RIJS

Level 7, Hilmer Building (E10) (Office) / Level 6, SEB (E8) (Lab)

T: 9385 57876 E: [n.rijs@unsw.edu.au](mailto:n.rijs@unsw.edu.au)

### STRUCTURAL CHEMISTRY WITH MASS SPECTROMETRY & ION-MOBILITY

Can you envisage a world where building a new catalyst or an artificial enzyme is like following an architectural plan for building a house? This is difficult as we don't even fully understand the construction materials! We research the properties of molecular building-blocks and their "constructed" aggregates, towards drawing up these type of blueprints.

We use high resolution ion-mobility spectrometry and mass spectrometry, computational chemistry, combinatorial libraries and robotics; along with wet chemistry to understand structure and function. We are targeting many types of chemistry involving metal ions. The way we are currently using these techniques is unique.

Electrospray ionization-mass spectrometry (ESI-MS) is rapid, sensitive, precise and well-controlled. Ion mobility (IMS) separates **much quicker** than chromatographic techniques, in milliseconds rather than minutes or hours. It is ideal for measuring the size and shape of molecules and complexes.<sup>[1]</sup> It also seamlessly interfaces with mass spectrometry. We use these methods together to monitor target reactions both simple and complex reactions with ease.

We are interested in structure, structure-function relationships, mechanism, theory underpinning chemical reactions and catalysis, chemical data crunching, and methodological development for mass spectrometry and ion-mobility. We are particularly interested in the way metals, ligands and organic molecules aggregate or react.

A An ion mobility spectrometer separates isomers in the gas phase by their mobility in a drift region	B Our liquid handling robot, uses customised components & python script
<p><math>K \propto v</math> <math>K \propto 1/t_D</math> where <math>K</math> = mobility <math>v</math> = velocity <math>t_D</math> = drift time</p>	

**During honours, researchers can become experts at:**

- different types of electrospray ionisation mass spectrometry & ion-mobility techniques (A)
- robotic preparation and analysis of solutions (B),
- screening of chemical data sets and analysis,
- electronic methods of computation, such as DFT, for structure and function
- collaborative projects with synthetic research groups


*No prior experience is necessary, these skills are in demand in industry. Projects are tailored to interests.*

It would be great to work with students on the following projects:

### Catching Structures within Dynamic Combinatorial Libraries


Dynamic combinatorial libraries (DCL) are self-assembling components in equilibrium (as in **C**). Depending on the ligand elbow different shaped oligomers are possible. DCL allow a high throughput screening for useful shaped molecules. Many can lead to useful supermolecules.

In this project, robotically generated DCLs using diketonate ligands will be monitored for the stoichiometry & shape of the evolving molecular assemblies. We will push the reactions in different ways, using additives and other changes to solution conditions.



### Assemblies of Relevance to Agriculture and Human Protection

#### D Non-protein amino acids & structural analogues



Non-protein amino acids and their analogues are significant in our environment, from ubiquitous herbicides like Glyphosate, to chemical warfare agents, to toxins produced by cyanobacteria. Recently  $\beta$ -(N-Methylamino)-L-alanine (BMAA) has been implicated in ALS and Alzheimer's disease and is not uncommonly found in waterways in regional NSW due to algal blooms.


In this project a combinatorial approach based on robotics will be used to screen the metal complexes of BMAA and its structural analogue 2,4-diaminobutyric acid (2,4-DAB) (**D**) to discover how they behave dynamically and interact with metal ions in solution.

#### Encapsulation Exquisitely Probed by Ion-Mobility

Cryptophanes (**E**) are known for their extraordinary complexation properties in water. They can capture molecules, such as methane or metal cations. They are targets for gas sensing, environmental remediation of thallium in water, and delivery of agents for MRI contrast.

In this project, ion mobility mass spectrometry will be used to study the encapsulation properties of diverse cryptophane complexes, to help explain the origins of their complexation.

#### E Example cryptophanes



### Clusters as Model Systems for Enzymatic Sites

Our ability to control chemical reactions is determined by the ability to observe them. Enzymatic reactions are notoriously difficult to observe. In this project, metals-ligand assemblies of urea, guanidine, formamide, & nucleobases will be analysed as well-defined models for enzymatic sites.

### Antibiotics and the Significance of their Metal Complexes

Metal complexes of antibiotics are important, e.g. they may inactivate their mode of action, or be critical to steps in breaking down bacterial biofilms. In this project, the structures of metal complexes of antibiotics of global importance will be analysed to better understand their role.

[1] O. H. Lloyd Williams and N. J. Rijs, *Front. Chem.*, 2021, 9 [DOI:10.3389/fchem.2021.682743](https://doi.org/10.3389/fchem.2021.682743).



## DR. FELIX RIZZUTO

Level 7, Hilmer Building (E10), 739

E: [f.rizzuto@unsw.edu.au](mailto:f.rizzuto@unsw.edu.au)

W: [rizzutogroup.org](http://rizzutogroup.org)


### DNA NANOTECHNOLOGY AND ASSEMBLY

Our research looks at new ways of synthesising and manipulating soft materials built from DNA and RNA. We take nucleic acids out of their biological context and use them to construct 'Lego' building blocks and nanorobots for sensing, autonomous devices, and replacement tissues. Our group is interested in fundamental and applied chemistry, and how we can harness chemical systems to mimic life-like processes, like self-healing, stimuli-responsiveness and, ultimately, evolution. Feel free to email me if you have any questions!

**It would be great to work with Honours students on the following projects:**


#### (a) Hierarchical DNA origami assembly

Just like you fold paper into complex shapes like swans and frogs, we can fold DNA into arbitrary geometries, like the bricks shown on the right below. We do this by adding several hundred 'staples' to a large 'scaffold' strand. We are interested in new methods to control this process, and ways in which we can bring these building blocks together to make nanomaterials and DNA polymers. The right is a TEM image where we have built long strings of these structures simply by tuning the sequence of the DNA. We are developing more 'tricks' to connect these bricks together, and to start programming the formation of nanomaterials using only single strands of DNA.



#### (b) Making 'perfect' nanomaterials


No material is perfect – even the most robust can have defects, cracks, and scars at the nanoscale. We can tune the energy landscape of materials assembly by 'pushing' these structures away from their equilibrium positions. As these systems relax to their energy minima, components are released slowly, healing defects and producing more structured, higher-performance nanomaterials. In this project we will develop new pathways for using chemical fuels and light-activated switches that rip nanomaterials apart. Instead of simply heating up and cooling down, we will develop new methods using chemical energy to direct the formation of functional nanomaterials. A sample of what we can currently do is shown to the right – using



this technology we can modulate the properties of biopolymers – including DNA and peptides – converting highly interwoven materials (before) into nanocable superstructures (right).

### (c) Light-activated and dissipative systems


Light is a powerful stimulus for self-assembly – it can be applied with high spatial and temporal control and does not generate waste, making it highly sustainable. We are interested in using light to control the self-assembly processes of DNA. Recently, we showed that double-stranded DNA could be ripped apart and folded into useful secondary structures using light-activated chemistry (see right). We have current



projects available that explore this process more thoroughly, specifically for catalysis, kinetic modulation, and templated synthesis. This project is in continued collaboration with A/Prof Jon Beves.

### (d) Building self-assembled polymers with DNA

Double stranded DNA is an archetype of programmability: the base pairs in DNA mean that we can construct two- and three-dimensional architectures relatively simply. But the range of geometries such structures can take is dictated by the inherent double helix of DNA, limiting the structural and functional diversity of DNA nanomaterials. This project will use small molecules and metal ions to reprogram how DNA self-assembles, producing new structural motifs for nanotechnology applications. We will explore a range of small molecules capable of hydrogen-bonding to common DNA bases and use these structures to build 2 and 3 dimensional constructs that we can image using state-of-the-art microscopy techniques.



Students in my group will also have the opportunity to collaborate with labs in England and Canada, as well as other labs in Chemistry and Medicine here at UNSW. All our projects are highly interdisciplinary, spanning chemistry, nanotechnology, biochemistry, medicine, and bioengineering. If anything here sparks interest, do get in touch – we have lots more going on than what I've shown here!

**Skills learnt in my group:** Biomaterials analysis, non-covalent chemistry, polymer chemistry, self-assembly, stimuli-responsive nanomaterials, microscopy

**Read more about our stuff:** *JACS*, 2023, 145, 2088; *Nat. Chem.*, 2021, 13, 843; *JACS*, 2022, 144, 12272-12279; *Nat. Mater.*, 2020, 19, 1012-1018; *Chem. Soc. Rev.*, 2020, 49, 4220-4233; *Nat. Rev. Chem.*, 2019, 3, 204-222.



## PROF. TIMOTHY SCHMIDT

Level 6, Science and Engineering Building (E8)

T: 04 39 386 109 E: [schmidtim@gmail.com](mailto:schmidtim@gmail.com)


### EXCITED MOLECULES

The Schmidt group undertakes research on electronically excited states of molecules. We undertake experiments on molecules in the condensed phase, with applications to renewable energy research, and in the gas phase, with applications in astronomy and astrophysics. We use laser systems with high temporal resolution ( $10^{-13}$  s) to watch energy flow in molecule systems, and high spectral resolution ( $0.07$   $\text{cm}^{-1}$ ) to compare with astronomical observations. We underpin all of our experimental results with appropriate theory and modelling. We have many projects on offer, so come and have a chat!

It would be great to work with Honours students on the following projects:

#### (a) Photochemical Upconversion


What do solar cells, organic light emitting diodes (OLEDs), and hydrogen production have in common? They can all be made more efficient by converting lower-energy excited states into high energy excitations through a process known as photochemical upconversion. By using chromophores to make red light into blue light (or infrared into yellow *etc.*), we can harness more of the solar spectrum, convert dark states in OLEDs into bright states, and increase solar hydrogen production efficiency. To achieve these goals, efficient solid-state upconverters are required.



Our group recently showed that by attaching chromophores to a porous scaffold we can achieve efficient solid-state sensitization of upconversion:

(<https://pubs.acs.org/doi/full/10.1021/acsenergylett.3c01678>).

In this project we will expand on this work to include new wavelength regions that will benefit solar cells, LEDs, and the photocatalytic production of hydrogen.




**What you will do:** Organic synthesis, laser spectroscopy, device fabrication, modelling (pick and mix)

#### (b) Singlet Fission (with Dr Peeks and/or A/Prof. Beves)

UNSW has led the development of silicon photovoltaics (Si-PV). However, Si-PV are fast approaching their maximum theoretical efficiency of 29%. To realise greater efficiencies in Si-PV the fundamental loss mechanism of thermalisation, which accounts for 33% of total losses, needs to be addressed. Thermalisation losses arise because a Si-PV can only use about 1.1 eV of an absorbed photon, no matter its energy. It would be nice if we could split the energy of high energy ( $>2.2$  eV) photons in two!

Singlet fission (SF) is such a process. By incorporating a SF capable chromophore with Si-PVs the theoretical efficiency of 29% can be surpassed.



Through the \$6M ARENA-funded OMEGA Silicon project (<https://www.omegasilicon.solar/>) we are developing the next generation of solar cells based on SF. There are multiple potential projects involving SF. If you are interested, please get in contact for more information!


**What you will do:** Organic synthesis, laser spectroscopy, device fabrication, modelling (pick and mix)

### (c) Interstellar Molecules (with Dr Chris Hansen)



The *Molecular Photonics Laboratories* house sophisticated lasers and equipment with which we can discover new transient chemical species of importance in the gas phase chemistries of our atmosphere and the interstellar medium. As stars die, they eject complex organic molecules into the interstellar medium, where they live out millennia before being incorporated into new stars and planetary systems. These organic molecules are

the seeds of life, but, as yet, we do not know the chemical make-up of the interstellar medium from which planetary systems are formed. Using a star as a lamp, we can peer into this medium using telescopes by observing molecular absorption spectra. However, despite there being hundreds of nibbles taken out of the visible stellar spectra of stars occluded by diffuse clouds, only a few molecules have been unambiguously detected by their visible spectra. The unidentified features are known as the *diffuse interstellar bands* (DIBs), and are the longest standing mystery in astrophysical spectroscopy. In this project, we will develop techniques to capture the spectra of isolated, never-seen-before aromatic cations and radicals (which are the leading candidates for carrying the DIBs), and (hopefully) solve this long-standing problem.



**What you will do:** Laser spectroscopy, electric discharges, vacuum techniques, quantum chemistry



## A/PROF. NEERAJ SHARMA

Room 129, Dalton Building (F12)

T: 9385 4714 E: [neeraj.sharma@unsw.edu.au](mailto:neeraj.sharma@unsw.edu.au)


### BATTERIES, SOLID STATE AND MATERIALS CHEMISTRY

- We chemically tune the atomic arrangement (crystal structure) of solid state materials to enhance their physical properties such as energy storage capacity, ionic conductivity or thermal expansion.
- We use a combination of techniques to characterise our materials, including but not limited to X-ray and neutron diffraction (at the Australian Synchrotron and ANSTO), solid state NMR, electrochemical and impedance analysis, and electron microscopy.
- Our goal is to fully characterise materials, place them into real-world devices such as batteries and solid oxide fuel cells, and then characterise how they work in these devices.

It would be great to work with Honours students on the following projects:

#### (a) Towards the next generation of batteries: Sodium-ion batteries

Lithium-ion batteries are ubiquitous in our daily lives, e.g. mobile phones and laptop computers, but their limitations have restricted wide-scale use in applications requiring higher power, e.g. electric vehicles and energy storage of renewable energy. This project will target new battery chemistries, in particular sodium-ion batteries, by developing and characterising new electrode and electrolyte materials. We will work to develop a reliable and affordable room-temperature sodium-ion battery to provide sufficient power for large-scale energy storage from intermittent renewable power sources. Students will work on one of the following parts of a battery and test their component in idealized batteries.



#### *Positive electrode materials*

These electrodes provide the source of the sodium-ions and represent the largest cost and energy limitations for lithium-ion batteries. Here, new sodium-containing transition metal oxides, phosphates or sulfates will be synthesized and characterized to determine the relationship between crystal structure and battery performance. **We are working towards scaffolding layered electrode materials in order to dramatically improve performance.**

#### *Electrolytes*

Sodium-ion conducting ceramics or glassy-ceramics are known to be excellent electrolytes at high temperatures (>300°C). This project works towards making materials with sufficient sodium-ion conduction at room temperature.




### Negative electrode materials

Negative electrodes are the least investigated component in a sodium-ion battery and the compounds used for lithium-ion batteries show poor performance in sodium-ion batteries. By developing new negative electrodes and understanding their limitations towards reversible sodium insertion/extraction we will be able to enable the next generation of devices.

#### (b) Tuning negative thermal expansion to produce zero thermal expansion materials

The majority of materials expand during heating *via* thermal expansion and this process is responsible for billions of dollars per year in maintenance, re-manufacture and replacement costs due to wear and tear on both moving parts (*e.g.* in aircraft gas turbines), and components that are designed to be static (*e.g.* in optics, coatings, electronics). If a zero thermal expansion (ZTE) material can be made, a material that neither expands nor contracts upon heating, this could dramatically reduce industrial costs. In order to achieve this, the opposite extreme of materials are considered in this project - negative thermal expansion (NTE) is a property exhibited by a small group of materials predominantly due to transverse vibrations of atom groups or cooperative rotations of units (*e.g.*  $-\text{CN}-$  or  $\text{WO}_6$ ). These materials typically feature large crystallographic voids and cations with variable oxidation states. So why not use a battery as a synthesis tool? In this project we will controllably insert Li and Na into the voids of the NTE materials, via a battery, in order to tune the cooperative rotations to produce ZTE materials.



#### (c) Improving solid-state electrolytes by understanding their formation characteristics and phase evolution

Safety is an important aspect of high power batteries. Using a solid-state electrolyte has significant advantages to the highly flammable liquid electrolytes that are commercially available. Unfortunately the ionic conductivities of solid-state compounds are generally lower than the liquid counterparts, especially under ambient conditions. At the other extreme, solid oxide fuel cells often operate at approximately  $1000^\circ\text{C}$  as the operating temperatures are essentially determined by the ionic conductivity of the electrolyte. In both examples, electrolyte ionic conductivity is a critical hurdle in preventing further development and use of these technologies. The ionic conductivity is directly related to the crystal structures adopted by the electrolytes and how they evolve with temperature. In this project lithium-ion and oxide-ion conducting materials will be synthesized and their ionic conductivities characterized. Importantly, variable temperature time-resolved neutron powder diffraction will be used to study the formation (from starting reagents) of these ionic conductors under varied conditions. This will shed light on the formation processes and optimal conditions required for synthesis.

#### (d) Other projects

Depending on your interests, other solid state projects, *e.g.* making new superconductors, can be designed. Please consult with Neeraj for further details.



## SCIENTIA PROF. MARTINA STENZEL

Level 7, Science and Engineering Building (E8)

T: 9385 4656 E: [m.stenzel@unsw.edu.au](mailto:m.stenzel@unsw.edu.au)


### NANOPARTICLES AND TAILORED POLYMERS FOR CANCER TREATMENT

- The delivery of drugs can be improved by packaging the drug into nanoparticles. Nanoparticles for drug delivery have typically sizes below 100 nm and can be prepared using various materials including polymers. In our group, we synthesize various polymers to create core-shell nanoparticles – the core holds the drug, mainly anti-cancer drugs, while the shell makes the particles soluble and determines the interaction with cells.
- In our group, we work on different aspects starting from organic synthesis to polymer nanoparticle preparation to testing these particles on cancer cell lines. We have meeting with clinicians and we discuss their drug delivery problems.

It would be great to work with Honours students on the following projects:

#### (a) Making really small polymer nanoparticles

Most nanoparticles described in literature used for drug delivery have sizes above 20 nm. There are however various reasons to make nanoparticles that are even smaller as they seemed to have better circulation time in the blood stream and can diffuse into tumour tissue. To achieve this, we are preparing polymers with attached drugs of various amount and sequence. These polymers are then collapsed into single chain nanoparticles, which are typically below 5 nm in size.




*Collapsing of polymer chains into single chain nanoparticles*

The unusual part about this project is that we will use flow chemistry to assemble the nanoparticles

#### (b) Drug carriers inspired by nature: Nanoparticles with sugar antennae


Carbohydrates are involved in a number of biological communication events as they carry sugar-specific receptors. This specific sugar-receptor interaction can be used to deliver nanoparticles specifically to receptor-expressing cells, which can result in improved biodistribution. Synthetic polysaccharides, coined glycopolymers, have been shown to be superior to single sugars as they can bind simultaneously to several receptors. In this project we would like to develop glycopolymers for the delivery of drugs to brain cancer. We will entrap anti-cancer drugs that are specifically developed in the school of medicine against brain cancer. If time permits, the drug carrier will be tested on brain-cancer cell lines.



*Synthesis of glycopolymers and their self-assembly into nanoparticles*

**c) Delivering siRNA to combat viruses** Collaboration with Prof Bill Rawlinson (Medicine)

Safe and efficient transfection of nucleic acid-based drugs (like oligonucleotide and short interfering RNA (siRNA)) to desired site, with the assistance of nanoparticles, has been of considerable interest in the nanomedicine field in the past few decades. The Pfizer mRNA vaccine has shown how nanoparticles can benefit the delivery of these otherwise fragile drugs. In this project, we will design nanocarriers based on cationic polymers to deliver siRNA against Human cytomegalovirus (CMV) infection




**Tissue-specific nanoparticle vehicles for targeted CMV antiviral delivery.**

**A)** CMV antivirals will be encapsulated within polymer nanoparticles decorated with placental specific antibodies for targeted placental delivery. **B)** Preliminary data using untargeted nanoparticles coupled with siRNA UL54 show CMV inhibition in placental cell cultures (\* $P < 0.05$ ).

**(d) Where is the drug? – Fluorescence studies to monitor drug release**

Key to the success of drug delivery with nanoparticles is the entrapment of the drug inside the nanoparticle, only to be released when the nanoparticle is inside the targeted cell. One way of monitoring the release is to monitor the fluorescence of the drug carrier and that of the drug. If the drug is entrapped inside the carrier, Förster resonance energy transfer (FRET) occurs. If the drug is released from the nanoparticle, fluorescence changes. FRET can be used to monitor the drug release in live cells, which can help us make the decision how to improve our drug delivery system.





## A/PROF. JOHN STRIDE

Level 1, Dalton Building (F12)  
T: 9385 4672 E: [j.stride@unsw.edu.au](mailto:j.stride@unsw.edu.au)

### MATERIALS CHEMISTRY AT THE NANOSCALE

My group focuses on making and understanding new materials that are often focused on some of the major challenges facing us today: energy, water and sustainability. We make use of a range of techniques that include X-ray and neutron scattering in truly multi-disciplinary projects. Key to these studies is the notion of hierarchical emergent properties and complexity - the world around us derives from simple inter-molecular interactions; we aim for a greater understanding of these fundamental processes in order to deliver new materials displaying novel properties.


**It would be great to work with Honours students on the following projects:**

**(a) Metal organic frameworks (MOFs): coordination chemistry of the 21<sup>st</sup> century**

Over the last 20 years, inorganic chemistry has taken on board a number of new concepts and approaches that have reinvigorated the subject – one area showing particular promise is polymeric coordination compounds or MOFs. These topologically beautiful materials display intimate long range ordering and immense compositional flexibility, along with structural rigidity; they are ideal hosts for a range of molecular guests, opening up many potential applications.


*Sorting and storing molecules - how to select for one molecule over another*

This research project is specifically targeted at very real challenges faced in industry - effective separations of mixed gas streams and facile storage of gaseous fuels such as H<sub>2</sub>. Highly porous MOFs make excellent host materials for small molecules such as CH<sub>4</sub> or H<sub>2</sub>. By tuning their properties MOFs can become efficient storage vessels or effective gas-selective membranes such as the H<sub>2</sub> selecting MOF shown here.




*Quantum phenomena in magnetic materials*

Magnetic materials have revolutionised the way in which we store and use information and have a key role to play in quantum computing; they have also been a navigational aid for centuries and are even pretty useful at securing notes to the fridge door. It is fascinating therefore that we still do not fully understand the behaviour of such materials, especially when dimensionality is constrained. MOFs can have single chains (1D) or sheets (2D) of metal ions embedded into a non-magnetic matrix, making them ideal materials in which to study the effects of magnetic quantum confinement.



## (b) Order and disorder in molecular materials

Solid state materials are often thought of in terms of the long range ordering of motifs into lattice structures; however what occurs upon phase transitions when molecular ordering may change or even order gives way to disorder? Welcome to the world of phase transitions, in which entropy and enthalpy play important roles in determining the behaviour of molecular motifs. Planar molecules, such as small aromatics, are of particular interest in that approximating to oblate discs, their reduced dimensionality directly influences their intermolecular interactions and orientations. They are also ideal systems to study; not too big, amenable to computational simulations, ubiquitous and very stable.




### *Inter-molecular hydrogen-bonding*

Identified by Linus Pauling around 80 years ago, the hydrogen bond is the champion of intermolecular interactions, the basis of biology and our watery world. However there is a lot to still learn and to problems to study when it comes to H-bonding - we have been looking at a number of model H-bonded systems, making use of solid state NMR, X-ray and neutron diffraction and inelastic neutron scattering. This work is highly collaborative, requiring high-end research infrastructure and sophisticated numerical modelling - it is ideally to students with an inquisitive mind, seeking deep insights into the fundamentals of our every day life.

### *Donor-Acceptor stacks: heterojunction photovoltaics to molecular magnets*

The intermolecular interactions between efficient electron donors (D) and acceptors (A) yield optically active charge transfer materials that can act as organic semiconductors, photovoltaics, ferroelectrics and light emitting diodes. Complete electron transfers can result in bulk magnetic materials. We aim to investigate the interactions of simple D...A stacks whilst modifying the peripheral functional groups, known to contribute to molecular packing. In this way, self-healing semi-conducting liquid crystalline materials can be produced that show remarkable anisotropy, enabling uniaxial conduction under greater load. With the wide range of suitable D and A molecules available, these materials have tremendous promise in their capacity to be tuned for specific applications, whether it be for emission in the visible spectrum (OLEDs) or broad-range absorption (OPVs). Being relatively small molecules, they are also suited to computational studies that are highly informative in terms of the electronic interactions and  $\pi$ - $\pi$  stacking interactions.



## (c) Other projects

Other projects involving materials-based chemistry, nanotechnology, graphene, crystallography and spectroscopy are available and can be tailored to your interests. Feel free to come and discuss possible research projects.



## DR. SCOTT A. SULWAY

Room 113, Level 1, Dalton Building (F12)

T: 9385 5236 E: [s.sulway@unsw.edu.au](mailto:s.sulway@unsw.edu.au)

### SYNTHETIC INORGANIC CHEMISTRY – LANTHANIDE COORDINATION COMPLEXES

Lanthanides are a commonly overlooked area of coordination chemistry – people often say “*But we know everything there is to know and how they react*”... This isn't so, lanthanide complexes are incredibly interesting and have a range of potential applications. Lanthanides have uses in catalytic cycles, luminescent devices & interesting magnetic properties that could be utilised in data storage devices or qubits in quantum computing.<sup>1</sup> This is where the research in the Sulway group comes in, we are exploring the synthesis and characterisation of new lanthanide containing coordination compounds that could be used in the technology of the future.



Skills you will learn:

- Manipulation of air- and moisture-sensitive compounds
- Organic and Inorganic synthetic chemistry
- Structure elucidation – NMR spectroscopy ( $^1\text{H}$ ,  $^{13}\text{C}$ ), IR spectroscopy, SQUID magnetometry and XRD (Yeap, we grow crystals)!

It would be great to work with Honours students on the following projects:


#### (a) Scorpionate Ligands in Lanthanide SIMs

Ongoing research into Single-Ion Magnets (SIMs) is of great interest due to their potential applications in molecular storage devices and as Qbits in quantum computing. SIMs retain their orientation of magnetisation in the absence of a magnetic field due to having a barrier to said reorientation. Record SIM barriers utilise cyclopentadienyl ligands, these have several drawbacks and as such in the Sulway group one aspect that we explore are isolobal systems such as scorpionate ligands.<sup>2</sup>




### (b) Exploring Alternate Cyclopentadienyl Ligands in SIMs

As described in project (a) the current record barriers in SIMs utilise the cyclopentadienyl moiety, we also look into these systems focusing upon bulkier ligands. These systems are great if you want to combine some inorganic and organic synthesis.



### (c) Equatorial Ligands in Prolate Lanthanide Ions

How about something a little different? SIM research often focusses on axial systems, equatorial systems are very rarely studied and not very well understood. We work on trying to expand the understand in this emerging area by looking into the “neglected” lanthanide ions.



### (d) Have your own ideas?

I'm open to discussing other potential ideas that you have after all it is your Honours year you should work on something you are interested in, just send me an e-mail...

1. (a) Goodwin, C. A. P., *Dalton Trans.*, 2020, 49, 14320-37, (b) Woodruff, D. N.; Wimpenny, R. E. P.; Layfield, R. A., *Chem. Rev.*, 2013, 113, 5110-48, (c) Rinehart, J. D.; Long, J. R., 2011, 2, 2078-85.  
2. Thomas, J. R.; Sulway, S. A., *RSC Adv.*, 2021, 11, 16158-60.



## PROF. PALLI (PALLI) THORDARSON

Level 7 (room 741), Hilmer Building (E10)

T: 0417 459 348 E: [p.thordarson@unsw.edu.au](mailto:p.thordarson@unsw.edu.au)

### RNA CHEMISTRY, ORIGIN OF LIFE AND NANOMEDICINE

- **RNA Chemistry** with focus on understanding how RNA interacts with peptides other molecules and how these interactions can be applied in RNA science and therapeutics
- **Origin of Life and Systems Chemistry**, exploring the role of self-assembly in how life originated and how we can make life-like systems.
- **Development** of 3D Cell Culture materials for use in **catalysis** and **medical research**
- **Synthesis** of novel **peptides** for **nanomedicine**, including drug delivery and tissue engineering

It would be great to work with Honours students on the following projects:


(a) **Peptide-RNA interactions – solving pressing problems in prebiotic chemistry and medicine** (*Potential for collaborations with Dr Albert Fahrenbach & Dr Anna Wang School of Chemistry, Prof. Martin Van Kranendonk, BEES & A/Prof. Archa Fox, University of Western Australia*).

Peptides/proteins and RNA are two of the key building blocks of life. Recently it has become proteins and RNA drive the formation of lava lamp or vinaigrette “droplets”<sup>1</sup> within the cell but biologists are now just uncovering how important droplet- or gel-like protein-RNA complex are in biology and medicine. At the same time, Origin of Life research<sup>2</sup> has started to turn its attention to a new hypothesis for how complexity could have arisen from a the “pre-biotic soup” of chemicals, particularly peptides and short RNA's.<sup>3</sup> We aim to solve key problems on both fronts by synthesising short RNA and peptides and investigate the structures they form. This would then give clues towards how we could develop medical treatment that modulate these interactions and how we could address one of the most important questions in science, *i.e.*,

**how did life originate**. If you join our team to work on these challenges, you would not help us tackling these problems but you will also gain valuable experience in synthesis, self-assembly and the chemistry of RNA and peptide biomolecules such as the peptide shown here:




Did the cell start of as a collection of peptide-RNA “droplets”? And is this how the cell is really organised? (from E. Dolgin, Nature 2018, 555, 300.






**(b) Novel switchable and hybrid peptide-based materials for catalysis and 3D cell cultures** (Potential for collaborations A/Prof. Jonathon Beves, A/Prof. T. Vinh Nguyen and A/Prof. Kris Kilian, School of Chemistry).

Self-assembled peptide gels have already been proven to be useful as 3D material for growing living cells, even neurons.<sup>4</sup> We have extended this work to include the formation of gels that can be changed through a photo-switch, <sup>5</sup> or mixing with a biological material such as collagen. In more recent work we also been able to demonstrate that self-assembled gels can be used as a scaffold for catalysis in chemical synthesis. Projects involving developing novel photo-switchable and hybrid gels applications in cell biology and catalysis are available for those with interest in medicinal chemistry, nanomedicine, supramolecular and synthetic chemistry.



**(c) Novel RNA therapeutics** (Potential for collaborations with Dr Chantelle Ahlenstiel, Kirby Institute UNSW, A/Prof. Joshua McCarroll and Prof. Maria Kavallaris, Children's Cancer Institute Australia).

Ribonucleic acid (RNA) is now recognised to play a much more important role in biology than previously thought. It is not merely a “messenger” (mRNA) but has many other functions in the cell. From a chemists point of view the structural diversity of these different types of RNA molecules is fascinating (see Figure from a recent review by the Disney group,<sup>6</sup>). Many of these RNA molecules play a crucial role in diseases. Inhibiting or regulating RNA function through the application of small molecules, peptides or specially designed RNA molecules such as small interfering RNA (siRNA) therefore represents a powerful strategy to develop new and better therapeutics for cancer, infectious diseases (including HIV and COVID-19) and various disorders that have genetic origin or relate to gene expression misregulation. Projects, including with our collaborators, involve developing novel peptide binders for disease-causing RNA motifs, synthesis of novel siRNA molecules and siRNA-ligand conjugates are available for anyone that wants to combine synthetic chemistry, supramolecular chemistry and medicinal chemistry in their research training.



1. Elie Dolgin, Cell biology's new phase, *Nature*, **2018**, 555, 300-302.
2. Pall Thordarson, "Emergence of Life" in *Encyclopedia of Supramolecular Chemistry*; eds: Jerry L. Atwood, Jonathan W. Steed, Marcel Dekker Inc., New York, **2004**, 528-534.
3. Martin Van Kranendonk, David W. Deamer and Tara Djokic, Life Springs, *Scientific American*, August **2017**, 28-35.
4. Adam D. Martin, Sook Wern Chua, Carol G. Au, Holly Stefen, Magdalena Przybyla, Yijun Lin, Josefina Bertz, Pall Thordarson, Thomas Fath, Yazdi D. Ke and Lars M. Ittner, Peptide nanofiber substrates for long-term culturing of primary neurons, *ACS Applied Materials & Interfaces*, **2018**, 10, 25217-25134.
5. Fayaz Ali Larik, Lucy L. Fillbrook, Sandra S. Nurtilla, Adam D. Martin, Rhiannon P. Kuchel, Karrar Al Taief, Mohan Bhadbhade, Jonathon E. Beves\* and Pall Thordarson\*, Ultra-Low Molecular Weight Photoswitchable Hydrogels, *Angewandte Chemie International Edition*, **2021**, 60, 6764-6770.
6. S. M. Meyer, C. C. Williams, Y. Akahori, T. Tanaka, H. Aikawa, Y. Tong, J. L. Childs-Disney, M. D. Disney, Small molecule recognition of disease-relevant RNA structure, *Chem. Soc. Rev.*, **2020**, 49, 71767-7199.



## PROF. RICHARD TILLEY

B89, Chemical Sciences Building (F10)

T: 9385 4435 E: [r.tilley@unsw.edu.au](mailto:r.tilley@unsw.edu.au)

### NANOPARTICLE SYNTHESIS & ELECTRON MICROSCOPY

Our group is world leading in the synthesis of the highest performing nanoparticle catalysts and medical imaging agents. Our synthesis expertise allows us to engineer complex nanoparticle catalysts that with atomic level precision. As Director of the Electron Microscope Unit you will use state-of-the-art electron microscopes that are the best in Australia to characterise cutting edge nanoparticles.

#### Magnetic nanoparticles for cancer detection using Magnetic Particle Imaging

As the first to have a Magnetic Particle Imaging (MPI) instrument in Australia, we are in a unique position to detect early stage tumours and cancerous cells with the most sensitive and precise imaging. The exceptional magnetic properties of iron and iron oxide nanoparticles make these ideal candidates for this state-of-the-art application. These key magnetic properties are dictated by the size, crystallinity and composition of the magnetic nanoparticles.






Figure 1: MRI images from iron nanoparticles injected into a mouse to enhance the contrast of a tumour.

Using the leading edge of solution phase synthesis, precise control over the nanoparticles and their magnetic properties can be achieved (Figure 2). In this project, well-defined nanoparticles with controlled crystalline domains will be studied for MPI. You will use transmission electron microscopy and collaborate with leading researchers in MPI from Australia. Overall, this work will tune nanoparticle size with precise synthetic control to optimise magnetic properties of iron and iron oxide nanoparticles for MPI.

Figure 2: Transmission electron microscopy images of iron nanocubes and their magnetic properties for use in MPI.<sup>1</sup>



1. Gloag, L. *et al.* Zero valent iron core–iron oxide shell nanoparticles as small magnetic particle imaging tracers. *Chem. Commun.* **56**, 3504–3507 (2020).

#### Controlling nanoparticle structure for active and stable catalysts in renewable energy storage

The oxygen evolution reaction (OER) is crucial for the storage and conversion of H<sub>2</sub> fuel and requires highly active and highly stable catalysts to drive it. Our expertise in nanoparticle synthesis has allowed us to create the most active and stable nanocatalysts for OER reported to date.<sup>1</sup> We achieved this by

synthesizing 3D branched Ru nanoparticles with structural features that both prevent dissolution and improve oxidation catalysis (Figure 1).




Figure 1: Energy dispersive X-ray spectroscopy elemental mapping of Pd-Ru branched nanoparticles and TEM images of individual nanoparticles. Models show the controlled direction of growth of Ru from Pd seed.


In this project, Ru nanoparticles will be synthesized with low index facets which are critical for achieving stable reaction kinetics that prevent dissolution of Ru and enhance the catalytic activity. This work will combine the development of synthetic methods to control the size, shape and composition of Ru-based nanocatalysts, with advanced characterisation using high-resolution transmission electron microscope and also evaluation of their electrocatalytic performance. This allows for the relationships between nanoparticle structure and catalytic performance to be fundamentally understood and tuned to create leading nanocatalyst materials.

1. Gloag, L. *et al.* A cubic-core hexagonal-branch mechanism to synthesize bi-metallic branched and faceted Pd-Ru nanoparticles for oxygen evolution reaction electrocatalysis. *J. Am. Chem. Soc.* **140**, 12760–12764 (2018).

### Synthesising strained Pt on metal nanoparticles for enhanced electrocatalytic activity in hydrogen fuel cells

In order to convert to sustainable energy cells in a hydrogen economy, nanocatalysts need to be high-performing and use minimal amounts of scarce Pt. Strained Pt on the surface of a metal nanoparticle is a promising structure for highly active fuel cell catalysts. Depositing Pt directly onto Ni nanoparticles creates highly strained Pt that maximises the specific and minimises the amount of expensive Pt that is used to provide the highest mass activities reported to date (Figure 1).<sup>1</sup>

Figure 1: Relationship between strain and HER activity and elemental map of a Pt on Ni nanoparticle.<sup>2</sup>



In this project, nanoparticles will be decorated with small clusters of Pt atoms for use as high performance catalysts. By controlling the position of Pt atoms on different metal nanoparticle structures, both electrocatalytic activity and stability will be optimised to create the most advanced and effective nanoparticle catalysts.

1. Alinezhad, A. *et al.* Direct Growth of Highly Strained Pt Islands on Branched Ni Nanoparticles for Improved Hydrogen Evolution Reaction Activity. *J. Am. Chem. Soc.* **141**, 16202–16207 (2019)



## DR. ANNA WANG

Office: Level 7, Hilmer (E10) | Lab: Level 6, SEB (E8)

E: [anna.wang@unsw.edu.au](mailto:anna.wang@unsw.edu.au) | [annawanglab.com](http://annawanglab.com)

### SOFT MATTER BIOPHYSICS AND THE ORIGINS OF LIFE

- We tackle problems at the nexus of chemistry, physics, biology, and materials science.
- One of our main interests is in lipid bilayers (Fig 1), which encase our cells but are also used for drug delivery.
- One of our other main interests is in how matter interacts with light, which we study with holographic imaging and light scattering calculations.
- Our group has students from many backgrounds – interdisciplinary problems require multidisciplinary teams of problem solvers.
- Students typically work with biomaterials like lipids, RNA, and gels, and do microscopy, optics, image analysis, data analysis and calculations with Python.
- We have ongoing international collaborations, including with industry.

It would be great to work with students on the project topics (a) – (e):

#### (a) Building an artificial cell life cycle

Being able to create a self-perpetuating artificial cell (Fig 2) reflects an unprecedented understanding of lipid bilayers and living systems, and is an overarching goal of our group's. Outstanding questions include:

- How can we control the fusion of lipid compartments by modulating their composition?
- How do we get these compartments to grow and divide?
- How do different populations of lipid vesicles/liposomes compete with each other?
- How does crowding inside cells and artificial cells affect diffusion rates and membrane shape changes?

The answers to these questions are important for understanding how evolution could have been kickstarted at the origins of life (Fig 2). They will also reveal fundamental membrane biophysics, with implications in drug delivery processes.

Students will learn lipid manipulation techniques, lipid vesicle/liposome processing techniques, fluorescence spectrophotometry, microscopy, and biophysics assays.




Fig. 1 Liposomes containing RNA




Fig. 2 A dynamic growing system


Recent papers include [“Passive endocytosis in model protocells”](#) (PNAS, 2023), [“Lipid exchange promotes fusion of model protocells”](#) (Small Methods, 2023), [“Subtle Changes in pH Affect the Packing and Robustness of Fatty Acid Bilayers”](#) (Soft Matter 2022), [“Bulk, Self-Assembly of Giant, Unilamellar Vesicles”](#) (ACS Nano, 2020).

**(b) Characterising liquid-liquid phase separation with holographic microscopy**

Liquid-liquid phase separation (LLPS) is a supramolecular phenomenon whereby macromolecules interact and condense into one liquid phase (dispersed in another). RNA and peptides, for example, undergo LLPS in cells. LLPS also occurs in secondary organic aerosols.

LLPS is also a first step in many diseases, including Alzheimer's, because LLPS droplets are only metastable. How do such droplets transition to a disease state?

The student will develop a mechanistic understanding of LLPS, which involves the characterisation of LLPS at a single-droplet level. This project will use holographic microscopy to characterise LLPS systems, revealing how their size density evolves over time. Holographic imaging of colloidal particles trapped in LLPS droplets will reveal the viscosity of the internal droplet environment. Recent papers include "[Using holographic microscopy to measure the effect of confinement on crowding agents in lipid vesicles](#)" (ChemBioChem, 2023)



**Fig. 3 Hologram of droplet**


**(c) Designing particles with novel optical properties for printing, cosmetics, and electronics**

This project involves using light scattering software to help design particles with intriguing optical properties e.g. transparency in the visible and high scattering in the UV. These novel particles could increase the efficiency of inks, sunscreens, electronics, and more. No background in Python needed, we'll teach you what you need to know. Recent papers include "[In-line holographic microscopy with model-based analysis](#)" (Nature Reviews Methods Primers, 2022).

**(d) Measuring cellular forces for improved material design (in collaboration with A/Prof. Kris Kilian)**

The mechanical environment of cells often determines their fate. To design better tissue engineering scaffolds and materials, we must first measure how cells push and pull on their environments. We propose using holographic imaging (Fig. 4) to solve this problem.


The student will pioneer and then use holographic traction force microscopy to investigate focal adhesion and traction stress propagation from adipose derived stem cells (ADSCs) adherent to the surface. Relationships between cell generated traction and differentiation to adipocyte, chondrocyte, and osteoblast lineages will guide the design of materials for tissue engineering and regenerative medicine



**Fig. 4 "Holographic imaging" in the movies**

**(e) A project of your choosing (and imagination)**

There are many more possible projects in our group pertaining to artificial cells, origins of life, soft matter, microscopy, and more – speak to Anna to see what's possible, we collaborate with many groups in the School.



## DR. SIOBHÁN WILLS

Office: Level 1, Dalton Building (Rm 130)

E: [siobhan.wills@unsw.edu.au](mailto:siobhan.wills@unsw.edu.au)

### CHEMISTRY AND SCIENCE EDUCATION

- Chemistry is central to understanding our world. For us to pass on and build upon our collective knowledge, we must fully comprehend how everyone learns best in science subjects. Science education research (SER) helps us optimise the students learning experience and communicate effectively. I am interested in student learning processes and how those can be developed and supported. SER can encompass a wide range of themes, so my projects are many and varied from classroom activities to assessment design, educational technology, and chemistry outreach. Projects can cover both secondary and tertiary education.
- Education research is a multidisciplinary area, and we welcome students from any university background to bring new ideas and opinions.
- I am part of the wider UNSW science and chemistry education research network and have ongoing collaborations at USydney, UWA, and Curtin Uni.
- We are keen to explore all aspects of science education and learning, and this means there are endless possible projects. If you have any curious questions you want to explore, please get in touch to discuss how your research interests could grow to be an Honours project!

**It would be great to work with students on the following projects (a) – (e):**

**(a) Designing authentic chemistry assessments with effective feedback cycles for learning**

*(in collaboration with Dr Shannan Maisey, Dr Sara Kyne, and Dr Jeffrey Black)*



Assessment is a main pillar of university learning, both to support progress in learning and to measure comprehension of a topic. Recent research has shown that assessment can be very formulaic, which underprepares students to be “job-ready” at the end of their degree. In addition, feedback process can be confusing and sub-optimal.

This research area looks at improving assessment and feedback practices in chemistry.

There are several sub-projects in this area:

- Designing higher year, project-based assessments contextualised to real world problems – providing effective feedback for learning and properly testing comprehension
- Designing and evaluating feedback cycle prototypes to enhance student learning – in workshops, on lab reports, and after tests and exams.
- New approaches to assessment using educational technology.
- Exploring the diversity of first year exam questions and formats to improve clarity and effectiveness of digital and paper exams.

These projects will use qualitative methods such as interviews and focus groups to collect data on what an authentic chemistry assessment should include to help and assess student learning. Scope could include interviewing students and academics on ‘ideal’ assessments and core graduate skills, running focus workshops on building assessments and feedback protocols, and analysing examples of successful/unsuccessful assessments.

## (b) Designing and evaluating a new collection of high school extension experiments

*(In collaboration with Dr Stephen George-Williams from the University of Sydney)*



Experiments in high school can be simple and unexciting. The new 7-12 chemistry syllabus calls for more nuanced, context-based experiments. Working with high school teachers from the Royal Australian Chemistry Institute Chemistry Education group and Dr George-Williams, we will aim to generate and evaluate a selection of experiments to support the new curriculum.

This project will include lab time for experiment development, building supporting teacher and students resources, and visits to schools or hosting students for evaluative testing of the labs using validated instruments.

Final experiments will be shared with teachers across NSW.

## (c) Metacognition in learning chemistry



Experts tackle problem solving almost instinctively, easily breaking down a problem to identify approaches to a solution. Novices (such as undergraduate students) require support to build proficiency, so answering the following research questions will help them become more accomplished as such tasks.

How do students think about solving problems (metacognition)? Where are the sticking points and misconceptions? How can we aid in overcoming these barriers to build problem-solving skills?

Certain concepts in chemistry can be troublesome for many students and could be better taught. Using your own experience as well as interviews from other students and assessment analysis, you will identify challenging concepts that can then be further explore to design support interventions. Some ideas are how we interpret NMR spectra for compound identification<sup>2</sup>, how we design synthetic pathways, and how we propose mechanisms<sup>3</sup>. One current subproject in collaboration with Dr Laura McKemmish is assessing the metacognition around writing creative chemistry study questions.

## (d) Sensory perceptions in the lab

*(In collaboration with Dr Ron Haines)*




The chemistry teaching laboratory can be a sensory and cognitive overload for both students and instructors alike.<sup>4</sup> The overload may negatively affect meaningful and successful learning experiences in the lab. This project can encompass developing a sensory audit of lab spaces, or student and staff surveys to identify sensory overload perceptions and assess impact, or the development and evaluation of interventions to alleviate sensory overload.

<sup>2</sup>Connor, M. C.; Finkenstaedt-Quinn, S. A.; Shultz, G. V., Constraints on organic chemistry students' reasoning during IR and <sup>1</sup>H NMR spectral interpretation. *Chemistry Education Research and Practice* 2019, 20 (3), 522-541.

<sup>3</sup>Caspari, I.; Kranz, D.; Graulich, N., Resolving the complexity of organic chemistry students' reasoning through the lens of a mechanistic framework. *Chemistry Education Research and Practice* 2018, 19 (4), 1117-1141.

<sup>4</sup>Flaherty, A., The Chemistry Teaching Laboratory: A Sensory Overload Vortex for Students and Instructors? *J. Chem. Educ.* 2022, 99, 4, 1775-1777



## PROF. CHUAN ZHAO

Level 1, Room 127, Dalton Building (F12)

T: 9385 4645 E: [chuan.zhao@unsw.edu.au](mailto:chuan.zhao@unsw.edu.au)

## CLEAN ENERGY TECHNOLOGIES AND ELECTROCHEMICAL SYNTHESIS

Clean, renewable energy has enormous implications for the future prosperity of humankind. As a thriving civilisation, living better and longer has been our instinctive pursuit, and advanced biomedical technology is therefore always highly demanded. Research in our lab addresses these problems by using electrochemical technology, nanotechnology and biotechnology. Our research areas include solar water splitting, CO<sub>2</sub> reduction, fuel cells, ammonia synthesis, gas sensors, and proton batteries.

**It would be great to work with Honours students on the following projects:**


### (a). Solar Hydrogen Fuel Production From Seawater

Production of hydrogen fuels from water using electricity generated from renewable energy sources such as solar and wind can provide a sustainable and clean fuel supply for human use. Conventional water splitting is typically carried out in freshwater containing an added supporting electrolyte to conduct electricity, such as potassium hydroxide. However, freshwater only represents a microcosm of the total forms of water found on Earth. The vast majority of water on Earth is seawater (approximately 97%), which contains naturally present salts, predominately sodium chloride. Current hurdles in seawater electrolysis lies in the release of toxic chlorine gas due to the kinetically favoured chlorine evolution over oxygen evolution. The project will develop novel electrodes made of Earth-abundant materials and a prototype water splitting cell for hydrogen production directly from seawater without chlorine evolution.



### (b) Electrocatalytic Synthesis of Ammonia from Renewable Hydrogen and Atmospheric Nitrogen

Ammonia (NH<sub>3</sub>) is one of the most important and widely produced chemicals worldwide for fertiliser production and is also a promising liquid hydrogen carrier to be used as a carbon-free fuel. N<sub>2</sub> has a very strong triple bond and is extremely inert. Currently, the synthesis of NH<sub>3</sub> is still dominated by the high-temperature and high-pressure Haber-Bosch process developed in the early 1900s, which is one of the top largest chemical processes in terms of energy consumption and greenhouse gas emissions.






This project aims to develop a sustainable electrochemical nitrogen reduction reaction (NRR) at ambient conditions powered by renewable energy sources. Our group has recently made breakthrough in developing metal-organic framework (MOF) based catalysts for NRR. In this project, the student will have opportunity to work on these advanced electrocatalysts and evaluated their performance for ammonia synthesis using renewable electricity, hydrogen and atmospheric nitrogen.

### (c). Conversion of CO<sub>2</sub> to Fuels with Renewable Electricity and Earth Abundant Catalysts

Fossil fuels have historically been the primary feedstock for petroleum based products and industrial chemicals. Apart from the impact that fossil fuels pose on the environment, they are generally mined in remote locations and require massive infrastructure for processing and distribution before they are even refined. One promising solution is to reduce CO<sub>2</sub> itself to petrochemical feedstock, which could cater to the unprecedented consumerism of society and simultaneously reduce the anthropogenic emissions of CO<sub>2</sub> in the atmosphere to restore the natural carbon cycle. To improve the CO<sub>2</sub> reduction efficiency, advanced catalysts that are efficient, selective, stable, and low cost need to be developed. This project will design a class of inexpensive, non-metallic electrocatalysts based on nanoporous graphene. The electrocatalysts will be integrated into a prototype device for converting CO<sub>2</sub> into useful fuels.




### (d) Nonprecious Metal Catalysts for Hydrogen Fuel Cells: Towards Affordable Hydrogen Powered Electric Vehicles

Hydrogen fuel cell powered vehicles have been regarded to be the ultimate solution to the future of transportation, and are particularly attractive for larger (e.g. SUV) and longer-range vehicles. Low-temperature hydrogen fuel cells producing electricity using hydrogen and air, with water as the only by-product offer the advantages of simplicity and zero greenhouse gas emission. However, an affordable low-cost fuel cell with catalysts capable of working at industrial scales is yet to be developed. The primary challenges for this project are to discover low-cost electrocatalysts that are active and stable to replace the benchmark catalysts based on precious metals such as platinum for cathode catalyst for hydrogen fuel cells.



In this project the student would learn how to synthesize mesoporous nonprecious metal catalysts. The student will learn how to assemble, prepare and test a hydrogen fuel cell. The student will also have the opportunity to characterise the nonprecious metal catalyst materials using a range of characterisation techniques (XRD, TEM, XPS), and their electrochemical behaviours in operating hydrogen fuel cells.



# **HONOURS ALTERNATIVES**

## HONOURS AT UNSW CANBERRA

### School of Physical Environmental and Mathematical Sciences

---

Research into chemistry is also conducted within the School of Physical, Environmental and Mathematical Sciences at the Canberra campus of UNSW. Co-located with the Australian Defence Force Academy, UNSW Canberra maintains a diverse research program and is home to many research students. It is straight-forward for current UNSW Bachelor of Science students to transition to the Honours program at UNSW Canberra.

A selection of the chemistry research areas being actively pursued at UNSW Canberra are listed below, organised by research group leader. A range of Honours projects are available within each group, which can be adjusted to fit a student's interest. Group leaders may also be available for collaborative projects with research groups within the School of Chemistry, on a case-by-case basis.

Students with an undergraduate Weighted Average Mark (WAM) of 85 or higher are eligible for an \$8000 Honours scholarship if enrolled at UNSW Canberra.

Students who complete their Honours degree within the School of Chemistry should also consider the possibility of higher degree research at UNSW Canberra, as the research at this campus extends into areas complementary to those actively pursued within the School of Chemistry. PhD scholarships for study at UNSW Canberra are generally available to all UNSW students who achieve First Class Honours.

#### **Bio-inorganic Chemistry — Prof Grant Collins**

The current focus of our work is in the development of new, and very promising, classes of multinuclear ruthenium complexes as antimicrobial agents. The emergence of drug-resistant populations of microorganisms means there is clearly a need for the development of new antimicrobials — but more importantly, there is the need for the development of new *classes* of antimicrobials. Complementing the antimicrobial studies has been research on the toxicity and the biological processing of the ruthenium complexes in eukaryotic cells. Emerging directions include utilising ruthenium complexes more broadly as anti-parasite agents, particularly in the treatment of schistosomiasis.

#### **Supramolecular Chemistry — Dr Anthony Day**

The research interests of the Supramolecular Chemistry group are based in organic chemistry synthetic design and the development of new synthetic techniques – in particular molecular host/guest chemistry. The family of molecular hosts known as cucurbit[n]uril and derivatives of this family are a particular target. Specific areas of research interest involve drug delivery vehicles, sensors and supramolecular materials. The group has a secondary research area involving energetic materials, including insensitive munitions, detection, deactivation and environmental aspects. Applied aspects of this work with the Australian Defence Force and the Australian Federal Police are supported by strong links within the research team.

#### **Theoretical Chemistry — Dr Terry Frankcombe**

The group's research activities fall broadly under the umbrella of theoretical and computational chemistry, extending into chemical physics. Four main research themes are currently being pursued: Gaussian-based quantum dynamics methods, gas-surface dynamics, dielectric materials, and the structure and mechanism of photosystem II. Secondary areas of investigation are processes occurring on the surfaces

of spacecraft, simulating condensed matter spectroscopies, and “divide-and-conquer” strategies. Our research can be computationally intensive and involves the combination of chemistry, physics, mathematics and computer science.

Beyond these focus areas the group maintains active expertise in quantum chemistry in general, in molecular, condensed phase and surface adsorbate contexts.

#### **Optical and Laser Spectroscopy of the Solid State — Prof Hans Riesen**

The research of the Riesen group has been mostly focused on laser spectroscopy of transition metal ion doped insulators/wide band gap semiconductors. We are particularly interested in light-induced changes in the solid state that have potential applications in ultra-high density optical data storage and optical signal processing. Very recently Hans and some of his team have studied the generation of slow and fast light by transient hole-burning.

With the advent of the Australian Synchrotron, Hans has also become a user and strong supporter of synchrotron science. In recent years, the group have discovered a novel X-ray storage technology with applications in personal and clinical dosimetry, and medical imaging.

#### **Inorganic Chemistry and Electrochemistry — Dr Lynne Wallace**

The Wallace group research interests include the synthesis and study of redox-active and luminescent transition metal complexes, and applications of such complexes in sensor systems, light-activated molecular devices, supramolecular assemblies and therapeutic approaches. Electrosynthesis of green energetic materials is also a research focus.

#### **Statistical Mechanics — Assoc Prof Cliff Woodward**

The Woodward group conducts research in the field of statistical mechanical theories of condensed matter and complex fluids.

Some highlights of our research program include density functional theory and simulations of room temperature ionic liquids and polymer mediated interactions in polymer/particle mixtures, including many-body effects close to criticality. We have also used intensive simulations and theory to investigate biological systems, in particular, the riddle of “arginine magic”; the poorly understood mechanism that allows arginine-rich peptides to easily penetrate cell membranes. While most of the theoretical work carried out in the group is not reliant on massively large computational platforms, potential candidates should have a strong background in computational methods, physics or physical chemistry and mathematics.

## **JOINT PROJECTS AND COLLABORATIVE PROJECTS WITH OTHER SCHOOLS ACROSS UNSW**

---

School of Chemistry researchers collaborate broadly with researchers in other Schools, Faculties and Institutes. If you are a Chemistry major or are eligible for Honours and wish to do a project aligned between Chemistry and another discipline, please contact the Honours coordinator.