

Never Stand Still

Faculty of Science

School of Mathematics and Statistics

Solution Sheet 6, June 4, 2012

Answers

- 1. 6
- 2. only c and d are always true.
- 3. x = 170, y = 13 and x = 170, y = 3
- 4. $5\sqrt{2}$
- 5. Notice that using only the rules 1 and 2 ((2x, y) and (x, 2y) resp.) we can obtain all points of the form $(2^n, 2^m)$ and $gcd(2^n, 2^m) = 2^{|m-n|}$: a power of 2. Furthermore, the operations (x y, y) and (x, y x) (as used in Euclid's algorithm), preserve the gcd. Hence points with a gcd that is not a power of 2 cannot be reached.

Conversely, these are the only points that can be reached. If $gcd(a,b) = 2^m$, then $a = 2^m a', b = 2^n b'$ with gcd(a',b') = 1. The point (a,b) can be reached from (a',b') using rules 1 and 2 (apply each *m* and *n* times resp.).

Assume a' < b'. Since both a', b' are odd. a' + b' is even, and can be reached from the point $(a', \frac{a'+b'}{2})$. Notice that this point is closer to (1, 1) than (a', b') was.

Continue this process until a' = b', since gcd(a', b') = 1, this point is (1, 1).